arXiv:math-ph/0210012v3 12 Oct 2002

Tau Functions and Matrix Integrals

A. Yu. Orlov*

Abstract

We consider solvable matrix models. We generalize Harish-Chandra-Itzykson-Zuber and certain other integrals (Gross-Witten integral and integrals over complex matrices) using the notion of tau function of matrix argument. In this case one can reduce matrix integral to the integral over eigenvalues, which in turn is certain tau function. The resulting tau functions may be analyzed either by the method of orthogonal polynomials or by Schur functions expansion method.

Contents

1 Introduction **2**

2 Symmetric functions **4**
2.1 Schur functions and scalar product [4] 4
2.2 A deformation of the standard scalar product and series of hypergeometric type 7

3 Solitons, free fermions and fermionic realization of the scalar product **9**
3.1 Fermionic operators, Fock spaces, vacuum expectations, tau functions [7] 9
3.2 Partitions and free fermions 12
3.3 Vacuum expectation value as a scalar product. Symmetric function theory consideration. 13
3.4 KP tau-function $\tau_r(n, \mathbf{t}, \mathbf{t}^*)$ 15
3.5 Algebra of ΨDO on a circle and a development of $e^{A(\mathbf{t}^*)}, e^{\tilde{A}(\mathbf{t})}$ 16
3.6 Schur function expansion of tau function. Loop scalar products 18

4 Angle integration of tau functions of matrix argument. Integration over complex matrices **19**
4.1 Useful formulae 19
4.2 Certain matrix integrals and expansion in Schur functions 20
4.3 Tau functions of a matrix argument and angle integration. The integration of tau functions over complex matrices 23

5 Integral representation of the scalar product and the integrals over eigenvalues of random matrices **25**
5.1 Integral representation of the scalar product $<,>_{r,n}$ 25
5.2 Normal matrix model 28
5.3 Two-matrix model 29
5.4 Hermitian one matrix model 31

---

*Oceanology Institute, Nahimovskii prospect 36, Moscow, Russia, email address: orlovs@wave.sio.rssi.ru

5.5 Unitary two-matrix model 33
5.6 New multi-matrix models which can be solved by the method of orthogonal polynomials and the Schur function expansion 33
5.7 Standard scalar product 35
5.8 About spectral determinants 35

6 Appendices A **36**
6.1 Baker-Akhiezer functions and bilinear identities [7] 36
6.2 $H_0(\mathbf{T})$, fermions $\psi(T,z), \psi^*(T,z)$, bosonization rules [12] 36
6.3 Toda lattice tau-function $\tau(n,\mathbf{t},T,\mathbf{t}^*)$ [12] 38
6.4 Expansion of e^{H_0} and its trace in the Fock subspace F_n 39
6.5 Expressions and linear equations for tau-functions I (Hirota-Miwa variables) . . 40
6.6 The vertex operator action. Linear equations for tau-functions II 42
6.7 Determinant formulae I (Hirota-Miwa variables) [12] 43
6.8 Determinant formulae II (The case $\mathbf{r}(\mathbf{0}) = \mathbf{0}$.) 44
6.9 Integral representations I (Hirota-Miwa variables) [12] 44
6.10 Integral representations II 46
6.11 Rational r and their q-deformations 46
6.12 Examples of hypergeometric series [12] 48
6.13 Baker-Akhiezer functions and the elements of Sato Grassmannian related to the tau function of hypergeometric type 53
6.14 Gauss factorization problem, additional symmetries, string equations and ΨDO on the circle [12] 54

7 Appendices B **56**
7.1 A scalar product in the theory of symmetric functions [4] 56
7.2 A deformation of the scalar product and series of hypergeometric type 57
7.3 Scalar product of tau functions. A conjecture. 58
7.4 Integral representation of scalar products 59

8 Appendix C. Hypergeometric functions **60**

9 Further generalization. Examples of Gelfand-Graev hypergeometric functions **63**
9.1 Scalar product and series in plane partitions 63
9.2 Integral representation for $\tau_{\tilde{\mathbf{r}},\mathbf{r}}$ 64
9.3 Multi-matrix integrals 65
9.4 Fermionic representation [12] 65
9.5 Loop matrix integrals 68
9.6 The example of Gelfand,Graev and Retakh hypergeometric series [12] 68

10 Acknowledgements **71**

1 Introduction

This paper in its main part is based on the previous paper "Soliton theory, Symmetric Functions and Matrix Integrals" [1] and the papers [2],[3]. In the first quoted paper we considered certain class of tau-functions which we call tau-functions of hypergeometric type. The key point of the paper was to identify scalar products on the space of symmetric functions with vacuum expectations of fermionic fields. We considered some known matrix models, hypergeometric functions of matrix argument, Schur function expansion as the perturbation series for matrix integrals.

We enlarge the class of matrix integrals which turned out to be tau functions. (This presentation in particular means that one has a determinant representation, Schur function expansion for perturbation series and the possibility to apply the orthogonal polynomials method for non-perturbative analysis.)

Here we include a version of the previous paper [1] into the text to make it more self consistent.

Symmetric functions. Symmetric function is a function of variables $\mathbf{x} = x_1, \dots, x_n$ which is invariant under the action of permutation group S_n. Symmetric polynomials of n variables form a ring denoted by Λ_n. The number n is usually irrelevant and may be infinite. The ring of symmetric functions in infinitely many variables is denoted by Λ.

Power sums are defined as

$$p_m = \sum_{i=1} x_i^m, \quad m = 1, 2, \dots \tag{1.0.1}$$

Functions $p_{m_1} p_{m_2} \cdots p_{m_k}, \quad k = 1, 2, \dots$ form a basis in Λ.

There are different well investigated polynomials of many variables such as Schur functions, projective Schur functions, complete symmetric functions [4]. Each set form a basis in Λ.

There exists a scalar product in Λ such that Schur functions are orthonormal functions:

$$< s_\lambda, s_\mu >= \delta_{\lambda,\mu} \tag{1.0.2}$$

The notion of scalar product is very important in the theory of symmetric functions. The choice of scalar product gives different orthogonal and bi-orthogonal systems of polynomials.

Soliton theory. KP hierarchy of integrable equations, which is the most popular example, consists of semi-infinite set of nonlinear partial differential equations

$$\partial_{t_m} u = K_m[u], \quad m = 1, 2, \dots \tag{1.0.3}$$

which are commuting flows:

$$\left[\partial_{t_k}, \partial_{t_m}\right] u = 0 \tag{1.0.4}$$

The first nontrivial one is Kadomtsev-Petviashvili equation

$$\partial_{t_3} u = \frac{1}{4} \partial_{t_1}^3 u + \frac{3}{4} \partial_{t_1}^{-1} \partial_{t_2}^2 u + \frac{3}{4} \partial_{t_1} u^2 \tag{1.0.5}$$

which originally served in plasma physics [17]. This equation, which was integrated in [21],[6], now plays a very important role both, in physics (see [18]; see review in [19] for modern applications) and in mathematics. Each members of the KP hierarchy is compatible with each other one.

Another very important equation is the equation of two-dimensional Toda lattice (TL) [22],[7],[27]

$$\partial_{t_1} \partial_{t_1^*} \phi_n = e^{\phi_{n-1} - \phi_n} - e^{\phi_n - \phi_{n+1}} \tag{1.0.6}$$

This equation gives rise to TL hierarchy which contains derivatives with respect to higher times $t_1, t_2, \dots$ and $t_1^*, t_2^*, \dots$.

The key point of soliton theory is the notion of tau function, introduced by Sato (for KP tau-function see [7]). Tau function is a sort of potential which gives rise both to TL hierarchy and KP hierarchy. It depends on two semi-infinite sets of higher times $t_1, t_2, \ldots$ and $t_1^*, t_2^*, \ldots$, and discrete variable n: $\tau = \tau(n, \mathbf{t}, \mathbf{t}^*)$. More explicitly we have [7],[23, 20],[27]:

$$u = 2\partial_{t_1}^2 \log \tau(n, \mathbf{t}, \mathbf{t}^*), \quad \phi_n(\mathbf{t}, \mathbf{t}^*) = -\log \frac{\tau(n+1, \mathbf{t}, \mathbf{t}^*)}{\tau(n, \mathbf{t}, \mathbf{t}^*)} \tag{1.0.7}$$

In soliton theory so-called Hirota-Miwa variables $\mathbf{x}, \mathbf{y}$, which are related to higher times as

$$mt_m = \sum_i x_i^m, \quad mt_m^* = \sum_i y_i^m, \tag{1.0.8}$$

are well-known [24]. Tau function is a symmetric function in Hirota-Miwa variables, higher times t_m, t_m^* are basically power sums, see (1.0.1), see [42] as an example.

It is known fact that typical tau function may be presented in the form of double series in Schur functions [28, 29, 30]:

$$\tau(n, \mathbf{t}, \mathbf{t}^*) = \sum_{\lambda,\mu} K_{\lambda\mu} s_\lambda(\mathbf{t}) s_\mu(\mathbf{t}^*) \tag{1.0.9}$$

where coefficients $K_{\lambda\mu}$ are independent of higher times and solve very special bilinear equations, equivalent to Hirota bilinear equations.

In the previous paper we use these equations to construct hypergeometric functions which depend on many variables, these variables are KP and Toda lattice higher times. Here we shall use the general approach to integrable hierarchies of Kyoto school [7], see also [23, 20]. Especially a set of papers about Toda lattice [27, 28, 29, 30, 31, 32, 33] is important for us. Let us notice that there is an interesting set of papers of M.Adler and P. van Moerbeke, where links between solitons and random matrix theory were worked out from a different point of view (see [13],[14] and other papers).

Random matrices. In many problems in physics and mathematics one use probability distribution on different sets of matrices. We are interested mainly in two matrix models. The integration measure $dM_{1,2}$ is different for different ensembles

$$Z = \int e^{V_1(M_1)+V_2(M_2)+U(M_1,M_2)} dM_1 dM_2 \tag{1.0.10}$$

where

$$V_1(M_1) = \mathrm{Tr} \sum_{m=1}^{\infty} M_1^m t_m, \quad V_2(M_2) = \mathrm{Tr} \sum_{m=1}^{\infty} M_2^m t_m^* \tag{1.0.11}$$

while U may be different for different ensembles.

The crucial point when evaluating the integral (1.0.10) is the possibility to reduce it to the integral over eigenvalues of matrices M_1, M_2. Then (1.0.10) takes the form

$$Z = C \int e^{\sum_{i=1}^n \sum_{m=1}^\infty \left(x_i^m t_m + y_i^m t_m^*\right)} e^{U(\mathbf{x},\mathbf{y})} \prod_{i=1}^n dx_i dy_i \tag{1.0.12}$$

In the present paper we are interested in the perturbation series of different matrix models which are series in parameters t_m, t_m^*.

2 Symmetric functions

2.1 Schur functions and scalar product [4]

Partitions. Polynomial functions of many variables are parameterized by partitions. A *partition* is any (finite or infinite) sequence of non-negative integers in decreasing order:

$$\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_r, \ldots), \quad \lambda_1 \geq \lambda_2 \geq \ldots \geq \lambda_r \geq \ldots \tag{2.1.1}$$

Non-zero λ_i in (2.1.1) are called the *parts* of λ. The number of parts is the *length* of λ, denoted by $l(\lambda)$. The sum of the parts is the *weight* of λ, denoted by $|\lambda|$. If $n = |\lambda|$ we say that λ is the *partition of* n. The partition of zero is denoted by 0.

The *diagram* of a partition (or *Young diagram*) may be defined as the set of points (or nodes) $(i,j) \in Z^2$ such that $1 \leq j \leq \lambda_i$. Thus Young diagram is viewed as a subset of entries in a matrix with $l(\lambda)$ lines and λ_1 rows. We shall denote the diagram of λ by the same symbol λ.

For example



is the diagram of $(3,3,1)$. The weight of this partition is 7, the length is equal to 3.

The *conjugate* of a partition λ is the partition λ' whose diagram is the transpose of the diagram λ, i.e. the diagram obtained by reflection in the main diagonal. The partition $(3,2,2)$ is conjugated to $(3,3,1)$, the corresponding diagram is



There are different notations for partitions. For instance sometimes it is convenient to indicate the number of times each integer occurs as a part:

$$\lambda = (1^{m_1} 2^{m_2} \ldots r^{m_r} \ldots) \tag{2.1.2}$$

means that exactly m_i of the parts of λ are equal to i. For instance partition $(3,3,1)$ is denoted by $(1^1 3^2)$.

Another notation is due to Frobenius. Suppose that the main diagonal of the diagram of λ consists of r nodes $(i,i) \quad (1 \leq i \leq r)$. Let $\alpha_i = \lambda_i - i$ be the number of nodes in the ith row of λ to the right of (i,i), for $1 \leq i \leq r$, and let $\beta_i = \lambda'_i - i$ be the number of nodes in the ith column of λ below (i,i), for $1 \leq i \leq r$. We have $\alpha_1 > \alpha_2 > \cdots > \alpha_r \geq 0$ and $\beta_1 > \beta_2 > \cdots > \beta_r \geq 0$. Then we denote the partition λ by

$$\lambda = (\alpha_1, \ldots, \alpha_r | \beta_1, \ldots, \beta_r) = (\alpha|\beta) \tag{2.1.3}$$

One may say that Frobenius notation corresponds to hook decomposition of diagram of λ, where the biggest hook is $(\alpha_1|\beta_1)$, then $(\alpha_2|\beta_2)$, and so on up to the smallest one which is $(\alpha_r|\beta_r)$. The corners of the hooks are situated on the main diagonal of the diagram. For instance the partition $(3,3,1)$ consists of two hooks $(2,2)$ and $(0,1)$:



and



In Frobenius notation this is $(2,0|2,1)$.

Schur functions, complete symmetric functions and power sums. We consider polynomial symmetric functions of variables $x_i, i = 1,2,\ldots,n$, where the number n is irrelevant and may be infinity. The collection of variables $x_1, x_2, \ldots$ we denote by $\mathbf{x}^N$. When N is irrelevant we denote it by $\mathbf{x}$.

The symmetric polynomials with rational integer coefficients in n variables form a ring which is denoted by Λ_n. A ring of symmetric functions in countably many variables is denoted by Λ (a precise definition of Λ see in [4]).

Let us remind some of well-known symmetric functions.

For each $m \geq 1$ the mth *power sum* is

$$p_m = \sum_i x_i^m \tag{2.1.4}$$

The collection of variables $p_1, p_2, \ldots$ we denote by $\mathbf{p}$.

If we define

$$p_\lambda = p_{\lambda_1} p_{\lambda_2} \cdots \tag{2.1.5}$$

for each partition $\lambda = (\lambda_1, \lambda_2, \ldots)$, then the p_λ form a basic of symmetric polynomial functions with rational coefficients.

Having in mind the links with soliton theory it is more convenient to consider the following "power sums":

$$\gamma_m = \frac{1}{m} \sum_i x_i^m \tag{2.1.6}$$

which is the same as Hirota-Miwa change of variables (1.0.8). The vector

$$\gamma = (\gamma_1, \gamma_2, \ldots) \tag{2.1.7}$$

is an analog of the higher times variables in KP theory.

Complete symmetric functions h_n are given by the generating function

$$\prod_{i \geq 1} (1 - x_i z)^{-1} = \sum_{n \geq 1} h_n(\mathbf{x}) z^n \tag{2.1.8}$$

If we define $h_\lambda = h_{\lambda_1} h_{\lambda_2} \cdots$ for any λ, then h_λ form a Z-basis of Λ.

The complete symmetric functions are expressed in terms of power sums with the help of the following generating function:

$$\exp \sum_{m=1}^{\infty} \gamma_m z^m = \sum_{n \geq 1} h_n(\gamma) z^n \tag{2.1.9}$$

Now suppose that n, the number of variables x_i, is finite.

Given partition λ *the Schur function* s_λ is the quotient

$$s_\lambda(\mathbf{x}) = \frac{\det \left(x_i^{\lambda_i + n - j} \right)_{1 \leq i,j \leq n}}{\det \left(x_i^{n-j} \right)_{1 \leq i,j \leq n}} \tag{2.1.10}$$

where denominator is the Vandermond determinant

$$\det \left(x_i^{n-j} \right)_{1 \leq i,j \leq n} = \prod_{1 \leq i < j \leq n} (x_i - x_j) \tag{2.1.11}$$

Both the numerator and denominator are skew-symmetric, therefore Schur function in variables $x_1, \ldots, x_n$ is symmetric.

For zero partition one puts $s_0(\mathbf{x}) = 1$.

The Schur functions $s_\lambda(x_1, \ldots, x_n)$, *where* $l(\lambda) \leq n$, *form a* Z*-basis of* Λ_n.

Each Schur function s_λ can be expressed as a polynomial in the complete symmetric functions h_n:

$$s_\lambda = \det\left(h_{\lambda_i - i + j}\right)_{1 \le i,j \le n} \tag{2.1.12}$$

where $n \ge l(\lambda)$.

Orthogonality. One may define a scalar product on Λ by requiring that for any pair of partitions λ and μ we have

$$< p_\lambda, p_\mu >= \delta_{\lambda\mu} z_\lambda \tag{2.1.13}$$

where $\delta_{\lambda\mu}$ is a Kronecker symbol and

$$z_\lambda = \prod_{i \ge 1} i^{m_i} \cdot m_i! \tag{2.1.14}$$

where $m_i = m_i(\lambda)$ is the number of parts of λ equal to i.

For any pair of partitions λ and μ we also have

$$< s_\lambda, s_\mu >= \delta_{\lambda,\mu} \tag{2.1.15}$$

so that the s_λ form an *orthonormal* basis of Λ. The s_λ such that $|\lambda| = n$ form an orhtonormal basis for all symmetric polynomials of degree n.

2.2 A deformation of the standard scalar product and series of hypergeometric type

Scalar product $<,>_{r,n}$. Let us consider a function r which depends on a single variable n, the n is integer. Given partition λ let us define

$$r_\lambda(x) = \prod_{i,j \in \lambda} r(x + j - i) \tag{2.2.1}$$

Thus $r_\lambda(n)$ is a product of functions r over all nodes of Young diagram of the partition λ where argument of r is defined by entries i, j of a node. The value of $j - i$ is zero on the main diagonal; the value $j - i$ is called the content of the node. For instance, for the partition $(3, 3, 1)$ the diagram is



so our $r_\lambda(x)$ is equal to $r(x+2)(r(x+1))^2(r(x))^2 r(x-1)r(x-2)$.

For zero partition one puts $r_0 \equiv 1$.

Given integer n and a function on the lattice $r(n), n \in Z$, define a scalar product, parameterized by r, n as follows

$$< s_\lambda, s_\mu >_{r,n}= r_\lambda(n)\delta_{\lambda\mu} \tag{2.2.2}$$

where $\delta_{\lambda\mu}$ is Kronecker symbol.

Let $n_i \in Z$ are zeroes of r, and

$$k = \min |n - n_i|. \tag{2.2.3}$$

The product (2.2.2) is non-degenerated on Λ_k. Really, if $k = n - n_i > 0$ due to the definition (2.2.1) the factor $r_\lambda(n)$ never vanish for the partitions which length is no more then k. In this case the Schur functions of k variables $\{s_\lambda(\mathbf{x}^k), l(\lambda) \le k\}$ form a basis on Λ_k. If $n - n_i = -k < 0$

then the factor $r_\lambda(n)$ never vanish for the partitions $\{\lambda : l(\lambda') \le k\}$, where λ' is the conjugated partition, then $\{s_\lambda(\mathbf{x}^k), l(\lambda') \le k\}$ form a basis on Λ_k. If r is non-vanishing function then the scalar product is non-degenerated on Λ_∞.

Example The **case** $\mathbf{r}(\mathbf{k}) = \mathbf{k}$ plays a role in applications to two-matrix models. In this case we have a realization of this product similar to the realization of standard scalar product, see (3.3.2) below. We shall prove that for the symmetric functions of n variables $x_1, \dots, x_n$ we have the following realization

$$< f(\mathbf{x}), g(\mathbf{x}) >_{r,n} = \frac{1}{n!}\Delta(\partial) f(\partial) \cdot \Delta(\mathbf{x}) g(\mathbf{x})|_{\mathbf{x}=0}, \quad \Delta(\partial) = \prod_{1\le i<j\le n}\left(\frac{\partial}{\partial x_i} - \frac{\partial}{\partial x_j}\right) \tag{2.2.4}$$

where $\Delta(\mathbf{x})$ is the Vandermond determinant (2.1.11), $f(\partial)$ is the symmetric function where all x_i are replaced by $\frac{\partial}{\partial x_i}$ and where $r(k) = k$. The proof follows from definition of the Schur functions (2.1.10) and the formula

$$\frac{1}{n!}\det\left(\partial_{x_i}^{\lambda_j+n-j}\right)_{i,j=1,\dots,n} \cdot \det\left({x_i}^{\mu_j+n-j}\right)_{i,j=1,\dots,n}|_{\mathbf{x}=0} = \delta_{\lambda\mu}\prod_{1\le i<j\le n}(\lambda_i+n-i)! =< s_\lambda, s_\mu >_{r,n} \tag{2.2.5}$$

It is interesting to compare the scalar product (2.2.4) with that considered in [67].□

Let us generalize the example (2.2.4)

A representation of the scalar product if $\mathbf{r}(\mathbf{0}) = \mathbf{0}$. Let us take

$$\hat{x} = \frac{1}{x} r(D), \quad D = x\frac{d}{dx} \tag{2.2.6}$$

Then

$$\hat{x}^m \cdot x^n|_{\mathbf{x}=0} = \delta_{mn} r(1) r(2) \cdots r(n) \tag{2.2.7}$$

Proposition 1 *We have the following representation of scalar product*

$$< f(\mathbf{x}), g(\mathbf{x}) >_{r,n} = \frac{1}{n!}\Delta(\hat{x}) f(\hat{x}) \cdot \Delta(\mathbf{x}) g(\mathbf{x})|_{\mathbf{x}=0}, \quad \Delta(\hat{x}) = \prod_{1\le i<j\le n}(\hat{x}_i - \hat{x}_j) \tag{2.2.8}$$

The proof follows from

$$\frac{1}{n!}\det\left({\hat{x}_i}^{\lambda_j+n-j}\right)_{i,j=1,\dots,n} \cdot \det\left({x_i}^{\mu_j+n-j}\right)_{i,j=1,\dots,n}|_{\mathbf{x}=0} = \delta_{\lambda\mu} r_\lambda(n) =< s_\lambda, s_\mu >_{r,n} \tag{2.2.9}$$

A useful formula. First let us remind the so-called Caughy-Littlewood formulae [4]:

$$\prod_{i,j}(1 - x_i y_j)^{-1} = \sum_\lambda s_\lambda(\mathbf{x}) s_\lambda(\mathbf{y}) \tag{2.2.10}$$

where the number of x, y variables is irrelevant, and the Schur functions are defined via the formula (2.1.10).

If $\gamma = (\gamma_1, \gamma_2, \dots)$ and $\gamma^* = (\gamma_1^*, \gamma_2^*, \dots)$ be two independent semi-infinite set of variables. Then we have *the following useful version of Cauchy-Littlewood formula*:

$$\exp\sum_{m=1}^{\infty} m\gamma_m\gamma_m^* = \sum_\lambda s_\lambda(\gamma) s_\lambda(\gamma^*) \tag{2.2.11}$$

where the sum is going over all partitions including zero partition and s_λ are constructed via (2.1.12) and (2.1.9). Since I did not meet the relation (2.2.11) in the literature I need to prove it (see (3.2.1) below). This proof follows from bosonic (l.h.s) and from fermionic (r.h.s) representations of the vacuum TL function.

Series of hypergeometric type. Let us consider the following pair of functions of variables p_m

$$\exp\sum_{m=1}^{\infty} m\gamma_m t_m, \quad \exp\sum_{m=1}^{\infty} m\gamma_m t_m^* \tag{2.2.12}$$

where t_m and t_m^* are formal parameters. Considering p_m as power sums and using formulae (2.2.11) and (2.2.2), we evaluate the scalar product of functions (2.2.12):

$$< \exp\sum_{m=1}^{\infty} m\gamma_m t_m, \exp\sum_{m=1}^{\infty} m\gamma_m t_m^* >_{r,n} = \sum_{l(\lambda)\le k} r_\lambda(n) s_\lambda(\mathbf{t}) s_\lambda(\mathbf{t}^*) = \sum_\lambda r_\lambda(n) s_\lambda(\mathbf{t}) s_\lambda(\mathbf{t}^*) \tag{2.2.13}$$

where sum is going over all partitions λ of the length $l(\lambda) \le k$, see (2.2.3), and including zero. The Schur functions $s_\lambda(\mathbf{t}), s_\lambda(\mathbf{t}^*)$ are defined with the help of

$$s_\lambda(\mathbf{t}) = \det h_{\lambda_i - i + j}(\mathbf{t})_{1\le i,j\le l(\lambda)}, \quad \exp\sum_{m=1}^{\infty} z^m t_m = \sum_{k=1}^{\infty} z^k h_k(\mathbf{t}) \tag{2.2.14}$$

We call the series (2.2.13) *the series of hypergeometric type*. It is not necessarily convergent series. If one chooses r as rational function he obtains so-called hypergeometric functions of matrix argument, or, in other words, hypergeometric functions related to $GL(N,C)/U(N)$ zonal spherical functions [39],[26]. If one takes trigonometric function r we gets q-deformed versions of these functions, suggested by I.G.Macdonald and studied by S.Milne in [38].

In case $\mathbf{r(0) = 0}$, which we shall show is important in applications to models of random matrices, we obtain

$$\tau_r(n,\mathbf{t},\mathbf{t}^*) = e^{\sum_{m=0}^{\infty}\sum_{i=1}^{n} t_m \hat{x}_i^m}\Delta(\hat{x}) \cdot e^{\sum_{m=0}^{\infty}\sum_{i=1}^{n} t_m^* x_i^m}\Delta(x)|_{\mathbf{x}=0}, \quad \hat{x}_i = \frac{1}{x_i} r\left(x_i \frac{\partial}{\partial x_i}\right) \tag{2.2.15}$$

Scalar product of series of hypergeometric type. Considering the scalar product of functions of γ_m and using formulae (2.2.13) and (2.2.2), we evaluate the scalar product of series of hypergeometric type (2.2.13), which are related to the different choice of the function r (which we denote by r_1 and r_2; don't miss it with the notation r_λ of (2.2.1), labelled by a partition!) . We find that the scalar product is a series of hypergeometric type again:

$$< \tau_{r_1}(k,\mathbf{t},\gamma), \tau_{r_2}(m,\gamma,\mathbf{t}^*) >_{r,n} = \tau_{r_3}(0,\mathbf{t},\mathbf{t}^*) \tag{2.2.16}$$

where

$$r_3(i) = r_1(k+i) r(n+i) r_2(m+i) \tag{2.2.17}$$

3 Solitons, free fermions and fermionic realization of the scalar product

In this section we use the language of free fermions as it was suggested in [23],[7].

3.1 Fermionic operators, Fock spaces, vacuum expectations, tau functions [7]

Free fermions. We have the algebra of free fermions *the Clifford algebra* **A** over **C** with generators $\psi_n, \psi_n^* (n \in Z)$, satisfying the defining relations:

$$[\psi_m, \psi_n]_+ = [\psi_m^*, \psi_n^*]_+ = 0; \qquad [\psi_m, \psi_n^*]_+ = \delta_{mn}. \tag{3.1.1}$$

Any element of $W = (\oplus_{m\in Z}\mathbf{C}\psi_m) \oplus (\oplus_{m\in Z}\mathbf{C}\psi_m^*)$ will be referred as a *free fermion.* The Clifford algebra has a standard representation (*Fock representation*) as follows. Put $W_{an} = (\oplus_{m<0}\mathbf{C}\psi_m) \oplus (\oplus_{m\geq 0}\mathbf{C}\psi_m^*)$, and $W_{cr} = (\oplus_{m\geq 0}\mathbf{C}\psi_m) \oplus (\oplus_{m<0}\mathbf{C}\psi_m^*)$, and consider left (resp. right) $\mathbf{A}$-module $F = \mathbf{A}/\mathbf{A}W_{an}$ (resp $F^* = W_{cr}\mathbf{A}\backslash\mathbf{A}$). These are cyclic $\mathbf{A}$-modules generated by the vectors $|0\rangle = 1 \bmod \mathbf{A}W_{an}$ (resp. by $\langle 0| = 1 \bmod W_{cr}\mathbf{A}$), with the properties

$$\psi_m|0\rangle = 0 \qquad (m<0), \qquad \psi_m^*|0\rangle = 0 \qquad (m\geq 0), \tag{3.1.2}$$

$$\langle 0|\psi_m = 0 \qquad (m\geq 0), \qquad \langle 0|\psi_m^* = 0 \qquad (m<0). \tag{3.1.3}$$

Vectors $\langle 0|$ and $|0\rangle$ are refereed as left and right vacuum vectors. Fermions $w \in W_{an}$ eliminate left vacuum vector, while fermions $w \in W_{cr}$ eliminate right vacuum vector.

The Fock spaces F and F^* are dual ones, the pairing is defined with the help of a linear form $\langle 0||0\rangle$ on $\mathbf{A}$ called the *vacuum expectation value.* It is given by

$$\langle 0|1|0\rangle = 1, \quad \langle 0|\psi_m\psi_m^*|0\rangle = 1 \quad m<0, \quad \langle 0|\psi_m^*\psi_m|0\rangle = 1 \quad m\geq 0, \tag{3.1.4}$$

$$\langle 0|\psi_m\psi_n|0\rangle = \langle 0|\psi_m^*\psi_n^*|0\rangle = 0, \quad \langle 0|\psi_m\psi_n^*|0\rangle = 0 \quad m\neq n. \tag{3.1.5}$$

and by **the Wick rule**, which is

$$\langle 0|w_1\cdots w_{2n+1}|0\rangle = 0, \quad \langle 0|w_1\cdots w_{2n}|0\rangle = \sum_\sigma sgn\sigma\langle 0|w_{\sigma(1)}w_{\sigma(2)}|0\rangle\cdots\langle 0|w_{\sigma(2n-1)}w_{\sigma(2n)}|0\rangle, \tag{3.1.6}$$

where $w_k \in W$, and σ runs over permutations such that $\sigma(1) < \sigma(2), \ldots, \sigma(2n-1) < \sigma(2n)$ and $\sigma(1) < \sigma(3) < \cdots < \sigma(2n-1)$.

Lie algebra $\widehat{gl}(\infty)$. Consider infinite matrices $(a_{ij})_{i,j\in Z}$ satisfying the condition: there exists an N such that $a_{ij} = 0$ for $|i-j| > N$; these matrices are called generalized Jacobian matrices. The generalized Jacobian matrices form Lie algebra, Lie bracket being the usual commutator of matrices $[A,B] = AB - BA$.

Let us consider a set of linear combinations of quadratic elements $\sum a_{ij} : \psi_i\psi_j^* :$, where the notation :: means *the normal ordering* which is defined as $: \psi_i\psi_j^* := \psi_i\psi_j^* - \langle 0|\psi_i\psi_j^*|0\rangle$. These elements together with 1 span an infinite dimensional Lie algebra $\widehat{gl}(\infty)$:

$$[\sum a_{ij} : \psi_i\psi_j^* :, \sum b_{ij} : \psi_i\psi_j^* :] = \sum c_{ij} : \psi_i\psi_j^* : + c_0, \tag{3.1.7}$$

$$c_{ij} = \sum_k a_{ik}b_{kj} - \sum_k b_{ik}a_{kj}, \tag{3.1.8}$$

and the last term

$$c_0 = \sum_{i<0,j\geq 0} a_{ij}b_{ji} - \sum_{i\geq 0,j<0} a_{ij}b_{ji}. \tag{3.1.9}$$

commutes with each quadratic term.

We see that Lie algebra of quadratic elements $\sum a_{ij} : \psi_i\psi_j^* :$ is different of the algebra of the generalized Jacobian matrices; the difference is the *central extension* c_0.

Bilinear identity. Now we define the operator g which is an element of the group corresponding to the Lie algebra $\widehat{gl}(\infty)$:

$$g = \exp\sum a_{nm} : \psi_n\psi_m : \tag{3.1.10}$$

Using (3.1.7) it is possible to derive the following relation

$$g\psi_n = \sum_m \psi_m A_{mn} g, \qquad \psi_n^* g = g\sum_m A_{nm}\psi_m^*. \tag{3.1.11}$$

where coefficients A_{nm} are defined by a_{nm}. In turn (3.1.11) yields

$$[\sum_{n\in Z}\psi_n\otimes\psi_n^*, g\otimes g]=0 \tag{3.1.12}$$

The last relation is very important for applications in soliton theory, and it is equivalent to the so-called Hirota equations.

Higher times. Let us introduce

$$H_n=\sum_{k=-\infty}^{\infty}\psi_k\psi_{k+n}^*,\quad n\neq 0,\quad H(\mathbf{t})=\sum_{n=1}^{+\infty}t_nH_n,\quad H^*(\mathbf{t}^*)=\sum_{n=1}^{+\infty}t_n^*H_{-n}. \tag{3.1.13}$$

$H_n\in\widehat{gl}(\infty)$, and $H(\mathbf{t}),H^*(\mathbf{t}^*)$ belong to $\widehat{gl}(\infty)$ if one restricts the number of non-vanishing parameters t_m,t_m^*. For H_n we have Heisenberg algebra commutation relations:

$$[H_n,H_m]=n\delta_{m+n,0}. \tag{3.1.14}$$

Let us mark that

$$H_n|0\rangle=0=\langle 0|H_{-n},\quad n>0 \tag{3.1.15}$$

For future purposes we introduce the following fermions

$$\psi(z)=\sum_k\psi_kz^k,\qquad \psi^*(z)=\sum_k\psi_k^*z^{-k-1}dz \tag{3.1.16}$$

Using (3.1.1) and (3.1.13) one obtains

$$e^{H(\mathbf{t})}\psi(z)e^{-H(\mathbf{t})}=\psi(z)e^{\xi(\mathbf{t},z)},\quad e^{H(\mathbf{t})}\psi^*(z)e^{-H(\mathbf{t})}=\psi^*(z)e^{-\xi(\mathbf{t},z)}, \tag{3.1.17}$$

$$e^{-H^*(\mathbf{t}^*)}\psi(z)e^{H^*(\mathbf{t}^*)}=\psi(z)e^{-\xi(\mathbf{t}^*,z^{-1})},\quad e^{-H^*(\mathbf{t}^*)}\psi^*(z)e^{H^*(\mathbf{t}^*)}=\psi^*(z)e^{\xi(\mathbf{t}^*,z^{-1})}. \tag{3.1.18}$$

Using

$$\sum_{n\in Z}\psi_n\otimes\psi_n^*=\mathtt{res}_{z=0}\psi(z)\otimes\psi^*(z) \tag{3.1.19}$$

and formulae (3.1.17),(3.1.18) result in

$$[\sum_{n\in Z}\psi_n\otimes\psi_n^*,e^{H(\mathbf{t})}\otimes e^{H(\mathbf{t})}]=0,\quad [\sum_{n\in Z}\psi_n\otimes\psi_n^*,e^{H^*(\mathbf{t}^*)}\otimes e^{H^*(\mathbf{t}^*)}]=0 \tag{3.1.20}$$

Therefore if g solves (3.1.12) then $e^{H(\mathbf{t})}ge^{H^*(\mathbf{t}^*)}$ also solves (3.1.12):

$$\left[\mathtt{res}_{z=0}\psi(z)\otimes\psi^*(z),e^{H(\mathbf{t})}ge^{H^*(\mathbf{t}^*)}\otimes e^{H(\mathbf{t})}ge^{H^*(\mathbf{t}^*)}\right]=0 \tag{3.1.21}$$

The KP and TL tau functions First let us define vacuum vectors labelled by the integer M:

$$\langle n|=\langle 0|\Psi_n^*,\qquad |n\rangle=\Psi_n|0\rangle, \tag{3.1.22}$$

$$\begin{aligned}&\Psi_n=\psi_{n-1}\cdots\psi_1\psi_0\quad n>0,\qquad \Psi_n=\psi_n^*\cdots\psi_{-2}^*\psi_{-1}^*\quad n<0,\\ &\Psi_n^*=\psi_0^*\psi_1^*\cdots\psi_{n-1}^*\quad n>0,\qquad \Psi_n^*=\psi_{-1}\psi_{-2}\cdots\psi_n\quad n<0.\end{aligned} \tag{3.1.23}$$

Given g, which satisfy bilinear identity (3.1.12), one constructs KP and TL tau-functions (1.0.7) as follows:

$$\tau_{KP}(n,\mathbf{t})=\langle n|e^{H(\mathbf{t})}g|n\rangle, \tag{3.1.24}$$

$$\tau_{TL}(n,\mathbf{t},\mathbf{t}^*) = \langle n|e^{H(\mathbf{t})} g e^{H^*(\mathbf{t}^*)}|n\rangle. \qquad (3.1.25)$$

If one fixes the variables $n, \mathbf{t}^*$, then τ_{TL} may be recognized as τ_{KP}.

The times $\mathbf{t} = (t_1, t_2, \dots)$ and $\mathbf{t}^* = (t_1^*, t_2^*, \dots)$ are called higher Toda lattice times [23, 27] (the first set $\mathbf{t}$ is in the same time the set of higher KP times. The first times of this set t_1, t_2, t_3 are independent variables for KP equation (1.0.5), which is the first nontrivial equation in the KP hierarchy).

Fermions and Schur functions. In the KP theory it is suitable to use the definition of the Schur function s_λ in terms of higher times

$$s_\lambda(\mathbf{t}) = \det(h_{n_i - i + j}(\mathbf{t}))_{1\le i,j\le r}, \qquad (3.1.26)$$

where $h_m(\mathbf{t})$ is the elementary Schur polynomial defined by the Taylor's expansion:

$$e^{\xi(\mathbf{t},z)} = \exp(\sum_{k=1}^{+\infty} t_k z^k) = \sum_{n=0}^{+\infty} z^n h_n(\mathbf{t}). \qquad (3.1.27)$$

Lemma 1 [7]
For $-j_1 < \cdots < -j_k < 0 \le i_s < \cdots < i_1$, $s-k \ge 0$ the next formula is valid:

$$\langle s-k|e^{H(\mathbf{t})}\psi^*_{-j_1}\cdots\psi^*_{-j_k}\psi_{i_s}\cdots\psi_{i_1}|0\rangle = (-1)^{j_1+\cdots+j_k+(k-s)(k-s+1)/2} s_\lambda(\mathbf{t}), \qquad (3.1.28)$$

where the partition $\lambda = (n_1, \dots, n_{s-k}, n_{s-k+1}, \dots, n_{s-k+j_1})$ is defined by the pair of partitions:

$$(n_1, \dots, n_{s-k}) = (i_1 - (s-k) + 1, i_2 - (s-k) + 2, \dots, i_{s-k}), \qquad (3.1.29)$$

$$(n_{s-k+1}, \dots, n_{s-k+j_1}) = (i_{s-k+1}, \dots, i_s | j_1 - 1, \dots, j_k - 1). \qquad (3.1.30)$$

The proof is achieved by direct calculation. Here $(\dots|\dots)$ is the Frobenius notation for a partition, see (2.1.3).

3.2 Partitions and free fermions

First let us prove the formula (2.2.11). Consider the so-called vacuum TL tau function

$$\langle 0|e^{H(\mathbf{t})}e^{H^*(\mathbf{t}^*)}|0\rangle \qquad (3.2.1)$$

By the Heisenberg algebra commutation relation (3.1.14) and by (3.1.15) it is equal to

$$e^{\sum_m m t_m t_m^*} \qquad (3.2.2)$$

On the other hand one may develop $e^{H^*(\mathbf{t}^*)}$ in Taylor series, then use the explicit form of H^* in terms of free fermions (see (3.1.13), and then use Lemma 1 of previous subsection. He gets

$$\sum_\lambda s_\lambda(\mathbf{t}) s_\lambda(\mathbf{t}^*) \qquad (3.2.3)$$

where sum is going over all partitions including zero partition $\Box$.

Lemma 2 Given partition $\lambda = (i_1, \dots, i_s | j_1 - 1, \dots, j_s - 1)$ let us introduce the notation

$$|\lambda\rangle = (-1)^{j_1+\cdots+j_s}\psi^*_{-j_1}\cdots\psi^*_{-j_s}\psi_{i_s}\cdots\psi_{i_1}|0\rangle \qquad (3.2.4)$$

$$\langle\lambda| = (-1)^{j_1+\cdots+j_s}\langle 0|\psi^*_{i_1}\cdots\psi^*_{i_s}\psi_{-j_s}\cdots\psi_{-j_1} \qquad (3.2.5)$$

Then

$$\langle\lambda|\mu\rangle = \delta_{\lambda,\mu} \tag{3.2.6}$$

where δ is Kronecker symbol, and

$$s_\lambda(\mathbf{H}^*)|0\rangle = |\lambda\rangle \tag{3.2.7}$$

$$\langle 0|s_\lambda(\mathbf{H}) = \langle\lambda| \tag{3.2.8}$$

here s_λ is defined according to (2.2.14) where instead of $\mathbf{t} = (t_1, t_2, \dots)$ we have

$$\mathbf{H}^* = (H_{-1}, \frac{H_{-2}}{2}, \dots, \frac{H_{-m}}{m}, \dots) \tag{3.2.9}$$

$$\mathbf{H} = (H_1, \frac{H_2}{2}, \dots, \frac{H_m}{m}, \dots) \tag{3.2.10}$$

□

The proof of (3.2.7) and of (3.2.8) follows from using Lemma 1 and the development (2.2.11)

$$e^{\sum_{m=1}^\infty H_m t_m} = \sum_\lambda s_\lambda(\mathbf{H}) s_\lambda(\mathbf{t}) \tag{3.2.11}$$

Lemma 3

$$s_\lambda(-\mathbf{A})|0\rangle = r_\lambda(0)|\lambda\rangle \tag{3.2.12}$$

where s_λ is defined according to (2.2.14) where instead of $\mathbf{t} = (t_1, t_2, \dots)$ we have

$$\mathbf{A} = (A_1, \frac{A_2}{2}, \dots, \frac{A_m}{m}, \dots) \tag{3.2.13}$$

$$A_k = \sum_{n=-\infty}^{\infty} \psi^*_{n-k}\psi_n r(n) r(n-1)\cdots r(n-k+1), \quad k = 1, 2, \dots. \tag{3.2.14}$$

□

The proof follows from the following. First the components of vector $\mathbf{A}$ mutually commute: $[A_m, A_k] = 0$. Then we apply the development

$$e^{\sum_{m=1}^\infty A_m t_m} = \sum_\lambda s_\lambda(-\mathbf{A}) s_\lambda(\mathbf{t}) \tag{3.2.15}$$

and apply the both sides to the right vacuum vector:

$$e^{\sum_{m=1}^\infty A_m t_m}|0\rangle = \sum_\lambda s_\lambda(\mathbf{t}) s_\lambda(-\mathbf{A})|0\rangle \tag{3.2.16}$$

Then we develop $e^{\sum_{m=1}^\infty A_m t_m}|0\rangle$ using Taylor expansion of the exponential function and also use the explicit form of (3.2.14), Lemma 1, the definition of $|\lambda\rangle$ and also (3.2.6).

3.3 Vacuum expectation value as a scalar product. Symmetric function theory consideration.

Let us remember the definition of scalar product (2.1.13) on the space of symmetric functions. It is known that one may present it directly in terms of the variables γ and the derivatives with respect to these variables:

$$\tilde{\partial} = (\partial_{\gamma_1}, \frac{1}{2}\partial_{\gamma_2}, \dots, \frac{1}{n}\partial_{\gamma_n}, \dots) \tag{3.3.1}$$

Then it is known, that the scalar product of polynomial functions, say, f and g, may be defined as

$$\langle f, g\rangle = \left(f(\tilde{\partial})\cdot g(\gamma)\right)|_{\gamma=0} \tag{3.3.2}$$

For instance, using (3.3.2), one easily gets

$$\langle \gamma_n, \gamma_m\rangle = \frac{1}{n}\delta_{n,m}, \quad \langle p_n, p_m\rangle = n\delta_{n,m} \tag{3.3.3}$$

which is a particular case of (2.1.13).

The following statement is important

Proposition 2 *Let scalar product is defined as in (3.3.2). Then*

$$\langle f, g\rangle = \langle 0|f(\mathbf{H})g(\mathbf{H}^*)|0\rangle \tag{3.3.4}$$

where

$$\mathbf{H} = \left(H_1, \frac{H_2}{2}, \dots, \frac{H_n}{n}, \dots\right), \quad \mathbf{H}^* = \left(H_{-1}, \frac{H_{-2}}{2}, \dots, \frac{H_{-n}}{n}, \dots\right) \tag{3.3.5}$$

It follows from the fact that higher times and derivatives with respect to higher times gives a realization of Heisenberg algebra $H_kH_m - H_mH_k = k\delta_{k+m,0}$:

$$\partial_{\gamma_k}\cdot m\gamma_m - m\gamma_m\cdot\partial_{\gamma_k} = k\delta_{k+m,0} \tag{3.3.6}$$

and from the comparison of

$$\partial_{\gamma_m}\cdot 1 = 0, \qquad Free \quad term \quad of \quad \partial_{\gamma_m} \quad is \quad equal \quad to \quad 0 \tag{3.3.7}$$

with

$$H_m|n\rangle = 0, m > 0, \qquad \langle n|H_m = 0, m < 0 \tag{3.3.8}$$

respectively.

Now let us consider deformed scalar product (2.2.2)

$$\langle s_\mu, s_\lambda\rangle_{r,n} = r_\lambda(n)\delta_{\mu,\lambda} \tag{3.3.9}$$

Each polynomial function is a linear combination of Schur functions. We have the following realization of the deformed scalar product

Proposition 3

$$\langle f, g\rangle_{r,n} = \langle n|f(\mathbf{H})g(-\mathbf{A})|n\rangle \tag{3.3.10}$$

where

$$\mathbf{H} = \left(H_1, \frac{H_2}{2}, \dots, \frac{H_m}{m}, \dots\right), \quad \mathbf{A} = \left(A_1, \frac{A_2}{2}, \dots, \frac{A_m}{m}, \dots\right) \tag{3.3.11}$$

H_k and A_k are defined respectively by (3.1.13) and (3.2.14).

3.4 KP tau-function $\tau_r(n, \mathbf{t}, \mathbf{t}^*)$

For the collection of independent variables $\mathbf{t}^* = (t_1^*, t_2^*, \dots)$ we denote

$$A(\mathbf{t}^*) = \sum_{m=1}^{\infty} t_m^* A_m. \tag{3.4.1}$$

where A_m were defined above in (3.2.14).

Using the explicit form of A_m (3.2.14) and anti-commutation relations (3.1.1) we obtain

$$e^{A(\mathbf{t}^*)}\psi(z)e^{-A(\mathbf{t}^*)} = e^{-\xi_r(\mathbf{t}^*, z^{-1})} \cdot \psi(z) \tag{3.4.2}$$

$$e^{A(\mathbf{t}^*)}\psi^*(z)e^{-A(\mathbf{t}^*)} = e^{\xi_{r'}(\mathbf{t}^*, z^{-1})} \cdot \psi^*(z) \tag{3.4.3}$$

where operators

$$\xi_r(\mathbf{t}^*, z^{-1}) = \sum_{m=1}^{+\infty} t_m \left(\frac{1}{z} r(D)\right)^m, \quad D = z\frac{d}{dz}, \quad r'(D) = r(-D) \tag{3.4.4}$$

act on all functions of z on the right hand side according to the rule which was described in the beginning of the subsection 3.4, namely $r(D)\cdot z^n = r(n)z^n$. The exponents in (3.4.2),(3.4.3),(3.4.4) are considered as their Taylor series.

Using relations (3.4.2),(3.4.3) and the fact that inside res_z the operator $\frac{1}{z}r(D)$ is the conjugate of $\frac{1}{z}r(-D) = \frac{1}{z}r'(D)$, we get

Lemma The fermionic operator $e^{A(\mathbf{t}^*)}$ solves the bilinear identity (3.1.12):

$$\left[\mathrm{res}_{z=0}\psi(z) \otimes \psi^*(z), e^{A(\mathbf{t}^*)} \otimes e^{A(\mathbf{t}^*)}\right] = 0 \tag{3.4.5}$$

□

Of cause this Lemma is also a result of a general treating in [7].

Now it is natural to consider the following tau function:

$$\tau_r(n, \mathbf{t}, \mathbf{t}^*) := \langle n| e^{H(\mathbf{t})} e^{-A(\mathbf{t}^*)} |n\rangle. \tag{3.4.6}$$

Let us use the fermionic realization of scalar product (3.3.10). We immediately obtain

Proposition 4

$$\tau_r(n, \mathbf{t}, \mathbf{t}^*) = \langle e^{\sum_{m=1}^{\infty} m t_m \gamma_m}, e^{\sum_{m=1}^{\infty} m t_m^* \gamma_m} \rangle_{r,n} \tag{3.4.7}$$

Due to (2.2.13) we get

Proposition 5

$$\tau_r(n, \mathbf{t}, \mathbf{t}^*) = \sum_{\lambda} r_\lambda(n) s_\lambda(\mathbf{t}) s_\lambda(\mathbf{t}^*). \tag{3.4.8}$$

We shall not consider the problem of convergence of this series. The variables $M, \mathbf{t}$ play the role of KP higher times, $\mathbf{t}^*$ is a collection of group times for a commuting subalgebra of additional symmetries of KP (see [35, 34, 13] and Remark 7 in [36]). From different point of view (3.4.8) is a tau-function of two-dimensional Toda lattice [27] with two sets of continuous variables $\mathbf{t}, \mathbf{t}^*$ and one discrete variable M. Formula (3.4.8) is symmetric with respect to $\mathbf{t} \leftrightarrow \mathbf{t}^*$. For given r we define the function r':

$$r'(n) := r(-n). \tag{3.4.9}$$

Tau function τ_r has properties

$$\tau_r(n,\mathbf{t},\mathbf{t}^*)=\tau_r(n,\mathbf{t}^*,\mathbf{t}),\quad \tau_{r'}(-n,-\mathbf{t},-\mathbf{t}^*)=\tau_r(n,\mathbf{t},\mathbf{t}^*). \tag{3.4.10}$$

Also $\tau_r(n,\mathbf{t},\mathbf{t}^*)$ does not change if $t_m \to a^m t_m, t_m^* \to a^{-m}t_m^*, m=1,2,\ldots,$, and it does change if $t_m \to a^m t_m, m=1,2,\ldots, r(n)\to a^{-1}r(n)$. (3.4.10) follows from the relations

$$r'_\lambda(n)=r_{\lambda'}(-n),\quad s_\lambda(\mathbf{t})=(-)^{|\lambda|}s_{\lambda'}(-\mathbf{t}). \tag{3.4.11}$$

Let us notice that tau-function (3.4.8) can be viewed as a result of action of additional symmetries [36],[20],[35] on the vacuum tau-function .

3.5 Algebra of ΨDO on a circle and a development of $e^{A(\mathbf{t}^*)}, e^{\tilde{A}(\mathbf{t})}$

Given functions $r,\tilde{r}$, the operators

$$A_m=-\sum_{n=-\infty}^{\infty} r(n)\cdots r(n-m+1)\psi_n\psi_{n-m}^*,\quad m=1,2,\ldots, \tag{3.5.1}$$

$$\tilde{A}_m=\sum_{n=-\infty}^{\infty}\tilde{r}(n+1)\cdots\tilde{r}(n+m)\psi_n\psi_{n+m}^*,\quad m=1,2,\ldots \tag{3.5.2}$$

belong to the Lie algebra of pseudo-differential operators on a circle with central extension. These operators may be also rewritten in the following form

$$A_m=\frac{1}{2\pi\sqrt{-1}}\oint :\psi^*(z)\left(\frac{1}{z}r(D)\right)^m\cdot\psi(z): \tag{3.5.3}$$

$$\tilde{A}_m=-\frac{1}{2\pi\sqrt{-1}}\oint :\psi^*(z)\left(\tilde{r}(D)z\right)^m\cdot\psi(z): \tag{3.5.4}$$

where $D=z\frac{d}{dz}$, pseudo-differential operators $r(D),\tilde{r}(D)$ act on functions of z to the right. This action is given as follows: $r(D)\cdot z^n=r(n)z^n$, where $z^n, n\in Z$ is the basis of holomorphic functions in the punctured disk $1>|z|>0$. Central extensions of the Lie algebra of generators (3.5.3) (remember the subsection 3.1) are described by the formulae

$$\omega_n(A_m,A_k)=\delta_{mk}\tilde{r}(n+m-1)\cdots\tilde{r}(n)r(n)\cdots r(n-m+1),\quad \omega_n-\omega_{n'}\sim 0 \tag{3.5.5}$$

where n is an integer. These extension which differs by the choice of n are cohomological ones. The choice corresponds to the choice of normal ordering $::$, which may chosen in a different way as $:a: \ := a-\langle n|a|n\rangle$, n is an integer. (We choose it by $n=0$ see subsection 3.1). When functions $r,\tilde{r}$ are polynomials the operators $\left(\frac{1}{z}r(D)\right)^m, (\tilde{r}(D)z)^m$ belong to the W_∞ algebra, while the fermionic operators (3.2.14),(3.5.2) belong to the algebra with central extension denoted by $\widehat{W}_\infty$ [41].

We are interested in calculating the vacuum expectation value of different products of operators of type $e^{\sum A_m t_m^*}$ and $e^{\sum \tilde{A}_m t_m}$.

Lemma [12] Let partitions $\lambda=(i_1,\ldots,i_s|j_1-1,\ldots,j_s-1)$ and $\mu=(\tilde{i}_1,\ldots,\tilde{i}_r|\tilde{j}_1-1,\ldots,\tilde{j}_r-1)$ satisfy the relation $\lambda\supseteq\mu$. The following is valid:

$$\langle 0|\psi_{\tilde{i}_1}^*\cdots\psi_{\tilde{i}_r}^*\psi_{-\tilde{j}_r}\cdots\psi_{-\tilde{j}_1}e^{A(\mathbf{t}^*)}\psi_{-j_1}^*\cdots\psi_{-j_s}^*\psi_{i_s}\cdots\psi_{i_1}|0\rangle = \\ =(-1)^{\tilde{j}_1+\cdots+\tilde{j}_r+j_1+\cdots+j_s}s_{\lambda/\mu}(\mathbf{t}^*)r_{\lambda/\mu}(0) \tag{3.5.6}$$

$$\langle 0|\psi^*_{i_1}\cdots\psi^*_{i_s}\psi_{-j_s}\cdots\psi_{-j_1}e^{\tilde{A}(\mathbf{t})}\psi^*_{-\tilde{j}_1}\cdots\psi^*_{-\tilde{j}_r}\psi_{\tilde{i}_r}\cdots\psi_{\tilde{i}_1}|0\rangle = \\ = (-1)^{\tilde{j}_1+\cdots+\tilde{j}_r+j_1+\cdots+j_s}s_{\lambda/\mu}(\mathbf{t})\tilde{r}_{\lambda/\mu}(0) \qquad (3.5.7)$$

where $s_{\lambda/\mu}(\mathbf{t})$ is a skew Schur function [4]:

$$s_{\lambda/\mu}(\mathbf{t}) = \det\left(h_{\lambda_i-\mu_j-i+j}(\mathbf{t})\right), \qquad (3.5.8)$$

and

$$r_{\lambda/\mu}(n) := \prod_{i,j\in\lambda/\mu} r(n+j-i), \quad \tilde{r}_{\lambda/\mu}(n) := \prod_{i,j\in\lambda/\mu} \tilde{r}(n+j-i) \qquad (3.5.9)$$

□

The proof is achieved by a development of $e^A = 1 + A + \cdots, e^{\tilde{A}} = 1 + \tilde{A} + \cdots$ and the direct evaluation of vacuum expectations (3.5.6),(3.5.7), and a determinant formula for the skew Schur function of Example 22 in Sec 5 of [4].

Let us introduce vectors

$$|\lambda, n\rangle = (-1)^{j_1+\cdots+j_s}\psi^*_{-j_1+n}\cdots\psi^*_{-j_s+n}\psi_{i_s+n}\cdots\psi_{i_1+n}|n\rangle \qquad (3.5.10)$$

$$\langle\lambda, n| = (-1)^{j_1+\cdots+j_s}\langle n|\psi^*_{i_1+n}\cdots\psi^*_{i_s+n}\psi_{-j_s+n}\cdots\psi_{-j_1+n} \qquad (3.5.11)$$

As we see

$$\langle\lambda, n|\mu, m\rangle = \delta_{mn}\delta_{\lambda\mu} \qquad (3.5.12)$$

We see that Fock vectors (3.5.10) (and (3.5.11)) form a orthonormal basis in the Fock spaces F (and respectively) F^*.

Remark. From (3.1.16) we obtain

$$\oint\cdots\oint\psi(z_1)\cdots\psi(z_n)|0><0|\psi^*(z_n)\cdots\psi^*(z_1) = \sum_{\lambda,l(\lambda)\le n}|\lambda, n><\lambda, n| \qquad (3.5.13)$$

which projects any state to the component F_n of the Fock space $F = \bigoplus_{n\in Z} F_n$. □

Using Lemma 1 and the development (3.2.15) we get

Lemma

$$|\lambda, n\rangle = s_\lambda(\mathbf{H}^*)|n\rangle, \quad \langle\lambda, n| = \langle n|s_\lambda(\mathbf{H}) \qquad (3.5.14)$$

and

$$s_\lambda(-\mathbf{A})|n\rangle = r_\lambda(n)s_\lambda(\mathbf{H}^*)|n\rangle, \quad \langle n|s_\lambda(\tilde{\mathbf{A}}) = \tilde{r}_\lambda(n)\langle n|s_\lambda(\mathbf{H}) \qquad (3.5.15)$$

$$\mathbf{H} = \left(H_1, \frac{H_2}{2}, \ldots, \frac{H_m}{m}, \ldots\right), \quad \tilde{\mathbf{A}} = \left(\tilde{A}_1, \frac{\tilde{A}_2}{2}, \ldots, \frac{\tilde{A}_m}{m}, \ldots\right) \qquad (3.5.16)$$

where $A_m, \tilde{A}_m$ are given by (3.2.14), (3.5.2). □

There are the following developments:

Proposition 6

$$e^{-A(\mathbf{t}^*)} = \sum_{n\in Z}\sum_{\lambda\supseteq\mu}|\mu, n\rangle s_{\lambda/\mu}(\mathbf{t}^*)r_{\lambda/\mu}(n)\langle\lambda, n| \qquad (3.5.17)$$

$$e^{\tilde{A}(\mathbf{t})} = \sum_{n\in Z}\sum_{\lambda\supseteq\mu}|\lambda, n\rangle s_{\lambda/\mu}(\mathbf{t})\tilde{r}_{\lambda/\mu}(n)\langle\mu, n| \qquad (3.5.18)$$

where $s_{\lambda/\mu}(\mathbf{t})$ is the skew Schur function (3.5.8) and $r_{\lambda/\mu}(n), \tilde{r}_{\lambda/\mu}(n)$ are defined in (3.5.9).

The proof follows from the relation [4]

$$\langle s_{\lambda/\mu}, s_\nu\rangle = \langle s_\lambda, s_\mu s_\nu\rangle \tag{3.5.19}$$

Remark. It follows from (3.5.17), (3.5.18)

$$s_\nu(-A) = \sum_{n\in Z}\sum_{\lambda\supseteq\mu} C^\lambda_{\mu\nu} r_{\lambda/\mu}(n)|\mu, n\rangle\langle\lambda, n|, \quad s_\nu(\tilde{A}) = \sum_{n\in Z}\sum_{\lambda\supseteq\mu} C^\lambda_{\mu\nu}\tilde{r}_{\lambda/\mu}(n)|\lambda, n\rangle\langle\mu, n| \tag{3.5.20}$$

where coefficients $C^\lambda_{\mu\nu}$ define the decomposition of the product of Schur functions $s_\mu s_\nu = \sum_\lambda C^\lambda_{\mu\nu} s_\lambda$.

3.6 Schur function expansion of tau function. Loop scalar products

According to formulae (3.1.25) and (3.5.12) we have

$$\tau_g(n, \mathbf{t}, \mathbf{t}^*) = \langle n|e^{H(\mathbf{t})} g e^{H(\mathbf{t}^*)}|n\rangle = \sum_{\lambda,\mu} s_\lambda(\mathbf{t}) g_{\lambda\mu}(n) s_\mu(\mathbf{t}^*) \tag{3.6.1}$$

where $g_{\lambda\mu}$ is the matrix element

$$g_{\lambda\mu}(n) = \langle\lambda, n|g|\mu, n\rangle \tag{3.6.2}$$

Therefore we have the property:

$$(g_1 g_2)_{\lambda\mu}(n) = \sum_\nu (g_1)_{\lambda\nu}(n)(g_2)_{\nu\mu}(n) \tag{3.6.3}$$

We can introduce the following 'integrable' scalar product

$$< s_\lambda, s_\mu >_{g,n} = g_{\lambda\mu}(n) \tag{3.6.4}$$

In general this form is a degenerated one.

We see that we obviously have

$$< \tau_{g_1}(n_1, \mathbf{t}, \gamma), \tau_{g_1}(n_2, \gamma, \mathbf{t}^*) >_{g,n} = \tau_{g_3}(0, \mathbf{t}, \mathbf{t}^*), \quad g_3(k) = g_1(n_1+k) g(n+k) g_2(n_2+k) \tag{3.6.5}$$

In the previous consideration we actually took

$$g_{\lambda\mu}(n) = \delta_{\lambda\mu} r_\lambda(n) \tag{3.6.6}$$

Different examples of non diagonal $g_{\lambda\mu}$ expressed in terms of skew Schur functions one can obtain using the Proposition of previous subsection. It will be used in the section "Further generalization" below. Let us note that in fact non diagonal $g_{\lambda\mu}$ were constructed in the form of certain determinants in the papers [28],[29],[30].

Now let us consider the following scalar product of $\tau_{g'}$ with itself:

$$<< \tau_{g'}(n) >>_{g,n} := < \tau_{g'}(n, \gamma, \gamma) >_{g,n} = \sum_{\lambda,\mu} g_{\lambda\mu}(n) g'_{\mu\lambda}(n) \tag{3.6.7}$$

One can easily see that this scalar product is the trace of operator g in the component of the Fock space of the charge n, namely, F_n:

$$<< \tau_{g'}(n) >>_{g,n} = < \tau_{g'}(n, \gamma, \gamma) >_{g,n} = \sum_{\lambda,\mu} g_{\lambda,\mu}(n) g'_{\mu\lambda}(n) = \sum_\lambda \langle\lambda, n|g'g|\lambda, n\rangle = \mathrm{Trace}_{F_n}(g'g) \tag{3.6.8}$$

Let us consider the following **loop scalar product**:

$$<< \tau_{g_1}(n)\tau_{g_2}(n)\cdots\tau_{g_p}(n) >>_{g,n} := < \tau_{g_1}(n, \gamma^{(1)}, \gamma^{(2)})\tau_{g_2}(n, \gamma^{(2)}, \gamma^{(3)})\cdots\tau_{g_p}(n, \gamma^{(p)}, \gamma^{(1)}) >_{g,n} = \tag{3.6.9}$$

$$\sum_{\lambda_1,\dots,\lambda_p} g_{1\lambda_1\lambda_2}(n)\cdots g_{p\lambda_p\lambda_1}(n) = \mathrm{Trace}_{F_n}(g_1 g g_2 g\cdots g_p g) \tag{3.6.10}$$

As we shall see below the loop scalar product describes certain matrix integrals.

4 Angle integration of tau functions of matrix argument. Integration over complex matrices

In this section we shall generalize some known solvable matrix integrals. By the *solvable matrix integrals* we mean those which can be presented as tau functions and which can be evaluated by the method of orthogonal polynomials. Let us make a reference to the related paper [55], where by solvable matrix model the author mean the model which admits a reduction of number of integrals (from the order $\sim N^2$ to the order N) via a character expansion method. Let us refer to the papers [65],[48],[64] where different generalizations of HCIZ integral were considered.

4.1 Useful formulae

In this section we shall exploit the following formulae of an integration of Schur functions over the unitary group [4]. First is the formula

$$\int_{U(n)} s_\lambda(AUBU^{-1})d_*U = \frac{s_\lambda(A)s_\lambda(B)}{s_\lambda(I_n)}, \tag{4.1.1}$$

where d_*U is the Haar measure on the $U(n)$. Then we have

$$\int_{U(n)} s_\lambda(AU)s_\mu(U^{-1}B)d_*U = \frac{s_\lambda(AB)}{s_\lambda(I_n)}\delta_{\lambda\mu}. \tag{4.1.2}$$

If we have complex matrices Z then there are formulae [4]

$$\pi^{-n^2}\int_{-\infty}^{+\infty} s_\lambda(AZBZ^+)e^{-\mathrm{Tr}ZZ^+}\prod_{i,j=1}^{n} d\Re Z_{ij}d\Im Z_{ij} = \frac{s_\lambda(A)s_\lambda(B)}{s_\lambda(1,0,0,0,\dots)} \tag{4.1.3}$$

and

$$\pi^{-n^2}\int_{-\infty}^{+\infty} s_\lambda(AZ)s_\mu(Z^+B)e^{-\mathrm{Tr}ZZ^+}\prod_{i,j=1}^{n} d\Re Z_{ij}d\Im Z_{ij} = \frac{s_\lambda(AB)}{s_\lambda(1,0,0,0,\dots)}\delta_{\lambda\mu} \tag{4.1.4}$$

Then we need [4]

$$s_\lambda(1,0,0,0,\dots) = \frac{1}{H_\lambda}, \quad H_\lambda = \prod_{i,j}(\lambda_i + \lambda'_j - i - j + 1) \tag{4.1.5}$$

where H_λ is the hook product. The *Pochhammer's symbol related to a partition* $\lambda = (\lambda_1, \dots, \lambda_k)$ is the following product of the Pochhammer's symbols $(a)_\lambda := (a)_{\lambda_1}(a-1)_{\lambda_2}\cdots(a-k+1)_{\lambda_k}$. Then

$$(a)_\lambda = H_\lambda s_\lambda(\mathbf{t}(a)), \quad (n)_\lambda = H_\lambda s_\lambda(I_n) \tag{4.1.6}$$

In addition we have the relation (bosonic (l.h.s) and fermionic (r.h.s) representation of the vacuum TL tau function)

$$\exp\sum_{m=1}^{\infty} mt_m t_m^* = \sum_\lambda s_\lambda(\mathbf{t})s_\lambda(\mathbf{t}^*), \tag{4.1.7}$$

which is a general version of *Cauchy-Littlewood identity [4].*

Let $\mathbf{t}^*_\infty = (1,0,0,0,\ldots), \mathbf{t}(a) = (\frac{a}{1}, \frac{a}{2}, \frac{a}{3}, \ldots), r(k) = \prod_{i=1}^p (k+a_i) \prod_{i=1}^s (k+b_i)^{-1}$, $l(\lambda)$ is the number of nonvanishing parts of λ. Let us write down the formulae for the *hypergeometric function of a matrix argument [26]*, see Appendix C:

$$
{}_pF_s\left(\begin{matrix} a_1+M, \ldots, a_p+M \\ b_1+M, \cdots, b_s+M \end{matrix} \middle| X \right) = \tau_r(M, \mathbf{t}(\mathbf{x}^n), \mathbf{t}^*_\infty) =
$$

$$
\sum_{\substack{\lambda \\ l(\lambda)\le n}} \frac{\prod_{k=1}^p s_\lambda(\mathbf{t}(a_k+M))}{\prod_{k=1}^s s_\lambda(\mathbf{t}(b_k+M))} \left(s_\lambda(1,0,0,0,\ldots)\right)^{s-p+1} s_\lambda(\mathbf{x}^n). \tag{4.1.8}
$$

Now let $r(k) = \prod_{i=1}^p (k+a_i)(k+n-M)^{-1} \prod_{i=1}^s (k+b_i)^{-1}$. The *hypergeometric functions of two matrix arguments [26]*, see Appendix C, is

$$
{}_p\mathcal{F}_s\left(\begin{matrix} a_1+M, \ldots, a_p+M \\ b_1+M, \cdots, b_s+M \end{matrix} \middle| X, Y \right) = \tau_r(M, \mathbf{t}(\mathbf{x}^n), \mathbf{t}^*(\mathbf{y}^n)) =
$$

$$
\sum_{\substack{\lambda \\ l(\lambda)\le n}} \frac{\prod_{k=1}^p s_\lambda(\mathbf{t}(a_k+M))}{\prod_{k=1}^s s_\lambda(\mathbf{t}(b_k+M))} \left(s_\lambda(1,0,0,0,\ldots)\right)^{s-p+1} \frac{s_\lambda(\mathbf{x}^n) s_\lambda(\mathbf{y}^n)}{s_\lambda(I_n)}. \tag{4.1.9}
$$

Here $\mathbf{x}^n = (x_1, \ldots, x_n), \mathbf{y}^n = (y_1, \ldots, y_n)$ are eigenvalues of matrices X, Y.

Let us mark also the **determinant representations of (4.1.8) and of (4.1.9)** (details we find in Appendix A), which are of importance in applications of the method of orthogonal polynomials to the new matrix models, which we shall consider below.

The determinant representations have forms (the reader may find details in [1],[11]):

$$
\tau_r(M, \mathbf{t}(\mathbf{x}^n), \mathbf{t}^*) = \frac{\det\left(x_i^{n-k} \tau_r(M-k+1, \mathbf{t}(x_i), \mathbf{t}^*) \right)_{i,k=1}^n}{\Delta(\mathbf{x}^n)} \tag{4.1.10}
$$

and

$$
\tau_r(M, \mathbf{t}(\mathbf{x}^n), \mathbf{t}^*(\mathbf{y}^n)) = \frac{c_M}{c_{M-n}} \frac{\det\left(\tau_r(M-n+1, x_i, y_j) \right)_{i,j=1}^n}{\Delta(\mathbf{x}^n)\Delta(\mathbf{y}^n)}, \tag{4.1.11}
$$

where

$$
\Delta(\mathbf{x}^n) = \prod_{i<j} (x_i - x_j) = \det(x_i^{n-k}), \tag{4.1.12}
$$

$$
c_n = \prod_{k=0}^{n-1} \left(r(k)\right)^{k-n}, \tag{4.1.13}
$$

$$
\tau_r(M-n+1, x_i, y_j) = 1 + r(M-n+1)x_i y_j + r(M-n+1)r(M-n+2)x_i^2 y_j^2 + \cdots \tag{4.1.14}
$$

Below we need mainly the case $M = n$.

4.2 Certain matrix integrals and expansion in Schur functions

In this subsection we shall consider few integrals of exponential functions. We shall pick up the solvable models.

Integration over complex matrices. Different integrals over complex matrices were considered in the papers of I. Kostov, see [57],[58] as an example.

(A) Using relations (4.1.7) and (4.1.3) we evaluate the following integral over a complex matrix M as the expansion over Schur functions

$$
I_A = \int e^{\sum_{m=1}^\infty t_m \mathrm{Tr}\left(AZBZ^+\right)^m} e^{-\mathrm{Tr} ZZ^+} d^2 Z = \sum_{\lambda, l(\lambda)\le n} \frac{s_\lambda(\mathbf{t}) s_\lambda(A) s_\lambda(B)}{s_\lambda(1,0,0,0,\ldots)} \tag{4.2.1}
$$

where $d^2Z = \pi^{-n^2}\prod_{i,j=1}^n d\Re Z_{ij} d\Im Z_{ij}$.

Let us compare this series with the relation (4.1.9). To identify them we have the following prescription. He will specify at least one of the arguments of the Schur functions in the numerator with one of the sets: with $(1,0,0,0,\dots)$ or with $(\frac{a}{1},\frac{a}{2},\frac{a}{3},\dots)$. Let us notice that if we take $a=n$ we get the Schur function of the unit matrix I_n, which is the dimension of the representation labeled by λ. Also we mark that $s_\lambda(1,0,0,0,\dots)$ is the inverse of the hook product H_λ.

Thus we have few opportunities.

(A1) First, putting $\mathbf{t}=(1,0,0,0,\dots)$ we get

$$\int e^{\sum_{m=1}^\infty \mathrm{Tr}(AZBZ^+)} e^{-\mathrm{Tr}ZZ^+} d^2Z = \sum_{\lambda, l(\lambda)\le n} s_\lambda(A)s_\lambda(B) = \prod_{i,j}^n (1-a_ib_j)^{-1} = {}_1\mathcal{F}_0\left(n|A,B\right) \tag{4.2.2}$$

(A2) Second, putting $\mathbf{t}(a)=(\frac{a}{1},\frac{a}{2},\frac{a}{3},\dots)$, $\Re a<1$, we get the expansion of the following matrix integral

$$\int \det\left(1-\frac{1}{2}AZBZ^+\right)^{-a} e^{-\mathrm{Tr}ZZ^+} d^2Z = \sum_{\lambda, l(\lambda)\le n} (a)_\lambda s_\lambda(A)s_\lambda(B) = {}_2\mathcal{F}_0\left(a,n|A,B\right) \tag{4.2.3}$$

which in general is a divergent series.

(A3) Then let us choose the matrix B as diagonal matrix with $B_{kk}=\frac{1}{n}, k=1,\dots,n$. Let us consider the limit $n\to\infty$. In the limit with the help of (4.1.5) we get the hypergeometric function of matrix argument

$$\sum_{\lambda, l(\lambda)\le n} (a)_\lambda \frac{s_\lambda(A)}{H_\lambda} = {}_1F_0(a;A) \tag{4.2.4}$$

(B) With the help of (4.1.7), (4.1.4) we get the development

$$I_B = \int e^{\sum_{m=1}^\infty t_m \mathrm{Tr}(AZ)^m + \sum_{m=1}^\infty t_m^* \mathrm{Tr}(Z^+B)^m} e^{-\mathrm{Tr}ZZ^+} d^2Z = \sum_{\lambda, l(\lambda)\le n} \frac{s_\lambda(\mathbf{t})s_\lambda(\mathbf{t}^*)s_\lambda(AB)}{s_\lambda(1,0,0,0,\dots)} \tag{4.2.5}$$

Comparing it with (4.1.9) we obtain three solvable cases.

(B1) The first one is $\mathbf{t}^*=(1,0,0,0,\dots)$. Then we have

$$I_B = e^{\sum_{m=1}^\infty t_m \mathrm{Tr}(AB)^m} \tag{4.2.6}$$

(B2) The second one is $AB=I_n$, then we get

$$I_B = \sum_{\lambda, l(\lambda)\le n} (n)_\lambda s_\lambda(\mathbf{t})s_\lambda(\mathbf{t}^*) = {}_2\mathcal{F}_0\left(n,n|\mathbf{t},\mathbf{t}^*\right) \tag{4.2.7}$$

This expansion coincides with the expansion for the model of two Hermitian random matrices [**?**].

(B3) The third one is $\mathbf{t}^*=(\frac{a}{1},\frac{a}{2},\frac{a}{3},\dots)$, then

$$I_B = \sum_{\lambda, l(\lambda)\le n} (a)_\lambda s_\lambda(\mathbf{t})s_\lambda(AB) = {}_2\mathcal{F}_0\left(a,n|\mathbf{t},AB\right) \tag{4.2.8}$$

Integration over unitary matrices.

(C) The combination of the formula (4.1.1) with the formula (4.1.7) yields Schur function representation for different matrix integral:

$$\int_{U(n)} e^{\sum_{m=1}^{\infty} t_m \mathrm{Tr}\left(AUBU^{+}\right)^m} d_*U = \sum_{\lambda, l(\lambda)\le n} \frac{s_\lambda(\mathbf{t}) s_\lambda(A) s_\lambda(B)}{s_\lambda(I_n)} \tag{4.2.9}$$

Similar series were considered in the papers [55],[56].

Looking at this formula we see that one can choose $\mathbf{t}$ in such a way, that he gets the hypergeometric functions of matrix arguments (4.1.9).

As a consequence we evaluate the following matrix integrals

(C1) HCIZ integral

$$\int_{U(n)} e^{\mathrm{Tr}\left(AUBU^{+}\right)} d_*U = \sum_{\lambda, l(\lambda)\le n} \frac{s_\lambda(A) s_\lambda(B)}{H_\lambda s_\lambda(I_n)} = {}_0\mathcal{F}_0\left(A, B\right) = c_n \frac{\det(e^{a_i b_j})}{\Delta(a)\Delta(b)} \tag{4.2.10}$$

(which we obtain putting $t_1 = 1; t_m = 0, m > 1$) and which is the hypergeometric function of matrix arguments (4.1.9). For the last equality see (4.1.11).

(C2)

$$\int_{U(n)} \det\left(1 - AUBU^{+}\right)^{-a} d_*U = \sum_{\lambda, l(\lambda)\le n} (a)_\lambda \frac{s_\lambda(A) s_\lambda(B)}{H_\lambda s_\lambda(I_n)} = {}_1\mathcal{F}_0\left(a; A, B\right) = c_n \frac{\det(1 - a_i b_j)^{n-1-a}}{\Delta(a)\Delta(b)}, \tag{4.2.11}$$

which we obtain by putting $t_m = -a/m, m = 1, 2, \ldots$. For the last equality see (4.1.11).

(D) Let us remember the following integral , which is used for the study of two-dimensional QCD [68] and called Gross-Witten one plaquette model

$$I^{GW} = \int_{U(n)} e^{-\frac{1}{g_s}\mathrm{Tr}\left(U + U^{+}\right)} d_*U \tag{4.2.12}$$

About this integral see also [59].

A generalization of this integral was considered in [69]:

$$I^{GW}(J, J^{+}) = \int_{U(n)} e^{\mathrm{Tr}\left(JU + U^{+}J^{+}\right)} d_*U \tag{4.2.13}$$

where the method of orthogonal polynomials was applied and the links with the KP equation and with the so-called generalized Kontsevich model was shown.

Using the orthogonality of Schur functions we easily get the development

$$I^{GW}(J, J^{+}) = \sum_{\lambda, l(\lambda)\le n} \frac{s_\lambda(JJ^{+}) s_\lambda(1, 0, 0, \ldots)}{(n)_\lambda} = {}_0\mathcal{F}_0\left(JJ^{+}; (1, 0, 0, 0, \ldots)\right) \tag{4.2.14}$$

which is the TL tau function of hypergeometric type.

Let us further generalize the matrix integral (4.2.13) as follows

$$\int_{U(n)} e^{\sum_m t_m \mathrm{Tr}(AU)^m + \sum_m t^*_m \mathrm{Tr}(U^{-1}B)^m} d_*U = \sum_{\lambda, l(\lambda)\le n} \frac{s_\lambda(AB) s_\lambda(\mathbf{t}) s_\lambda(\mathbf{t}^*)}{s_\lambda(I_n)} \tag{4.2.15}$$

Now we consider two special cases of the last integral.

(D1) For the first example we take $t^*_m = b/m, m = 1, 2, \ldots \Re b < 1$), then the integral (4.2.15) is

$$\int_{U(n)} e^{\sum_m t_m \mathrm{Tr}(AU)^m} \det\left(1 - U^{-1}B\right)^{-b} d_*U = \sum_{\lambda, l(\lambda)\le n} (b)_\lambda \frac{s_\lambda(AB) s_\lambda(\mathbf{t})}{(n)_\lambda} = {}_1\mathcal{F}_0\left(b|\mathbf{t}, AB\right) \tag{4.2.16}$$

By (4.1.10) we have the determinant representation of (4.2.16).

(D2) For the second example we take $t_m = a/m, t^*_m = b/m, m = 1, 2, \dots (\Re a, \Re b < 1)$, then the integral (4.2.15) is

$$\int_{U(n)} \det (1 - AU)^{-a} \det \left(1 - U^{-1}B\right)^{-b} d_*U = \sum_{\lambda, l(\lambda) \leq n} \frac{(a)_\lambda (b)_\lambda}{(n)_\lambda} \frac{s_\lambda(AB)}{H_\lambda} = {}_2F_1 (a, b; n|AB) \tag{4.2.17}$$

which is Gauss hypergeometric function of the matrix argument AB. By (4.1.10) we have the determinant representation of (4.2.17). Let us mark that for integer a, b the logarithmic derivative of the hypergeometric function ${}_2F_1 (a, b; n|AB)$ solves Painleve V equation [14].

(D3) The third example is well-known model of unitary matrices ($A = B = 1$). See [45] about this model. Now we present the following representation

$$\int_{U(n)} e^{\sum_m t_m \mathrm{Tr} U^m + \sum_m t^*_m \mathrm{Tr} U^{-m}} d_*U = \sum_{\lambda, l(\lambda) \leq n} s_\lambda(\mathbf{t}) s_\lambda(\mathbf{t}^*) \tag{4.2.18}$$

Due to the restriction $l(\lambda) \leq n$ it is not the l.h.s of Cauchy-Littlewood formula (4.1.7). Let us mark that the right hand side of (4.2.18) is the subject of the so-called Gessel theorem if one take $\mathbf{t} = \mathbf{t}(\mathbf{x}^m), \quad \mathbf{t}^* = \mathbf{t}^*(\mathbf{y}^m), \quad n < m$. This case was considered in [15]. In this case we have different determinant representations by (4.1.10), by (4.1.11) and also by the formula for semi-infinite TL tau-function (6.8.1).

4.3 Tau functions of a matrix argument and angle integration. The integration of tau functions over complex matrices

It is suitable to introduce the notion of *tau function of matrix argument* by analogy with the hypergeometric function of matrix argument (4.1.8),(4.1.9):

$$\tau (n, X, \mathbf{t}^*) := \tau (n, \mathbf{t}(\mathbf{x}^n), \mathbf{t}^*) , \quad \tau (n, X, Y) := \tau (n, \mathbf{t}(\mathbf{x}^n), \mathbf{t}^*(\mathbf{y}^n)) \tag{4.3.1}$$

where Hirota-Miwa variables $x_1, \dots, x_n$ are just eigenvalues of a matrix X. We shall use large letters for this matrix argument. As it is in the case of the hypergeometric functions the tau function by definition depends only on eigenvalues of the matrix.

Generalization of HCIZ integral

Let us consider the averaging of the TL tau function $\tau_r(n, XUYU^+, \mathbf{t}^*)$:

$$I = \langle \tau_r \left(n, XUYU^+, \mathbf{t}^*\right) \rangle_U := \int_{U(n)} \tau_r \left(n, XUYU^+, \mathbf{t}^*\right) d_*U = \sum_{\lambda, l(\lambda) \leq n} \frac{s_\lambda(X) s_\lambda(Y) s_\lambda(\mathbf{t}^*)}{s_\lambda(I_n)} r_\lambda(n) \tag{4.3.2}$$

If we take $\mathbf{t}^* = \left(\frac{a}{1}, \frac{a}{2}, \dots\right)$ we get the following TL tau function $\tau_{\tilde{r}}(n, X, Y)$

$$\sum_{\lambda, l(\lambda) \leq n} \frac{(a)_\lambda}{(n)_\lambda} r_\lambda(n) s_\lambda(X) s_\lambda(Y) = c_n \frac{\det\{\tau_{\tilde{r}}(1, x_i, y_j)\}|_{i,j=1,\dots,n}}{\Delta(x)\Delta(y)}, \quad \tilde{r}(m) := \frac{a + m - 1}{n + m - 1} r(m) \tag{4.3.3}$$

which is certain hypergeometric function (4.1.9) in case r rational. The r.h.s is the determinant representation of the l.h.s. , see (4.1.11). $\Delta(x)$ is the Vandermond determinant.

Generalization of the integral over complex matrices

Let us consider the averaging of the TL tau function $\tau_r(n, AZBZ^+, \mathbf{t}^*)$:

$$I_C = \langle \tau_r \left(n, XZYZ^+, \mathbf{t}^*\right) \rangle_Z := \int \tau_r \left(n, XZYZ^+, \mathbf{t}^*\right) e^{-\mathrm{Tr} ZZ^+} d^2Z \tag{4.3.4}$$

We get

$$\int \tau_r \left(n, XZYZ^+, \mathbf{t}^*\right) e^{-\mathrm{Tr} ZZ^+} d^2Z = \sum_{\lambda, l(\lambda)\le n} \frac{s_\lambda(X)s_\lambda(Y)s_\lambda(\mathbf{t}^*)}{s_\lambda(1,0,0,\dots)} r_\lambda(n) \qquad (4.3.5)$$

If we take $\mathbf{t}^* = \left(\frac{a}{1}, \frac{a}{2}, \dots\right)$ we get new TL tau function $\tau_{\tilde r}(n, X, Y)$, where

$$\tilde r(m) = (a+m-1)r(m) \qquad (4.3.6)$$

Finely we obtain

$$I_C = \tau_{\tilde r}(n, \mathbf{x}, \mathbf{y}) = c_n \frac{\det\{\tau_{\tilde r}(1, x_i, y_j)\}|_{i,j=1,\dots,n}}{\Delta(x)\Delta(y)} \qquad (4.3.7)$$

see (4.1.11).

Generalization of Gross-Witten integral. This is the integral

$$I(XY) = \int_{U(n)} \tau_r\left(n, \mathbf{t}, XU\right) \tau_{\tilde r}\left(n, U^{-1}Y, \mathbf{t}^*\right) d_*U = \sum_{\lambda, l(\lambda)\le n} \frac{s_\lambda(XY)s_\lambda(\mathbf{t})s_\lambda(\mathbf{t}^*)}{s_\lambda(I_n)} \tilde r_\lambda(n) r_\lambda(n) \qquad (4.3.8)$$

For $r\tilde r = 1$ the integral (4.3.8) is equal to the integral (4.2.15).

Let $z_i, i = 1, \dots, n$ are eigenvalues of the matrix XY. The matrix integral (4.3.8) is TL tau function if

(a) $t^*_m = a/m, m = 1, 2, \dots$ (a is any complex number), then

$$I(XY) = \sum_{\lambda, l(\lambda)\le n} (a)_\lambda \tilde r_\lambda(n) r_\lambda(n) \frac{s_\lambda(XY)s_\lambda(\mathbf{t})}{(n)_\lambda} = \frac{\det\left(z_i^{n-k}\tau_{r'}(n-k+1, \mathbf{t}(z_i), \mathbf{t}^*)\right)|_{i,k=1}^n}{\Delta(\mathbf{z}^n)} \qquad (4.3.9)$$

where $r'(m) = \frac{a+m-1}{n+m-1} r(m)\tilde r(m)$. The last equality is due to (4.1.10).

(b) $t^*_m = \delta_{1,m}$, then

$$I(XY) = \sum_{\lambda, l(\lambda)\le n} \frac{s_\lambda(XY)s_\lambda(\mathbf{t})}{(n)_\lambda} \tilde r_\lambda(n) r_\lambda(n) = \frac{\det\left(z_i^{n-k}\tau_{r'}(n-k+1, \mathbf{t}(z_i), \mathbf{t}^*)\right)|_{i,k=1}^n}{\Delta(\mathbf{z}^n)} \qquad (4.3.10)$$

where $r'(m) = \frac{1}{n+m-1} r(m)\tilde r(m)$. The last equality is due to (4.1.10).

(c) and (d) one gets by $\mathbf{t} \leftrightarrow \mathbf{t}^*$.

(e) $XY = I_n$ then

$$I(XY) = \tau_{r\tilde r}(n, \mathbf{t}, \mathbf{t}^*) \qquad (4.3.11)$$

This answer may be also derived from (3.5.13).

Remark. Let us notice that the following averaging over unitary group $U(n)$

$$I(XY) = \int_{U(n)} \tau_r\left(n, XU, U^{-1}Y\right) d_*U = \sum_\lambda r_\lambda(n) \frac{s_\lambda(XY)}{s_\lambda(I_n)} \qquad (4.3.12)$$

is not TL tau function. □

In the similar way one can consider the integral over complex matrix Z

$$I(XY) = \int \tau_r\left(n, \mathbf{t}, XZ\right) \tau_{\tilde r}\left(n, Z^+Y, \mathbf{t}^*\right) e^{-\mathrm{Tr} ZZ^+} d^2Z = \sum_{\lambda, l(\lambda)\le n} \frac{s_\lambda(XY)s_\lambda(\mathbf{t})s_\lambda(\mathbf{t}^*)}{s_\lambda(1,0,0,0,\dots)} \tilde r_\lambda(n) r_\lambda(n) \qquad (4.3.13)$$

The reader will pick up the solvable cases in the way similar to the previous cases.

5 Integral representation of the scalar product and the integrals over eigenvalues of random matrices

5.1 Integral representation of the scalar product $<,>_{r,n}$

In case the function r has zero the scalar product (9.1.1) is degenerate one. For simplicity let us take $r(0)=0, r(k)\neq 0, k\le n$. The scalar product (9.1.1) non generate on the subspace of symmetric functions spanned by Schur functions $\{s_\lambda, l(\lambda)\le n\}$, $l(\lambda)$ is the length of partition λ.

To find the integral representation of this scalar product it is enough to find a function μ of one variable with the following property. If it depends on the product of two variables x and y, then moments of this function have a diagonal form:

$$\int_\Gamma \mu(xy)dxdy = A, \quad \int_\Gamma \mu(xy)x^n y^m dxdy = A\delta_{n,m} r(1)r(2)\cdots r(n), \quad n,m=1,2,\dots \tag{5.1.1}$$

where Γ is some integration domain to be defined later, and values of $r(k)$ are certain given numbers, and A is finite and does not vanish:

$$A\neq\infty, \quad A\neq 0, \tag{5.1.2}$$

The appropriate μ we shall denote by μ_r.

Example 1. The variable y is a complex conjugated of $x=z$. The integration domain is the whole complex plane, we suggest that (5.1.2) is fulfilled. Then obviously

$$\int_{\mathsf{C}} z^n \bar z^m \mu_r(|z|^2)dzd\bar z = A\delta_{nm} r(1)r(2)\cdots r(n), \quad n,m=1,2,\dots \tag{5.1.3}$$

Example 2. The variable x is real, and the variable y is imaginary one. The integration is going over the real and over the imaginary axes. We take $\mu_r(xy)=e^{-xy}$ and obtain that $r(n)=n$:

$$\int_\Re\int_\Im x^n y^m e^{-xy}dxdy = -2\pi\sqrt{-1}\delta_{nm} n!, \quad n,m=0,1,2,\dots \tag{5.1.4}$$

Example 3. Both variables x,y lie on the unit circle: $x=e^{\sqrt{-1}\phi}, y=e^{\sqrt{-1}\varphi}$. We take $\mu_r(xy)=(xy)^{-1}e^{(xy)^{-1}}$ and obtain that $r(n)=1/n$:

$$\oint\oint x^n y^m e^{(xy)^{-1}}\frac{dxdy}{xy} = -4\pi^2\delta_{nm}\frac{1}{n!}, \quad n,m=0,1,2,\dots \tag{5.1.5}$$

□

Remark. We can find an appropriate function μ_r in a following way. Let r be an analytical function of one variable and consider $r(D_x), D_x = x\frac{d}{dx}$ which acts on the power functions x^s by the multiplication by $r(s)$. One suggests that there exists a function $\mu_r(x)$ with the following properties

$$\left(1-\frac{1}{x}r(-D_x)\right)\mu_r(x)=0, \quad D_x = x\frac{d}{dx} \tag{5.1.6}$$

One may find solutions by using the Mellin transformation as

$$\mu_r(x)=\int_\gamma x^s e^{-T(-s-1)}ds \tag{5.1.7}$$

where

$$e^{T(s-1)-T(s)} = r(s) \tag{5.1.8}$$

The integration domain Γ in (5.1.1) should be chosen in such a way that the operator $\frac{1}{x}r(D), D = x\frac{d}{dx}$ is conjugated to the operator $\frac{1}{x}r(-D), D = x\frac{d}{dx}$:

$$\left(\frac{1}{x}r(D)\right)^* = \left(\frac{1}{x}r(-D)\right) \tag{5.1.9}$$

on the appropriate class of functions, as used in (5.1.1) below.

At last with the help of (5.1.6) and (5.1.9) we get (5.1.1). □

In case $\mathbf{r(0) = 0}$ the series

$$\mu_r(x) = 1 + \frac{x}{r(-1)} + \frac{x^2}{r(-1)r(-2)} + \cdots \tag{5.1.10}$$

is the formal solution of (5.1.6). This series is equal to the tau-function of hypergeometric type related to $1/r'$, where

$$r'(n) := r(-n) \tag{5.1.11}$$

$$\mu_r(xy) = \tau_{1/r'}\left(1, \mathbf{t}(x), \mathbf{t}^*(y)\right), \quad kt_k = x^k, \quad kt_k^* = y^k \tag{5.1.12}$$

Let us suggest that the series (5.1.10) is convergent (this fact depends on the choice of the function r) , and the integral (5.1.1) exists ($A \neq 0, \infty$), then one may choose this μ_r for (5.1.21).

Example 1a. We consider the case of Example 1 above. The hypergeometric function ${}_pF_s(a_1, \dots, a_p; b_1, \dots, b_s; -|z|^2), p \leq s$ may be taken as an example of the function μ_r of the Example 1. We put $|z|^2 = x$ below. We have

$$\int_0^\infty x^n {}_pF_s(a_1, \dots, a_p; b_1, \dots, b_s; -x)dx = (-)^{(n+1)(p-s)} n! \frac{\prod_{i=1}^s (1-b_i)_{n+1}}{\prod_{i=1}^p (1-a_i)_{n+1}}, \tag{5.1.13}$$

see for instance [40]. In this case

$$A = \frac{\prod_{i=1}^s (b_i - 1)}{\prod_{i=1}^p (a_i - 1)}, \quad r(n) = n\frac{\prod_{i=1}^s (b_i - n - 1)}{\prod_{i=1}^p (a_i - n - 1)} \tag{5.1.14}$$

It is known that the hypergeometric function ${}_pF_s, p \leq s$

$${}_pF_s\left(a_1, \dots, a_p; b_1, \dots, b_s; -x\right) = \sum_{n=0}^\infty \frac{(a_1)_n \cdots (a_p)_n}{(b_1)_n \cdots (b_s)_n} \frac{(-x)^n}{n!}. \tag{5.1.15}$$

is a convergent hypergeometric series. It is of form (5.1.10). This series is a solution of

$$\left(x + D_x \frac{\prod_{i=1}^s (D_x + b_i - 1)}{\prod_{i=1}^p (D_x + a_i - 1)}\right) {}_pF_s(a_1, \dots, a_p; b_1, \dots, b_s; -x) = 0 \tag{5.1.16}$$

which the hypergeometric equation (8.0.12) rewritten in the form (5.1.6), where xy is replaced by $-x$. The operator $r(-D_x)$ acts on each x^n of the convergent series as the multiplication by $r(-n)$.□

Example 1b. We choose $r(k) = -k(a-1-k)^{-1}$, then ${}_1F_0(a;x) = (1-x)^{-a}$ is the solution of (5.1.6), for the case $\Re a < 0$ the condition (5.1.9) is fulfilled inside the interval $[0,1]$, and for $\Re a < 1$ we have

$$\int_0^1 x^n {}_1F_0(a;x)dx = A\frac{n!}{(2-a)_n}; \quad A = (1-a)^{-1} \tag{5.1.17}$$

□

Now we shall turn to the **multiply integrals**. Let us evaluate the integral

$$I_{\lambda,\nu} = A^{-n}\int\cdots\int \Delta(\mathbf{x})\Delta(\mathbf{y})s_\lambda(\mathbf{x})s_\nu(\mathbf{y})\prod_{k=1}^n \mu_r(x_ky_k)dx_kdy_k \tag{5.1.18}$$

where $\mathbf{x} = (x_1,\ldots,x_n), \mathbf{y} = (y_1,\ldots,y_n)$

$$\Delta(\mathbf{z}) = \prod_{i<j}^n (z_i - z_j) \tag{5.1.19}$$

Using (5.1.1), the definition of the Schur functions (2.1.10) and the definition of $r_\lambda(n)$ (2.2.1) (see [2] for details) we finely get the relation

$$I_{\lambda\nu} = r_\lambda(n)\delta_{\nu,\lambda} \tag{5.1.20}$$

and we obtain

Proposition 7

$$\langle f,g\rangle_{r,n} = \frac{1}{A^n}\int_\Gamma\cdots\int_\Gamma \Delta(\mathbf{x})\Delta(\mathbf{y})f(\mathbf{x})g(\mathbf{y})\prod_{k=1}^n \mu_r(x_ky_k)dx_kdy_k \tag{5.1.21}$$

Then it follows that

Proposition 8

$$\langle f,g\rangle_{r,n} = \frac{1}{n!A^n}\int_\Gamma\cdots\int_\Gamma f(\mathbf{x})g(\mathbf{y})\det\left(\mu(x_iy_j)\right)_{i,j=1,\ldots,n}\Delta(\mathbf{x})\Delta(\mathbf{y})\prod_{k=1}^n dx_kdy_k \tag{5.1.22}$$

The proof of the last Proposition follows from the fact that the function $\det\left(\mu(x_iy_j)\right)_{i,j=1,\ldots,n}$ is an anti-symmetric function in variables $x_1,\ldots,x_n$. Then by changing notations of the variables x_i inside of the integral we obtain the Proposition.

Let us remember that if

$$\mathbf{t}(\mathbf{x}) = (t_1(\mathbf{x}),t_2(\mathbf{x}),\ldots), \quad \mathbf{t}^*(\mathbf{y}) = (t_1^*(\mathbf{y}),t_2^*(\mathbf{y}),\ldots), \quad mt_m = \sum_{i=1}^n x_i^m, \quad mt_m^* = \sum_{i=1}^n y_i^m \tag{5.1.23}$$

we have the following determinant representation [9] , [10] (see an Appendix below (6.7.4), (6.2.11), (6.2.12))

$$\Delta(\mathbf{x})\Delta(\mathbf{y})\tau_r\left(n,\mathbf{t}(\mathbf{x}),\mathbf{t}^*(\mathbf{y})\right) = c_n\det\left(\tau_r\left(1,x_i,y_k\right)\right)_{i,k=1}^n, \quad c_n = \prod_{k=0}^{n-1}\left(r(k)\right)^{k-n} \tag{5.1.24}$$

where $\mathbf{x} = (x_1,\ldots,x_n), \mathbf{y} = (y_1,\ldots,y_n)$ which is true on the level of formal series.

$\tau_{1/r'}\left(n,\mathbf{t}(\mathbf{x}),\mathbf{t}^*(\mathbf{y})\right)$ is the tau-function (3.4.8), where the function r is chosen as $1/r'$

$$\tau_{1/r'}\left(n,\mathbf{t}(\mathbf{x}),\mathbf{t}^*(\mathbf{y})\right) = 1 + \frac{\left(\sum_{i=1}^n x_i\right)\left(\sum_{i=1}^n y_i\right)}{r(-1)} + \cdots \tag{5.1.25}$$

Due to determinant representation (5.1.24) we see that this series is a convergent one, since we consider the series (5.1.12) to be convergent.

Proposition 9 *Let the tau function $\tau_{1/r'}$ of (3.4.8), where $r'(n) := r(-n)$, is a convergent series, and let $r(0) = 0$. Then we have the following realization of scalar product (2.2.2)*

$$\langle f, g\rangle_{r,n} = A^{-n}c_n^{-1}\int_\Gamma \cdots \int_\Gamma f(\mathbf{x})g(\mathbf{y})\tau_{1/r'}\left(n, \mathbf{t}(\mathbf{x}), \mathbf{t}^*(\mathbf{y})\right)\left(\Delta(\mathbf{x})\right)^2\left(\Delta(\mathbf{y})\right)^2\prod_{k=1}^n dx_k dy_k \tag{5.1.26}$$

The proof follows from the fact that the function $\tau_{1/r'}\left(n, \mathbf{t}(\mathbf{x}), \mathbf{t}^*(\mathbf{y})\right)$ is a symmetric function in variables $x_1, \ldots, x_n$. Then by changing notations of the variables x_i inside of the integral we obtain the Proposition.

Then it follows from (5.1.26),(2.2.13) that for the same conditions the following duality relation is true

$$\tau_r\left(n, \mathbf{t}, \mathbf{t}^*)\right) = \tag{5.1.27}$$

$$A^{-n}c_n^{-1}\int_\Gamma \cdots \int_\Gamma e^{\sum_{i=1}^n\sum_{m=1}^\infty (t_m x_i^m + t_m^* y_i^m)}\tau_{1/r'}\left(n, \mathbf{t}(\mathbf{x}), \mathbf{t}^*(\mathbf{y})\right)\left(\Delta(\mathbf{x})\right)^2\left(\Delta(\mathbf{y})\right)^2\prod_{k=1}^n dx_k dy_k \tag{5.1.28}$$

Also due to (2.2.16),(2.2.17) we have the following

$$\tau_{r_3}\left(0, \mathbf{t}, \mathbf{t}^*\right) = \tag{5.1.29}$$

$$A^{-n}c_n^{-1}\int_\Gamma \cdots \int_\Gamma \tau_{r_1}\left(k, \mathbf{t}, \mathbf{t}^*(\mathbf{x})\right)\tau_{1/r'}\left(n, \mathbf{t}(\mathbf{x}), \mathbf{t}^*(\mathbf{y})\right)\tau_{r_2}\left(m, \mathbf{t}(\mathbf{y}), \mathbf{t}^*\right)\Delta^2(\mathbf{x})\Delta^2(\mathbf{y})\prod_{k=1}^n dx_k dy_k \tag{5.1.30}$$

where

$$r_3(i) = r_1(k+i)r_2(m+i)r_(n+i) \tag{5.1.31}$$

5.2 Normal matrix model

Let us consider a model of normal matrices. A matrix is called *normal* if it commutes with Hermitian conjugated matrix.

Let M be n by n normal matrix. The following integral is called the model of normal matrices:

$$I^{NM}(n, \mathbf{t}, \mathbf{t}^*; \mathbf{u}) = \int dM dM^+ e^{U(MM^+)+V_1(M)+V_2(M^+)} = \tag{5.2.1}$$

$$C\int_\mathsf{C} \cdots \int_\mathsf{C} |\Delta(\mathbf{z})|^2 e^{\sum_{i=1}^n U(z_i\bar z_i)+V_1(z_i)+V_2(\bar z_i)}\prod_{i=1}^n dz_i d\bar z_i \tag{5.2.2}$$

where $dM = \prod_{i<k} d\Re M_{ik} d\Im M_{ik}\prod_{i=1}^N dM_{ii}$, in the last integral the integration is going over complex planes of eigenvalues z_i, C is a number which appears due to the matrix 'angle integration' over V ($M = VZV^{-1}$, where $Z = \mathrm{d}iag\{z_n\}$), and U is defined by $\mathbf{u} = (u_1, u_2, \ldots)$:

$$U(MM^+) = \sum_{n=1}^\infty u_n \mathrm{Tr}(MM^+)^n, \quad V_1(M) = \sum_{m=1}^\infty t_m \mathrm{Tr}M^m, \quad V_2(M) = \sum_{m=1}^\infty t_m^* \mathrm{Tr}\left(M^+\right)^m \tag{5.2.3}$$

$$\Delta(\mathbf{z}) = \prod_{i<k}^n (z_i - z_k) \tag{5.2.4}$$

and $\Delta(z) = 1$ for $n = 1$.

As we see the integral (5.2.2) has a form of (5.1.21), where $\mathbf{x} = \mathbf{z}, \mathbf{y} = \bar{\mathbf{z}}$, see Example 1 of this section, and

$$f(\mathbf{z}) = e^{\sum_{i=1}^n\sum_{m=1}^\infty t_m z_i^m}, \quad g(\bar{\mathbf{z}}) = e^{\sum_{i=1}^n\sum_{m=1}^\infty t_m^* \bar z_i^m}, \quad \mu_r(z_i\bar z_i) = e^{U(z_i\bar z_i)} \tag{5.2.5}$$

Now let us compare e^U with μ_r of (5.1.10). We choose the integration domain Γ to be the whole complex plane. We suppose that

$$\pi\sqrt{-1}\int_0^{+\infty} e^{\sum_{n=1}^\infty u_n z^n} dz = A \neq 0, \infty \tag{5.2.6}$$

The condition (5.1.2) is fulfilled. If he function $e^{U(|z|^2)}$ has a form of (5.1.10) (each Taylor coefficient is non-vanishing), then the set of variables $\mathbf{u} = (u_1, u_2, \ldots)$ is related to the set $\{r(n), n = -1, -2, \ldots\}$ as follows

$$h_m(\mathbf{u}) = \frac{1}{r(-1)} \cdots \frac{1}{r(-m)}, \quad r(-m) = \frac{h_{m-1}(\mathbf{u})}{h_m(\mathbf{u})}, \quad e^{\sum_{k=1}^\infty u_k z^k} = \sum_{m=0}^{\infty} h_m(\mathbf{u}) z^m \tag{5.2.7}$$

Example. $T_m = \delta_{m,1}$. Then

$$h_m(\mathbf{u}) = \frac{1}{\Gamma(m+1)}, \quad r(-m) = m \tag{5.2.8}$$

□

Finely formula (2.2.13), where $p_m = \sum_{i=1}^n z_i^m$, yields the series for (5.2.1):

$$I^{NM}(n, \mathbf{t}, \mathbf{t}^*; \mathbf{u}) =< e^{\sum_{i=1}^n \sum_{m=1}^\infty t_m z_i^m}, e^{\sum_{i=1}^n \sum_{m=1}^\infty t_m^* z_i^m} >_{r,n} = \sum_\lambda r_\lambda(n) s_\lambda(\mathbf{t}) s_\lambda(\mathbf{t}^*) \tag{5.2.9}$$

5.3 Two-matrix model

Details of this and of the next section one can find in [2].

Let us evaluate the following integral over n by n matrices M_1 and M_2, where M_1 is a Hermitian matrix and M_2 is an anti-Hermitian one

$$I^{2MM}(n, \mathbf{t}, \mathbf{t}^*) = \int e^{V_1(M_1)+V_2(M_2)} e^{-\mathrm{Tr} M_1 M_2} dM_1 dM_2 \tag{5.3.1}$$

where

$$V_1(M_1) = \sum_{m=1}^{\infty} t_m \mathrm{Tr} M_1^m, \quad V_2(M_2) = \sum_{m=1}^{\infty} t_m^* \mathrm{Tr} M_2^m \tag{5.3.2}$$

It is well-known [16],[5] that this integral reduces to the integral over eigenvalues x_i and y_i of matrices M_1 and M_2 respectively.

To do it first we diagonalize the matrices: $M_1 = U_1 X U_1^{-1}, M_2 = \sqrt{-1} U_2 Y U_2^{-1}$, where U_1, U_2 are unitary matrices and X, Y are diagonal matrices with real entries. Then one calculates the Jacobians of the change of variables $M_1 \to (X, U_1), M_2 \to (Y, U_2) : dM_1 = (\Delta(\mathbf{x}))^2 \prod_{k=1}^n dx_k \prod_{i<j} d\Re(U_1)_{ij} d\Im(U_1)_{ij}$ and $dM_2 = (\Delta(\mathbf{y}))^2 \prod_{k=1}^n dy_k \prod_{i<j} d\Re(U_2)_{ij} d\Im(U_2)_{ij}$. As a result we get

$$I^{2MM}(n, \mathbf{t}, \mathbf{t}^*) = \int e^{\sum_{i=1}^n \sum_{m=1}^\infty t_m x_i^m} e^{\sum_{i=1}^n \sum_{m=1}^\infty t_m y_i^m} I(\mathbf{x}, \mathbf{y}) \Delta^2(\mathbf{x}) \Delta^2(\mathbf{y}) \prod_{k=1}^{n} dx_k dy_k \int d_* U_1 \tag{5.3.3}$$

where

$$I(\mathbf{x}, \mathbf{y}) = \int e^{-\mathrm{Tr} XUYU^{-1}} d_* U \tag{5.3.4}$$

is the so-called **Harish-Chandra-Itzykson-Zuber (HCIZ) integral**. The integration (5.3.4) is over unitary group $U(n)$, $d_* U$ is Haar measure. This integral was calculated [16], [5] :

$$I(\mathbf{x}, \mathbf{y}) = \frac{\det \left(e^{-x_i y_k \sqrt{-1}} \right)_{i,k=1}^n}{\Delta(\mathbf{x}) \Delta(\mathbf{y})} \tag{5.3.5}$$

Due to (5.1.24) formula (5.3.5) is a manifest of the fact that the Harish-Chandra-Itzykson-Zuber integral actually is a tau-function (see a different approach to this fact in [47]). This is the tau-function τ_r which corresponds to $r(n) = -1/n$, in other words we can write

$$\frac{I(\mathbf{x},\mathbf{y})}{0!\cdots(n-1)!} = \tau_{1/r'}\left(n,\mathbf{t}(\mathbf{x}),\mathbf{t}^*(\mathbf{y})\right), \quad 1/r'(n) = -\frac{1}{n} \tag{5.3.6}$$

For $n = 1$, $x_1 = x, y_1 = y$ we have

$$\tau_{1/r'}\left(1,\mathbf{t}(\mathbf{x}),\mathbf{t}^*(\mathbf{y})\right) = e^{-xy\sqrt{-1}} \tag{5.3.7}$$

This was the subject of the Example (5.1.4).

Therefore due to (5.1.26) we obtain that the model of two Hermitian random matrices is the series of hypergeometric type (2.2.13), where $p_m = \sum_{i=1}^n z_i^m$ and $r(n) = n$:

$$I^{2MM}(n,\mathbf{t},\mathbf{t}^*) = \frac{C_n}{(2\pi)^n} < e^{\sum_{i=1}^n\sum_{m=1}^\infty t_m z_i^m}, e^{\sum_{i=1}^n\sum_{m=1}^\infty t_m^* z_i^m} >_{r,n} = \frac{C_n}{(2\pi)^n}\sum_\lambda (n)_\lambda s_\lambda(\mathbf{t})s_\lambda(\mathbf{t}^*) \tag{5.3.8}$$

where we remember that according to (2.2.1)

$$(n)_\lambda = \prod_{i,j\in\lambda}(n+j-i) = \frac{\Gamma(n+1+\lambda_1)\Gamma(n+\lambda_2)\cdots\Gamma(\lambda_n)}{\Gamma(n+1)\Gamma(n)\cdots\Gamma(1)} \tag{5.3.9}$$

and C_n incorporates the volume of unitary group.

The same result one can obtain with the help of (5.1.21), using the known result that integral (5.3.3) is equal to

$$C_n\int_{-\infty}^{\infty}\cdots\int_{-\infty}^{\infty} e^{\sum_{i=1}^n\sum_{m=1}^\infty t_m x_i^m} e^{\sum_{i=1}^n\sum_{m=1}^\infty t_m y_i^m} e^{-\sum_{i=1}^n x_i y_i\sqrt{-1}}\Delta(\mathbf{x})\Delta(\mathbf{y})\prod_{i=1}^n dx_i dy_i \tag{5.3.10}$$

where x_i and y_i are eigenvalues of M_1 and M_2 respectively.

This integral has a form of (5.2.5) if one takes the integrals over variables x_i be along the real axe, while the integrals over variables y_i be along the imaginary axe. Then we take $r(n) = n$ so that

$$\mu_r(xy) = \tau_{1/r'} = 1 - xy + \frac{x^2y^2}{2} + \cdots = e^{-xy} \tag{5.3.11}$$

$$\int_\Re\int_\Im e^{-xy}dxdy = 2\pi \tag{5.3.12}$$

Thus we have the following perturbation series [2]

$$\frac{I^{2MM}(n,\mathbf{t},\mathbf{t}^*)}{I^{2MM}(n,0,0)} = \sum_\lambda (n)_\lambda s_\lambda(\mathbf{t})s_\lambda(\mathbf{t}^*) \tag{5.3.13}$$

In case all higher coupling constants are equal to 0 (namely, in the case of Gauss matrix integral) this series can be easily evaluated as

$$I^{2MM}(1,t_1,t_2,0,0,\ldots;t_1^*,t_2^*,0,0,\ldots) = \sum_{m=0}^\infty m!h_m(\mathbf{t})h_m(\mathbf{t}^*) = \frac{\exp\frac{t_1t_1^*+t_2\left(t_1^*\right)^2+t_2^*(t_1)^2}{1-4t_2t_2^*}}{\sqrt{1-4t_2t_2^*}} \tag{5.3.14}$$

and for $n > 1$ one can use the formula (for example see [2])

$$\tau_r\left(n,\mathbf{t},\mathbf{t}^*\right) = \frac{1}{(n-1)!\cdots 0!}\det\left(\frac{\partial^{k+m}}{\partial_{t_1}^k\partial_{t_1^*}^m}\tau_r\left(1,\mathbf{t},\mathbf{t}^*\right)\right)_{k,m=0}^{n-1} \tag{5.3.15}$$

which is true for the case $r(0) = 0$.

5.4 Hermitian one matrix model

Below we shall follow [2].

We suggest a fermionic representation of the well-known partition function of one matrix model [8] (which generalizes the known Gaussian unitary ensemble (GUE), [5])

$$Z(N,g,g_4) = \int dM e^{-N\mathrm{Tr}\left(\frac{g}{2}M^2+g_4M^4\right)}, \tag{5.4.1}$$

this fermionic representation being different from [45],[60]. The reason is that, as it was before in the paper, we consider the matrix model in the sense of the perturbation series .

We chose the normalization of the matrix integral in such a way that it is equal to 1 when $g_4 = 0$.

Let us evaluate simplest perturbation terms for one matrix model using the series (5.3.14) [2]. Let us consider the model of two Hermitian matrices (5.3.1), and take all $t_k = 0$ except t_4, and all $t_k^* = 0$ except t_2^*. Then as it is known [49] if we put

$$g_4 = -4N^{-1}t_4, \quad g = (2Nt_2^*)^{-1} \tag{5.4.2}$$

we obtain that

$$I^{2MM}(0,0,0,t_4,0,\ldots;0,t_2^*,0,0,\ldots) = Z(N,g,g_4) \tag{5.4.3}$$

and therefore we get

$$\sum_\lambda (N)_\lambda s_\lambda(\mathbf{t}) s_\lambda(\mathbf{t}^*) = \int dM e^{-N\mathrm{Tr}\left(\frac{g}{2}M^2+g_4M^4\right)} = Z(N,g,g_4) \tag{5.4.4}$$

Remark. As it is well-known, see for instance [50], according to Feynman rules for one matrix model we have (a) to each propagator (double line) is associated a factor $1/(Ng)$ (which is $2t_2^*$ in our notations) (b) to each four leg vertex is associated a factor $(-Ng_4)$ (which is $4t_4$ in our notations) (c) to each closed single line is associated a factor N. Therefore one may say that the factors $(N)_\lambda$ in (5.4.4) is responsible for closed lines, the factors $s_\lambda(\mathbf{t} = 0, t_2, 0, \ldots)$ are responsible for propagators and the factors $s_\lambda(\mathbf{t}^*)$ are responsible for vertices. As we see we get the following structure for the Feynman diagrams, containing $k = |\lambda|/4$ vertices and $2k = |\lambda|/2$ propagators:

$$(Ng_4)^{|\lambda|/4}(Ng)^{-|\lambda|/2} \sum_{|\lambda|=4k} (N)_\lambda a_\lambda b_\lambda \tag{5.4.5}$$

where the number $|\lambda|$ is the weight of partition λ, and a_λ, b_λ are numbers:

$$a_\lambda = s_\lambda(0,0,0,1,0,\ldots), \quad b_\lambda = s_\lambda(0,1,0,\ldots) \tag{5.4.6}$$

Let us also notice that to evaluate the numbers a_λ, b_λ one can use either the definition of Schur functions (3.1.26)-(3.1.27), or one can use the following formulae, which we borrow in [56]:

$$b_\lambda = \prod_{k=1}^{N-1} k! \frac{\Delta(h^{even})\Delta(h^{odd})}{\prod_i \left(\frac{h^{even}}{2}\right) \prod_i \left(\frac{h^{odd}-1}{2}\right)} sign \left[\prod_{i,j}(h^{even} - h^{odd})\right] \tag{5.4.7}$$

$$a_\lambda = \prod_{k=1}^{N-1} k! \prod_{\epsilon=0}^{3} \frac{\Delta(h^\epsilon)}{\prod_i \left(\frac{h^\epsilon - \epsilon}{2}\right)} sign \left[\prod_{0\le\epsilon_1<\epsilon_2\le 3} \prod_{i,j}(h_i^{(\epsilon_2)} - h_j^{(\epsilon_1)})\right] \tag{5.4.8}$$

where we keep the notation h_i of [56] for shifted weights $\lambda_i - i$ (not miss with the complete symmetric function). Here integers h factor into 4 groups of $\frac{N}{4}$ integers h^ϵ with $\epsilon = 0, 1, 2, 3$ denoting their congruence modulo 4 [56]. □

Since only t_4 is non vanishing the first non vanishing Schur functions correspond to λ of weight 4, which are $(4), (3,1), (2,2), (2,1,1), (1,1,1,1)$. Let us notice that two last partition are conjugated to the two first, and the partition $(2,2)$ is self conjugated.

To evaluate l.h.s. of (5.4.4) we use (2.2.1) , (2.1.12) and take into account the property for conjugated partition λ' we have

$$s_{\lambda'}(-\mathbf{t}) = (-)^{|\lambda|} s_\lambda(\mathbf{t}), \quad r'_{\lambda'}(-n) = r_\lambda(n) \tag{5.4.9}$$

where $|\lambda|$ is the number of boxes in the Young diagram (or the same - the weight of the partition), and for r' see (3.4.9).

The partition $\lambda = (4)$ gives

$$N(N+1)(N+2)(N+3)t_4 \frac{(t_2^*)^2}{2!} = (N^4 + 6N^3 + 11N^2 + 6N)\left(-\frac{Ng_4}{4}\right)\left(\frac{1}{8N^2g^2}\right) \tag{5.4.10}$$

Due to (5.4.9) , (5.4.2) the conjugated partition $(1,1,1,1)$ gives the same answer but $N \to -N$. (Thus one needs to keep only even powers of N).

The partition $(3,1)$ gives

$$N(N+1)(N+2)(N-1)(-t_4)\frac{(-t_2^*)^2}{2!} = (N^4 + 2N^3 - N^2 - 2N)\left(-\frac{Ng_4}{4}\right)\left(\frac{1}{8N^2g^2}\right) \tag{5.4.11}$$

Again due to the contribution of the conjugated partition $(2,1,1)$ one needs to keep only even powers of N.

First Schur function in the l.h.s of (5.4.4) is equal to zero (due to (2.1.12) and since only t_4 is non vanishing).

Finely we find first perturbation terms as

$$1 - \left(\frac{N^2}{2} + \frac{1}{4}\right)\frac{g_4}{g^2} + \cdots \tag{5.4.12}$$

This answer coincides with the well-known answer one gets by Feynman diagram method, see [50]

For the next order one takes partitions of weight 8, as a consequence that it is only t_4 that is non vanishing. One can get that non vanishing contribution is only due to self conjugated partitions $(3,3,2)$, $(4,2,1,1)$, and to partitions $(8); (7,1), (4,4); (6,1,1), (4,3,1); (5,1,1,1)$ and conjugated to these ones. The enumerated partitions yields respectively h_4^2, h_4^2, then $h_8; -h_8, h_4^2; -h_8, -h_4^2; -h_8$ for $s_\lambda(\mathbf{t}^*)$. And it yields respectively $h_4^2 - h_4h_2^2, h_4^2 - h_4h_2^2$, then $h_8; -h_8, h_4^2; -h_8 - h_6h_2, -h_4^2 + h_6h_2; -h_8 + h_6h_2$ for $s_\lambda(\mathbf{t})$. For partitions $(3,3,2)$, $(4,2,1,1)$, and for partitions $(8); (7,1), (4,4); (6,1,1), (4,3,1); (5,1,1,1)$ we get respectively

$$N(N+1)(N+2)(N-1)(N)(N+1)(N-2)(N-1)\left(t_4^2\right)\left(\left(\frac{(t_2^*)^2}{2!}\right)^2 - \frac{(t_2^*)^2}{2!}(t_2^*)^2\right) \tag{5.4.13}$$

$$N(N+1)(N+2)(N+3)(N-1)(N)(N-2)(N-3)(t_4^2)\left(\left(\frac{(t_2^*)^2}{2!}\right)^2 - \frac{(t_2^*)^2}{2!}(t_2^*)^2\right) \tag{5.4.14}$$

and

$$N(N+1)(N+2)(N+3)(N+4)(N+5)(N+6)(N+7)\left(\frac{t_4^2}{2!}\right)\left(\frac{(t_2^*)^4}{4!}\right) \tag{5.4.15}$$

$$N(N+1)(N+2)(N+3)(N+4)(N+5)(N+6)(N-1)\left(-\frac{t_4^2}{2!}\right)\left(-\frac{(t_2^*)^4}{4!}\right) \tag{5.4.16}$$

$$N(N+1)(N+2)(N+3)(N-1)(N)(N+1)(N+2)\left(t_4^2\right)\left(\frac{(t_2^*)^2}{2!}\right)^2 \tag{5.4.17}$$

$$N(N+1)(N+2)(N+3)(N+4)(N+5)(N-1)(N-2)\left(\frac{t_4^2}{2!}\right)\left(\frac{(t_2^*)^4}{4!}-t_2^*\frac{(t_2^*)^3}{3!}\right) \tag{5.4.18}$$

$$N(N+1)(N+2)(N+3)(N-1)(N)(N+1)(N-2)\left(-t_4^2\right)\left(-\left(\frac{(t_2^*)^2}{2!}\right)^2+t_2^*\frac{(t_2^*)^3}{3!}\right) \tag{5.4.19}$$

$$N(N+1)(N+2)(N+3)(N+4)(N-1)(N-2)(N-3)\left(-\frac{t_4^2}{2!}\right)\left(-\frac{(t_2^*)^4}{4!}+t_2^*\frac{(t_2^*)^3}{3!}\right) \tag{5.4.20}$$

At the highest order at N which is N^6 one gets zero.

$$(32N^6+320N^4+488N^2)t_4^2(t_2^*)^4=(32N^6+320N^4+488N^2)\left(\frac{Ng_4}{4}\right)^2\left(\frac{2}{Ng}\right)^4= \tag{5.4.21}$$

$$\frac{g_4^2}{g^4}(32N^4+320N^2+488)$$

Finely

$$Z(N,g,g_4)=1-\frac{g_4}{g^2}\left(\frac{N^2}{2}+\frac{1}{4}\right)+\frac{g_4^2}{g^4}(32N^4+320N^2+488)+O\left(\frac{g_4^3}{g^6}\right) \tag{5.4.22}$$

5.5 Unitary two-matrix model

Let us consider the integral

$$I^{UM}(n,\mathbf{t},\mathbf{t}^*)=\int_{U(n)}e^{\sum_{m=1}^\infty t_m \mathrm{Tr}U_1^m}e^{\sum_{m=1}^\infty t_m^* \mathrm{Tr}U_2^m}e^{\mathrm{Tr}U_1^{-1}U_2^{-1}}d_*U_1d_*U_2 \tag{5.5.1}$$

This model was considered in [47]. By the Example 3 of the subsection 4.1 we have

$$I^{UM}(n,\mathbf{t},\mathbf{t}^*)=\sum_\lambda\frac{s_\lambda(\mathbf{t})s_\lambda(\mathbf{t}^*)}{(n)_\lambda}=c_n\frac{\det(e^{a_ib_j})}{\Delta(a)\Delta(b)} \tag{5.5.2}$$

(the last formula we obtain via (4.1.11)).This expression coincides with the series HCIZ integral (5.3.6). First this model was identified with HCIZ integral in [47].

5.6 New multi-matrix models which can be solved by the method of orthogonal polynomials and the Schur function expansion

Here we consider new solvable multi-matrix integrals which we obtain with the help of the subsection *Angle integration of tau functions. Integration over complex matrices.*

Let us remind that the multi-matrix model of Hermitian matrices

$$I=\int e^{V_1(M_1)+\cdots+V_N(M_N)}e^{\mathrm{Tr}M_1M_2}e^{\mathrm{Tr}M_2M_3}\cdots e^{\mathrm{Tr}M_{N-1}M_N}dM_1\cdots dM_N \tag{5.6.1}$$

where $M_1, \ldots, M_N$ are Hermitian n by n matrices and

$$V_k(M_k) = \sum_{m=1}^{\infty} t_m^{(k)} \mathrm{Tr} M_k^m, \quad k = 1, \ldots, N, \tag{5.6.2}$$

was shown to be multi-component KP tau function and solved by the method of orthogonal polynomials.

Now we consider more general integral over matrices $M_1, \ldots, M_N$

$$I = \int e^{V_1(M_1)+\cdots+V_N(M_N)} R_1(M_1M_2)R_2(M_2M_3)\cdots R_{N-1}(M_{N-1}M_N)dM_1\cdots dM_N \tag{5.6.3}$$

If we choose the interaction term as hypergeometric function of matrix argument (see (4.3.1))

$$R_k(M_kM_{k+1}) = \tau_{r_k}(n, \mathbf{t}, M_kM_{k+1}), \quad \mathbf{t} = (1, 0, 0, 0, \ldots) \tag{5.6.4}$$

then due to (4.3.2) it is possible to perform the angle integration over each $U_k(n)$ where $M_k = U_k X^{(k)} U_k^+, \quad X^{(k)} = diag(x_1^{(k)}, \ldots, x_n^{(k)})$. Really, (4.3.2) is a tau function which has a determinant representation (4.1.11). Finely we obtain the following integral over eigenvalues $x_i^{(k)} \quad (i = 1, \ldots, n; k = 1, \ldots, N)$:

$$I = c \int \prod_{i=1}^{n} \rho_k(x_i^{(1)}x_i^{(2)})\rho_2(x_i^{(2)}x_i^{(3)})\cdots\rho_{N-1}(x_i^{(N-1)}x_i^{(N)}) \prod_{k=1}^{N} e^{\sum_{k=1}^{N}\sum_{m=1}^{\infty} t_m^{(k)}(x_i^{(k)})^m} dx_i^{(k)} \tag{5.6.5}$$

where

$$\rho_k(x_i^{(k)}x_i^{(k+1)}) = \tau_{r_k}(1, \mathbf{t}, x_i^{(k)}x_i^{(k+1)}), \quad \mathbf{t} = (1, 0, 0, 0, \ldots) \tag{5.6.6}$$

(In the case of the familiar multi-matrix model (5.6.1) each $\rho_k(xy), k = 1, \ldots, N$ is e^{-xy}).

The integral (5.6.5) may be evaluated by the *method of orthogonal polynomials* $\{p_m(t^{\mathbf{N}}, x)\}$, $\{\pi_m(t^{\mathbf{N}}, x)\}$ which depend on the collection of times $t^{\mathbf{N}} = (\mathbf{t}^{(1)}, \mathbf{t}^{(2)}, \ldots, \mathbf{t}^{(N)})$

$$\int p_m(t^{\mathbf{N}}, x)\pi_n(t^{\mathbf{N}}, y)\omega(t^{\mathbf{N}}, x, y)dxdx = \delta_{nm}e^{\phi_n(t^{\mathbf{N}})}, \quad n, m = 0, 1, 2, \ldots \tag{5.6.7}$$

Each of polynomials is of a form $p_m(t^{\mathbf{N}}, x) = \sum_{n\le m} p_{nm}(t^{\mathbf{N}})x^n, \quad \pi_m(t^{\mathbf{N}}, x) = \sum_{n\le m} \pi_{nm}(t^{\mathbf{N}})x^n$, and the weight function is

$$\omega(t^{\mathbf{N}}, x, y) = e^{V_1(x)+V_N(y)} \int \rho_k(xx^{(2)})\rho_2(x^{(2)}x^{(3)})\cdots\rho_{N-1}(x^{(N-1)}y) \prod_{k=2}^{N-1} e^{V_k(x^{(k)})}dx^{(k)} \tag{5.6.8}$$

We obtain

$$I = c\prod_{k=1}^{n} e^{\phi_k(t^{\mathbf{N}})} \tag{5.6.9}$$

Let us notice that the orthogonal polynomials are Baker-Akhiezer functions of TL hierarchy. Gauss-Zakharov-Shabat factorization problem for TL equation [27] directly results from the relation (5.6.7)

$$K_+(t^{\mathbf{N}}) = K_-(t^{\mathbf{N}})e^{\Phi}G(t^{\mathbf{N}}), \quad \Phi = diag(\phi_1(t^{\mathbf{N}}), \phi_2(t^{\mathbf{N}}), \ldots) \tag{5.6.10}$$

where

$$(K_+)_{nm} = (\pi^{-1})_{nm}, \quad (K_-)_{nm} = p_{nm}, \quad G_{nm} = \int x^n y^m \omega(t^{\mathbf{N}}, x, y)dxdx \tag{5.6.11}$$

(as it was in the case of two-matrix model [44]).

The Schur function expansion for these matrix matrix models is given by (1.0.9), where $K_{\lambda\mu}$ are expressed via products of skew Schur functions.

5.7 Standard scalar product

Let us suggest a realization of the standard scalar product (3.3.2). This scalar product is a particular case of (3.3.9). In this case $r \equiv 1$. This function has no zeroes, therefore the consideration of the previous subsection is not valid. We realize (3.3.2) as follows

$$< f, g >= \int f(\mathbf{t}) e^{-\sum_{m=1}^{\infty} m|t_m|^2} g(\bar{\mathbf{t}}) \prod_{m=1}^{\infty} \frac{md^2t_m}{\pi} \qquad (5.7.1)$$

where $\bar{t}_m$ is the complex conjugated of t_m. The formula for the scalar product of two tau-functions is similar to (5.1.29):

$$< \tau_{r_1}(k), \tau_{r_2}(m) >= \tau_{r_3}(0) = \qquad (5.7.2)$$

where due to (3.4.8),(3.3.2) $r_3(i) = r_1(k+i)r_2(m+i)$, and

$$= \int \tau_{r_1}(k, \mathbf{t}, \gamma) e^{-\sum_{k=1}^{\infty} k|\gamma_k|^2} \tau_{r_2}(m, \bar{\gamma}, \mathbf{t}^*) \prod_{i=1}^{\infty} \frac{kd^2\gamma_k}{\pi} \qquad (5.7.3)$$

Different representation of the standard scalar product one can get in the following way. Consider

$$< \tau_{r_1}(k), \tau_{r_2}(m) >_n = \int_{U(n)} \tau_{r_1}(k, \mathbf{t}, U) \tau_{r_2}(m, U^{-1}, \mathbf{t}^*) d_*U = \sum_{l(\lambda)\le n} (r_1 r_2)_\lambda s_\lambda(\mathbf{t}) s_\lambda(\mathbf{t}^*), \qquad (5.7.4)$$

where the tau functions of matrix arguments are defined by (4.3.1). In case the restriction $l(\lambda) \le n$ is irrelevant the standard scalar product coincides with the scalar product $<,>_n$ of (5.7.4).

5.8 About spectral determinants

In many problems it is interesting to evaluate the following integral over n by n matrices

$$K(n : a, b, c, d) = \int \frac{\prod_{i=1}^{p_a} \det(a_i - M_1) \prod_{i=1}^{p_b} \det(b_i - M_1)}{\prod_{i=1}^{p_c} \det(c_i - M_2) \prod_{i=1}^{p_d} \det(d_i - M_2)} e^{V_1(M_1)+V_2(M_2)} e^{U(M_1,M_2)} dM_1 dM_2 \qquad (5.8.1)$$

where

$$V_1(M_1) = \sum_{m=1}^{\infty} t_m \mathrm{Tr} M_1^m, \quad V_2(M_2) = \sum_{m=1}^{\infty} t_m^* \mathrm{Tr} M_2^m \qquad (5.8.2)$$

The integral (5.8.1) is called a spectral determinant. In case of one matrix model the results for the product of the determinants were obtained in [63]. In the paper [66] the interesting results were obtained to evaluate the rational expressions of the determinants.

In case the integral

$$I(\mathbf{t}, \mathbf{t}^*) = \int e^{V_1(M_1)+V_2(M_2)} e^{U(M_1,M_2)} dM_1 dM_2 \qquad (5.8.3)$$

is a tau function of hypergeometric type, the spectral determinant is also a tau-function. By the Wick theorem it is equal to a determinant. Really $K(x, y) =$

$$\langle n| e^{H(\mathbf{t})} \psi(\frac{1}{a_1}) \cdots \psi(\frac{1}{a_{p_a}}) \psi^*(\frac{1}{c_{p_c}}) \cdots \psi^*(\frac{1}{c_1}) g \psi^*(b_1) \cdots \psi^*(b_{p_b}) \psi(d_{p_d}) \cdots \psi(d_1) e^{H(\mathbf{t}^*)} |n\rangle \qquad (5.8.4)$$

Look Appendix A below. The explicit expression is rather long and we shall write down it somewhere else.

6 Appendices A

6.1 Baker-Akhiezer functions and bilinear identities [7]

Vertex operators $V_\infty(\mathbf{t},z)$, $V_\infty^*(\mathbf{t},z)$ and $V_0(\mathbf{t}^*,z)$, $V_0^*(\mathbf{t}^*,z)$ act on the space $C[t_1,t_2,\ldots]$ of polynomials in infinitely many variables, and are defined by the formulae:

$$V_\infty(\mathbf{t},z) = z^M e^{\xi(\mathbf{t},z)} e^{-\xi(\tilde\partial,z^{-1})}, \quad V_\infty^*(\mathbf{t},z) = z^{-M} e^{-\xi(\mathbf{t},z)} e^{\xi(\tilde\partial,z^{-1})}, \tag{6.1.1}$$

$$V_0(\mathbf{t}^*,z) = z^M e^{-\xi(\mathbf{t}^*,z^{-1})} e^{\xi(\tilde\partial^*,z)}, \quad V_0^*(\mathbf{t}^*,z) = z^{-M} e^{\xi(\mathbf{t}^*,z^{-1})} e^{-\xi(\tilde\partial^*,z)}, \tag{6.1.2}$$

where $\tilde\partial = (\frac{\partial}{\partial t_1}, \frac12\frac{\partial}{\partial t_2}, \frac13\frac{\partial}{\partial t_3},\ldots)$, $\tilde\partial^* = (\frac{\partial}{\partial t_1^*}, \frac12\frac{\partial}{\partial t_2^*}, \frac13\frac{\partial}{\partial t_3^*},\ldots)$.

We have the rules of the bosonization:

$$\langle M+1|e^{H(\mathbf{t})}\psi(z) = V_\infty(\mathbf{t},z)\langle M|e^{H(\mathbf{t})}, \quad \langle M-1|e^{H(\mathbf{t})}\psi^*(z) = V_\infty^*(\mathbf{t},z)\langle M|e^{H(\mathbf{t})}\frac{dz}{z}, \tag{6.1.3}$$

$$\psi(z)e^{H^*(\mathbf{t}^*)}|M\rangle = V_0(\mathbf{t}^*,z)e^{H^*(\mathbf{t}^*)}|M-1\rangle, \psi^*(z)e^{H^*(\mathbf{t}^*)}|M\rangle = V_0^*(\mathbf{t}^*,z)e^{H^*(\mathbf{t}^*)}|M+1\rangle\frac{dz}{z} \tag{6.1.4}$$

The Baker-Akhiezer functions and conjugated Baker-Akhiezer functions are:

$$w_\infty(M,\mathbf{t},\mathbf{t}^*,z) = \frac{V_\infty(\mathbf{t},z)\tau}{\tau}, \quad w_\infty^*(M,\mathbf{t},\mathbf{t}^*,z) = \frac{V_\infty^*(\mathbf{t},z)\tau}{\tau}, \tag{6.1.5}$$

$$w_0(M,\mathbf{t},\mathbf{t}^*,z) = \frac{V_0(\mathbf{t}^*,z)\tau(M+1)}{\tau(M)}, \quad w_0^*(M,\mathbf{t},\mathbf{t}^*,z) = \frac{V_0^*(\mathbf{t}^*,z)\tau(M-1)}{\tau(M)}, \tag{6.1.6}$$

where

$$\tau(M) = \tau(M,\mathbf{t},\mathbf{t}^*) = \langle M|e^{H(\mathbf{t})} g e^{H^*(\mathbf{t}^*)}|M\rangle. \tag{6.1.7}$$

Both KP and TL hierarchies are described by the bilinear identity:

$$\oint w_\infty(M,\mathbf{t},\mathbf{t}^*,z)w_\infty^*(M',\mathbf{t}',\mathbf{t}'^*,z)dz = \oint w_0(M,\mathbf{t},\mathbf{t}^*,z^{-1})w_0^*(M',\mathbf{t}',\mathbf{t}'^*,z^{-1})z^{-2}dz, \tag{6.1.8}$$

which holds for any $\mathbf{t},\mathbf{t}^*,\mathbf{t}',\mathbf{t}'^*$ for any integers M, M'.

The Schur functions $s_\lambda(\mathbf{t})$ are well-known examples of tau-functions which correspond to rational solutions of the KP hierarchy. It is known that not any linear combination of Schur functions turns to be a KP tau-function, in order to find these combinations one should solve bilinear difference equation, see [23], which is actually a version of discrete Hirota equation. Below we shall present KP tau-functions which are infinite series of Schur polynomials, and which turn to be known hypergeometric functions (8.0.20),(8.0.26).

6.2 $H_0(\mathbf{T})$, fermions $\psi(T,z), \psi^*(T,z)$, bosonization rules [12]

Throughout this section we assume that $r, \tilde r \neq 0$.
Put

$$r(n) = e^{T_{n-1}-T_n}, \quad \tilde r(n) = e^{\tilde T_{n-1}-\tilde T_n} \tag{6.2.1}$$

New variables $T_n, n \in Z$ (and also $\tilde T_n$) are defined up to a constant independent of n.
We define a generator $H_0(T) \in \widehat{gl}(\infty)$:

$$H_0(T) := \sum_{n=-\infty}^{\infty} T_n :\psi_n^*\psi_n:, \tag{6.2.2}$$

which produces the following transformation:

$$e^{\mp H_0(T)}\psi_n e^{\pm H_0(T)} = e^{\pm T_n}\psi_n, \quad e^{\mp H_0(T)}\psi_n^* e^{\pm H_0(T)} = e^{\mp T_n}\psi_n^* \tag{6.2.3}$$

This is the transformation which sends $A_m, \tilde{A}_m$ of (3.2.14), (3.5.2) to H_{-m}, H_m respectively. Therefore

$$e^{H_0(\tilde{T})}\tilde{A}(T)e^{-H_0(\tilde{T})} = H(\mathbf{t}), \quad e^{-H_0(T)}A(\mathbf{t}^*)e^{H_0(T)} = -H^*(\mathbf{t}^*). \tag{6.2.4}$$

It is convenient to consider the following fermionic operators:

$$\psi(T,z) = e^{H_0(T)}\psi(z)e^{-H_0(T)} = \sum_{n=-\infty}^{n=+\infty} e^{-T_n} z^n \psi_n, \tag{6.2.5}$$

$$\psi^*(T,z) = e^{H_0(T)} \sum_{n=-\infty}^{n=+\infty} z^{-n-1}\psi_n^* e^{-H_0(T)} = \sum_{n=-\infty}^{n=+\infty} e^{T_n} z^{-n-1}\psi_n^*. \tag{6.2.6}$$

Now we consider Hirota-Miwa change of variables: $\mathbf{t} = \pm\mathbf{t}(\mathbf{x}^N)$ and $\mathbf{t}^* = \pm\mathbf{t}^*(\mathbf{y}^N)$.

Considering (6.1.3),(6.1.4) and using (6.2.4),(6.2.5),(6.2.6) (6.1.3),(6.1.4) we get the following bosonization rules

$$e^{-A(\mathbf{t}^*(\mathbf{y}^N))}|M\rangle = \frac{\psi(T,y_1)\cdots\psi(T,y_N)|M-N\rangle}{\Delta^+(M,N,T,\mathbf{y}^N)}, \tag{6.2.7}$$

$$e^{-A(-\mathbf{t}^*(\mathbf{y}^N))}|M\rangle = \frac{\psi^*(T,y_1)\cdots\psi^*(T,y_N)|M+N\rangle}{\Delta^-(M,N,T,\mathbf{y}^N)}, \tag{6.2.8}$$

$$\langle M|e^{\tilde{A}(T(\mathbf{x}^N))} = \frac{\langle M-N|\psi^*(-\tilde{\mathbf{T}},\frac{1}{x_N})\cdots\psi^*(-\tilde{\mathbf{T}},\frac{1}{x_1})}{\tilde{\Delta}^+(M,N,\tilde{\mathbf{T}},\mathbf{x}^N)}, \tag{6.2.9}$$

$$\langle M|e^{\tilde{A}(-\mathbf{t}(\mathbf{x}^N))} = \frac{\langle M+N|\psi(-\tilde{\mathbf{T}},\frac{1}{x_N})\cdots\psi(-\tilde{\mathbf{T}},\frac{1}{x_1})}{\tilde{\Delta}^-(M,N,\tilde{\mathbf{T}},\mathbf{x}^N)}. \tag{6.2.10}$$

The coefficients are

$$\Delta^\pm(M,N,T,\mathbf{y}^N) = \frac{\prod_{i<j}^N(y_i-y_j)}{(y_1\cdots y_N)^{N\mp M}} \times \frac{\tau(M,\mathbf{0},\mathbf{T},\mathbf{0})}{\tau(M\mp N,\mathbf{0},\mathbf{T},\mathbf{0})}, \tag{6.2.11}$$

$$\tilde{\Delta}^\pm(M,N,\tilde{\mathbf{T}},\mathbf{x}^N) = \frac{\prod_{i<j}^N(x_i-x_j)}{(x_1\cdots x_N)^{N-1\mp M}} \times \frac{\tau(M,\mathbf{0},\tilde{\mathbf{T}},\mathbf{0})}{\tau(M\mp N,\mathbf{0},\tilde{\mathbf{T}},\mathbf{0})}, \tag{6.2.12}$$

where in case $N=1$ the Vandermond products $\prod_{i<j}^N(y_i-y_j), \prod_{i<j}^N(x_i-x_j)$ are replaced by 1. The origin of the notation $\tau(M,\mathbf{0},\mathbf{T},\mathbf{0})$ is explained latter, see (6.3.5),(6.3.7), and used for

$$\tau(n,\mathbf{0},T,\mathbf{0}) = e^{T_{n-1}+\cdots+T_1+T_0} = e^{nT_0}\prod_{k=0}^{n-1}(r(k))^{k-n}, \quad n>0, \tag{6.2.13}$$

$$\tau(0,\mathbf{0},T,\mathbf{0}) = 1, \quad n=0, \tag{6.2.14}$$

$$\tau(n,\mathbf{0},T,\mathbf{0}) = e^{-T_n-\cdots-T_{-2}-T_{-1}} = e^{nT_0}\prod_{k=0}^{-n-1}(r(k))^{k+n}, \quad n<0. \tag{6.2.15}$$

Therefore in Hirota-Miwa variables one can rewrite the vacuum expectation of the exponents of (3.5.1) and (3.5.2):

$$\langle M|e^{\tilde{A}(\mathbf{t}(\mathbf{x}^N))}e^{-A(\mathbf{t}^*(\mathbf{y}^N))}|M\rangle =$$

$$\frac{\langle M-N|\psi^*(-\tilde{\mathbf{T}},\frac{1}{x_N})\ldots\psi^*(-\tilde{\mathbf{T}},\frac{1}{x_1})\psi(T,y_1)\cdots\psi(T,y_N)|M-N\rangle}{\tilde{\Delta}^+(M,N,\tilde{\mathbf{T}},\mathbf{x}^N)\Delta^+(M,N,T,\mathbf{y}^N)}, \tag{6.2.16}$$

$$\langle M|e^{\tilde{A}(-\mathbf{t}(\mathbf{x}^N))}e^{-A(-\mathbf{t}^*(\mathbf{y}^N))}|M\rangle = \frac{\langle M+N|\psi(-\tilde{\mathbf{T}},\frac{1}{x_N})\cdots\psi(-\tilde{\mathbf{T}},\frac{1}{x_1})\psi^*(T,y_1)\cdots\psi^*(T,y_N)|M+N\rangle}{\tilde{\Delta}^-(M,N,\tilde{\mathbf{T}},\mathbf{x}^N)\Delta^-(M,N,T,\mathbf{y}^N)}. \qquad (6.2.17)$$

This representation is suitable for an application of Wick rule (3.1.6) , see (6.7.4) below.

6.3 Toda lattice tau-function $\tau(n,\mathbf{t},T,\mathbf{t}^*)$ [12]

Here we consider *Toda lattice* (TL) [19],[7],[27]. Our notations $M,\mathbf{t},\mathbf{t}^*$ correspond to the notations $s,x,-y$ respectively in [27].

Let us consider a special type TL tau-function (3.1.25), which depends on the three sets of variables $\mathbf{t},T,\mathbf{t}^*$ and on $M\in Z$:

$$\tau(M,\mathbf{t},T,\mathbf{t}^*) = \langle M|e^{H(\mathbf{t})}\exp\left(\sum_{-\infty}^{\infty}T_n : \psi_n^*\psi_n :\right)e^{H^*(\mathbf{t}^*)}|M\rangle, \qquad (6.3.1)$$

where $:\psi_n^*\psi_n := \psi_n^*\psi_n - \langle 0|\psi_n^*\psi_n|0\rangle$. Since the operator $\sum_{-\infty}^{\infty} : \psi_n^*\psi_n :$ commutes with all elements of the $\widehat{gl}(\infty)$ algebra, one can put $T_{-1}=0$ in (6.3.1). With respect to the KP and the TL dynamics the variables T_n have a meaning of integrals of motion.

As we shall see the hypergeometric functions (8.0.10),(8.0.13),(8.0.20),(8.0.17) listed in the Appendix are ratios of tau-functions (6.3.1) evaluated at special values of times $M,\mathbf{t},T,\mathbf{t}^*$. It is true only in the case when all parameters a_k of the hypergeometric functions are non integers. For the case when at least one of the indices a_k is an integer, we will need a tau-function of an open Toda chain which will be considered in the Appendix.

Tau-function (6.3.1) is linear in each e^{T_n}. It is described by the Proposition

Proposition 10

$$\frac{\tau(M,\mathbf{t},T,\mathbf{t}^*)}{\tau(M,\mathbf{0},T,\mathbf{0})} = 1+\sum_{\lambda\neq\mathbf{0}} e^{(T_{M-1}-T_{\lambda_1+M-1})+(T_{M-2}-T_{\lambda_2+M-2})+\cdots+(T_{M-l}-T_{\lambda_l+M-l})}s_\lambda(\mathbf{t})s_\lambda(\mathbf{t}^*). \quad (6.3.2)$$

The sum is going over all different partitions

$$\lambda = (\lambda_1,\lambda_2,\dots,\lambda_l), \quad l=1,2,3,..., \qquad (6.3.3)$$

excluding the partition $\mathbf{0}$.

Let $r\neq 0$,$\tilde{r}\neq 0$. Then we put

$$r(n) = e^{T_{n-1}-T_n}, \quad \tilde{r}(n) = e^{\tilde{T}_{n-1}-\tilde{T}_n}. \qquad (6.3.4)$$

to show the equivalence of (3.4.8) and (6.3.2) in this case. We have

$$\tau(n,\mathbf{0},T,\mathbf{0}) = e^{-T_{n-1}-\cdots-T_1-T_0} = e^{-nT_0}\prod_{k=0}^{n-1}(r(k))^{n-k}, \quad n>0, \qquad (6.3.5)$$

$$\tau(0,\mathbf{0},T,\mathbf{0}) = 1, \quad n=0, \qquad (6.3.6)$$

$$\tau(n,\mathbf{0},T,\mathbf{0}) = e^{T_n+\cdots+T_{-2}+T_{-1}} = e^{-nT_0}\prod_{k=0}^{-n-1}(r(k))^{-k-n}, \quad n<0. \qquad (6.3.7)$$

Proposition 11 *Let* $\tau(n,\mathbf{t},T,\mathbf{t}^*)$ *is the Toda lattice tau function (6.3.1). Then*

$$\frac{\tau(n,\mathbf{t},\tilde{T}+T,\mathbf{t}^*)}{\tau(n,\mathbf{0},\tilde{T}+T,\mathbf{0})} = \langle n|e^{\tilde{A}(\mathbf{t})}e^{-A(\mathbf{t}^*)}|n\rangle = \tau_{\tilde{r}r}(n,\mathbf{t},\mathbf{t}^*). \tag{6.3.8}$$

where

$$A(\mathbf{t}^*) = \sum_{k=1}^{\infty} A_k t_k^*, \quad \tilde{A}(\mathbf{t}) = \sum_{k=1}^{\infty} \tilde{A}_k t_k, \tag{6.3.9}$$

$A_k, \tilde{A}_k$ *are defined by (3.5.3),(3.5.4). The functions* $\tilde{r}, r$ *are related to* $\tilde{T}, T$ *by (6.3.4), the product* $r\tilde{r}$ *has no zeroes at integer values of it's argument.*

This proposition follows from formulas(6.2.3)-(6.2.4).

We shall consider $\tilde{r} = 1$.

The notation $\tau_r(M,\mathbf{t},\mathbf{t}^*)$ will be used only for the KP tau-function (3.4.6). Notation $\tau(M,\mathbf{t},T,\mathbf{t}^*)$ is used for Toda lattice tau-function. These tau-functions are related via (6.3.8). KP tau-function $\tau(n) := \tau_r(n,\mathbf{t},\mathbf{t}^*)$ obeys the following Hirota equation:

$$\tau(n)\partial_{t_1^*}\partial_{t_1}\tau(n) - \partial_{t_1}\tau(n)\partial_{t_1^*}\tau(n) = r(n)\tau(n-1)\tau(n+1). \tag{6.3.10}$$

The equation

$$\partial_{t_1}\partial_{t_1^*}\phi_n = r(n)e^{\phi_{n-1}-\phi_n} - r(n+1)e^{\phi_n-\phi_{n+1}} \tag{6.3.11}$$

which is similar to the Toda lattice equation holds for

$$\phi_n(\mathbf{t},\mathbf{t}^*) = -\log\frac{\tau_r(n+1,\mathbf{t},\mathbf{t}^*)}{\tau_r(n,\mathbf{t},\mathbf{t}^*)} \tag{6.3.12}$$

Equations (6.3.11) and (6.3.10) are still true in case r has zeroes.
If the function r has no integer zeroes, using the change of variables

$$\varphi_n = -\phi_n - T_n, \tag{6.3.13}$$

we obtain Toda lattice equation in the standard form [27]:

$$\partial_{t_1}\partial_{t_1^*}\varphi_n = e^{\varphi_{n+1}-\varphi_n} - e^{\varphi_n-\varphi_{n-1}}. \tag{6.3.14}$$

As we see from (6.3.13) the variables T_n might have the meaning of asymptotic values of the fields ϕ_n for the class of tau-functions (6.3.1) which is characterized by the property $\varphi_n \to 0$ as $t_1 \to 0$.

6.4 Expansion of e^{H_0} and its trace in the Fock subspace F_n

$$e^{H_0(T)} = \sum_{n=-\infty}^{\infty} c_n \sum_{\lambda} r_\lambda |\lambda,n\rangle\langle\lambda,n| \tag{6.4.1}$$

It then follows that the loop

$$<< \tau_r(n) >>_{r,n} = \sum_{\lambda} r_\lambda(n) = \mathrm{Tr}_{F_n} e^{H_0(T)} \tag{6.4.2}$$

where the trace is taken in the n-th component of Fock space F_n.

6.5 Expressions and linear equations for tau-functions I (Hirota-Miwa variables)

It is well-known fact that tau-functions solves bilinear equations. The class of tau-functions under our consideration solves also linear equations.

The relations of this subsection correspond to the case when at least one set of variables (either $\mathbf{t}$ or $\mathbf{t}^*$ or both ones) is substituted with Hirota-Miwa variables.

In cases $\mathbf{t} = \mathbf{t}(\mathbf{x}^N)$ (using (6.2.9) and (3.4.3)) and $\mathbf{t} = -\mathbf{t}(\mathbf{x}^N)$ (using (6.2.10) and (3.4.2)) one gets the following representations

$$\tau_r(M, \mathbf{t}(\mathbf{x}^N), \mathbf{t}^*) = \Delta^{-1}\left(e^{\xi_{r'}(\mathbf{t}^*,x_1)} \cdots e^{\xi_{r'}(\mathbf{t}^*,x_N)} \cdot \Delta\right), \quad \Delta = \frac{\prod_{i<j}(x_i - x_j)}{(x_1 \cdots x_N)^{N-1-M}} \tag{6.5.1}$$

$$\tau_r(M, -\mathbf{t}(\mathbf{x}^N), \mathbf{t}^*) = \Delta^{-1}\left(e^{-\xi_r(\mathbf{t}^*,x_1)} \cdots e^{-\xi_r(\mathbf{t}^*,x_N)} \cdot \Delta\right), \quad \Delta = \frac{\prod_{i<j}(x_i - x_j)}{(x_1 \cdots x_N)^{N-1+M}} \tag{6.5.2}$$

where the exponents are formal series $e^\xi = 1 + \xi + \cdots$, and $r(D_i)\prod_k x_k^{a_k} = r(a_i)\prod_k x_k^{a_k}$. Let us write down that according to (3.4.4),(3.4.9)

$$\xi_{r'}(\mathbf{t}^*, x_i) = \sum_{m=1}^{+\infty} t_m^* \left(x_i r(D_i)\right)^m, \quad \xi_r(\mathbf{t}^*, x) = \sum_{m=1}^{+\infty} t_m^* \left(x_i r(-D_i)\right)^m, \quad D_i = x_i \frac{\partial}{\partial x_i} \tag{6.5.3}$$

Let us note that $[\xi_{r'}(\mathbf{t}^*, x_n), \xi_{r'}(\mathbf{t}^*, x_m)] = 0 = [\xi_r(\mathbf{t}^*, x_n), \xi_r(\mathbf{t}^*, x_m)]$ for all n, m.

(6.5.2) may be also obtained from (6.5.1) with the help of (3.4.10).

From (6.5.1) it follows for $m = 1, 2, \ldots$ that

$$\left(\frac{\partial}{\partial t_m^*} - \sum_{i=1}^{N} (x_i r(D_{x_i}))^m\right)\left(\Delta \tau_r(M, \mathbf{t}(\mathbf{x}^N), \mathbf{t}^*)\right) = 0, \quad \Delta = \frac{\prod_{i<j}(x_i - x_j)}{(x_1 \cdots x_N)^{N-1-M}}. \tag{6.5.4}$$

In case $\mathbf{t} = \mathbf{t}(\mathbf{x}^N), \mathbf{t}^* = \mathbf{t}^*(\mathbf{y}^{N^*})$ we get

$$\tau_r(M, \mathbf{t}(\mathbf{x}^N), \mathbf{t}^*(\mathbf{y}^{N^*})) = \sum_{\lambda \in P} r_{\mathbf{n}}(M) s_\lambda(\mathbf{x}^N) s_\lambda(\mathbf{y}^{N^*}) \tag{6.5.5}$$

$$= \frac{1}{\Delta(x)} \prod_{i=1}^{N} \prod_{j=1}^{N^*} (1 - y_j x_i r(D_{x_i}))^{-1} \cdot \Delta(x) \tag{6.5.6}$$

$$= \frac{1}{\Delta(y)} \prod_{i=1}^{N} \prod_{j=1}^{N^*} (1 - x_i y_j r(D_{y_j}))^{-1} \cdot \Delta(y) \tag{6.5.7}$$

where

$$\Delta(x) = \frac{\prod_{i<j}^{N}(x_i - x_j)}{(x_1 \cdots x_N)^{N-1-M}}, \quad \Delta(y) = \frac{\prod_{i<j}^{N^*}(y_i - y_j)}{(y_1 \cdots y_N)^{N^*-1-M}} \tag{6.5.8}$$

Looking at (6.5.6) ,(6.5.7) one derives the following system of linear equations

$$\left(D_{y_j} - \sum_{i=1}^{N} \frac{1}{1 - y_j x_i r(D_{x_i})} + N\right)\left(\Delta(x)\tau_r(M, \mathbf{t}(\mathbf{x}^N), \mathbf{t}^*(\mathbf{y}^{N^*}))\right) = 0, \quad j = 1, \ldots, N^* \tag{6.5.9}$$

$$\left(D_{x_i} - \sum_{j=1}^{N^*} \frac{1}{1 - x_i y_j r(D_{y_j})} + N^*\right)\left(\Delta(y)\tau_r(M, \mathbf{t}(\mathbf{x}^N), \mathbf{t}^*(\mathbf{y}^{N^*}))\right) = 0, \quad i = 1, \ldots, N \tag{6.5.10}$$

In some examples below we shall take specify $\mathbf{t}^*$ as follows.

(1) Let us take

$$mt_m = \sum_{k=1}^{N} x_k^m, \quad \mathbf{t}^* = (1,0,0,\dots) \tag{6.5.11}$$

Then one consider the equation (6.5.19) with $k=1$ (taking $t_1^* = 1$):

$$\tau_r(M,\mathbf{t}(\mathbf{x}^N),\mathbf{t}^*) = \frac{1}{\Delta} e^{x_1 r(D_1)+\dots+x_N r(D_N)} \cdot \Delta, \tag{6.5.12}$$

where

$$\Delta = \frac{\prod_{i<j}(x_i - x_j)}{(x_1 \cdots x_N)^{N-M-1}} \tag{6.5.13}$$

and $D_i = x_i \partial_{x_i}$. We get

$$\left(\sum_{i=1}^{N} D_i - \frac{1}{\Delta} \left(\sum_{i=1}^{N} x_i r(D_i) \right) \Delta \right) \tau_r(M,\mathbf{t}(\mathbf{x}^N),\mathbf{t}^*) = 0 \tag{6.5.14}$$

(2) Let us take

$$mt_m = \sum_{k=1}^{N} x_k^m, \quad mt_m^* = \frac{z^m}{1-q^m}, \quad m = 1,2,\dots \tag{6.5.15}$$

Let us note that the tau-function $\tau_r(\mathbf{t},\mathbf{t}^*)$ will not change if instead of (6.5.15) one take

$$mt_m = \sum_{k=1}^{N} (zx_k)^m, \quad mt_m^* = \frac{1}{1-q^m}, \quad m = 1,2,\dots \tag{6.5.16}$$

This gives the representation as follows

$$\tau_r(M,\mathbf{t}(\mathbf{x}^N),\mathbf{t}^*) = \frac{1}{\Delta(x)} \prod_{i=1}^{N} \frac{1}{(x_i r(D_{x_i});q)_\infty} \cdot \Delta(x) \tag{6.5.17}$$

We see that

$$\left(q^{\frac{N^2-N}{2}-NM} q^{D_1+\cdots D_N} - \prod_{i=1}^{N} (1 - x_i r(D_i)) \right) \left(\Delta(x) \tau_r(M,\mathbf{t}(\mathbf{x}^N),\mathbf{t}^*) \right) = 0 \tag{6.5.18}$$

We shall write down more linear equations, which follow from the explicit fermionic representation of the tau-function (3.4.6) via the bosonization formulae (6.2.16) and (6.2.17) These equations may be also viewed as the constraint which result in the string equations (6.14.4) and (6.14.5).

For the variables $\mathbf{x}^N$ in case $\mathbf{t} = \mathbf{t}(\mathbf{x}^N)$, one can use the relation $\langle M | A_m = 0$, and makes profit of the relation $A_m = e^{H_0} H_{-m} e^{-H_0}$ inside the fermionic vacuum expectation value (6.2.16). This way we get the partial differential equations for the tau-function (3.4.8):

$$\frac{\partial \tau_r(M,\mathbf{t}(\mathbf{x}^N),\mathbf{t}^*)}{\partial t_m^*} = \frac{1}{\Delta} \left(\sum_{i=1}^{N} (x_i r(D_{x_i}))^m \right) \Delta \tau_r(M,\mathbf{t}(\mathbf{x}^N),\mathbf{t}^*), \quad m = 1,2,3,\dots, \tag{6.5.19}$$

where Δ is proportional to $\tilde{\Delta}^+$ of (6.2.12):

$$\Delta = \frac{\prod_{i<j}(x_i - x_j)}{(x_1 \cdots x_N)^{N-1-M}}. \tag{6.5.20}$$

In variables $\mathbf{y}^{(\infty)}$, $\mathbf{t}^* = -\mathbf{t}^*(\mathbf{y}^{(\infty)})$, we can rewrite (6.5.19):

$$(-1)^k \sum_{i=1}^{+\infty} \frac{e_{k-1}\left(\frac{1}{y_1}, \dots, \frac{1}{y_{i-1}}, \frac{1}{y_{i+1}}, \dots\right)}{\prod_{j\neq i}(1-\frac{y_i}{y_j})} \frac{\partial \tau_r(M, -\mathbf{t}(\mathbf{x}^N), -\mathbf{t}^*(\mathbf{y}^{(\infty)}))}{\partial y_i} =$$
$$\frac{1}{\tilde{\Delta}} \left(\sum_{i=1}^{N} (x_i r(-D_{x_i}))^k \right) \tilde{\Delta} \tau_r(M, -\mathbf{t}(\mathbf{x}^N), -\mathbf{t}^*(\mathbf{y}^{(\infty)})), \tag{6.5.21}$$

where $e_k(\mathbf{y})$ is a symmetric function defined through the relation $\prod_{i=1}^{+\infty}(1+ty_i) = \sum_{k=0}^{+\infty} t^k e_k(\mathbf{y})$.

Also we have

$$\left(\sum_{k=1}^{M+N-1} k - \sum_{i=1}^{N} D_{x_i} \right) \tilde{\Delta}^- \tau(M, -\mathbf{t}(\mathbf{x}^N), T, -\mathbf{t}^*(\mathbf{y}^{(N')}))\Delta^- =$$
$$\left(\sum_{k=1}^{M+N'-1} k - \sum_{i=1}^{N'} \left(\frac{1}{y_i} D_{y_i} y_i \right) \right) \tilde{\Delta}^- \tau(M, -\mathbf{t}(\mathbf{x}^N), T, -\mathbf{t}^*(\mathbf{y}^{(N')}))\Delta^-, \tag{6.5.22}$$

where $\tilde{\Delta}^- = \tilde{\Delta}^-(M, N, \mathbf{0}, \mathbf{x}^N)$ and $\Delta^- = \Delta^-(M, N, \mathbf{0}, \mathbf{y}^{(N')})$. This formula is obtained by the insertion of the fermionic operator $res_z : \psi^*(z) z \frac{d}{dz} \psi(z) :$ inside the fermionic vacuum expectation.

These formulae can be also written in terms of higher KP and TL times, with the help of vertex operator action, see the *Subsection "Vertex operator action"*. Then the relations (6.5.19) are the infinitesimal version of (6.6.7), while the relation (6.5.22) is the infinitesimal version of (6.6.8).

In some examples below we shall take specify $\mathbf{t}^*$ as follows. (1) Let us take

$$m t_m = \sum_{k=1}^{N} x_k^m, \quad \mathbf{t}^* = (t_1^*, 0, 0, \dots) \tag{6.5.23}$$

One notes that that the tau-function $\tau_r(\mathbf{t}, \mathbf{t}^*)$ will not change if instead of (6.5.15) one take

$$m t_m = \sum_{k=1}^{N} (t_1 x_k)^m, \quad \mathbf{t}^* = (1, 0, 0, \dots) \tag{6.5.24}$$

Then one consider the equation (6.5.19) with $k = 1$, taking $t_1 = 1$. We put $D_i = x_i \partial_{x_i}$ and get

$$\left(\sum_{i=1}^{N} D_i - \frac{1}{\Delta} \left(\sum_{i=1}^{N} x_i r(D_i) \right) \Delta \right) \tau_r(\mathbf{t}(\mathbf{x}^N), \mathbf{t}^*) = 0 \tag{6.5.25}$$

where r is given by (6.12.1), and

$$\Delta = \frac{\prod_{i<j}(x_i - x_j)}{(x_1 \cdots x_N)^{N-M-1}} \tag{6.5.26}$$

6.6 The vertex operator action. Linear equations for tau-functions II

Now we present relations between hypergeometric functions which follow from the soliton theory, for instance see [13],[34]. (In [12] there were few misprints in the formulae presented below). Let us introduce the operators which act on functions of $\mathbf{t}$ variables:

$$\Omega_r^{(\infty)}(\mathbf{t}, \mathbf{t}^*) := -\lim_{\epsilon \to 0} \mathrm{res}_{z=0} V_\infty^*(\mathbf{t}, z+\epsilon) \xi_r(\mathbf{t}^*, z^{-1}) V_\infty(\mathbf{t}, z) \frac{dz}{z}, \tag{6.6.1}$$

$$\Omega_r^{(0)}(\mathbf{t}^*,\mathbf{t}) := \lim_{\epsilon\to 0}\mathrm{res}_{z=0}V_0^*(\mathbf{t}^*,z+\epsilon)\xi_r^{(0)}(\mathbf{t},z)V_0(\mathbf{t}^*,z)\frac{dz}{z}, \qquad (6.6.2)$$

where $V_\infty(\mathbf{t},z), V_\infty^*(\mathbf{t},z), V_0(\mathbf{t}^*,z), V_0^*(\mathbf{t}^*,z)$ are defined by (6.1.1), and

$$\xi_r(\mathbf{t}^*,z^{-1}) = \sum_{m=1}^{+\infty} t_m^*\left(\frac{1}{z}r(D)\right)^m, \quad \xi_r^{(0)}(\mathbf{t},z) = \sum_{m=1}^{+\infty} t_m\left(r(D)z\right)^m \qquad (6.6.3)$$

For instance

$$r=1: \qquad \Omega_r^{(\infty)}(\mathbf{t},\mathbf{t}^*) = \Omega_r^{(0)}(\mathbf{t}^*,\mathbf{t}) = \sum_{n>0} nt_nt_n^*. \qquad (6.6.4)$$

Also we consider

$$Z_{nn}(\mathbf{t}) = \mathrm{res}_{z=0}\mathrm{res}_{z^*=0}\frac{z^n}{z^{*n}}V_\infty^*(\mathbf{t},z^*)V_\infty(\mathbf{t},z)\frac{dzdz^*}{zz^*}, \qquad (6.6.5)$$

$$Z_{nn}^*(\mathbf{t}^*) = \mathrm{res}_{z=0}\mathrm{res}_{z^*=0}\frac{z^n}{z^{*n}}V_0^*(\mathbf{t}^*,z^*)V_0(\mathbf{t}^*,z)\frac{dzdz^*}{zz^*}. \qquad (6.6.6)$$

The bosonization formulae (6.1.3),(6.1.4) result in

Proposition 12 *We have shift argument formulae for the tau function (3.4.8):*

$$e^{\Omega_r^{(\infty)}(\mathbf{t},\gamma)}\cdot\tau_r(M,\mathbf{t},\mathbf{t}^*) = \tau_r(M,\mathbf{t},\mathbf{t}^*+\gamma), \quad e^{\Omega_r^{(0)}(\mathbf{t}^*,\gamma)}\cdot\tau_r(M,\mathbf{t},\mathbf{t}^*) = \tau_r(M,\mathbf{t}+\gamma,\mathbf{t}^*). \quad (6.6.7)$$

Also we have

$$e^{\sum_{-\infty}^{\infty}\gamma_{n+M}Z_{nn}(\mathbf{t})}\cdot\tau(M,\mathbf{t},T,\mathbf{t}^*) = e^{\sum_{-\infty}^{\infty}\gamma_{n+M}Z_{nn}^*(\mathbf{t}^*)}\cdot\tau(M,\mathbf{t},T,\mathbf{t}^*) = \tau(M,\mathbf{t},T+\gamma,\mathbf{t}^*), \quad (6.6.8)$$

For instance

$$e^{\Omega_r^{(\infty)}(\mathbf{t},\mathbf{t}^*)}\cdot 1 = e^{\Omega_r^{(0)}(\mathbf{t}^*,\mathbf{t})}\cdot 1 = \tau_r(M,\mathbf{t},\mathbf{t}^*) \qquad (6.6.9)$$

Also

$$e^{\sum_{-\infty}^{\infty}T_{n+M}Z_{nn}}\cdot\exp\left(\sum_{n=1}^{\infty}nt_nt_n^*\right) = e^{\sum_{-\infty}^{\infty}T_{n+M}Z_{nn}^*}\cdot\exp\left(\sum_{n=1}^{\infty}nt_nt_n^*\right) = \tau(M,\mathbf{t},T,\mathbf{t}^*). \quad (6.6.10)$$

6.7 Determinant formulae I (Hirota-Miwa variables) [12]

In the case $\mathbf{t}=\mathbf{t}(\mathbf{x}^N)$ one can apply Wick theorem [23] to obtain a determinant formulae.

Proposition 13 *A generalization of Milne's determinant formula*

$$\tau_r(M,\mathbf{t}(\mathbf{x}^N),\mathbf{t}^*) = \frac{\det\left(x_i^{N-k}\tau_r(M-k+1,\mathbf{t}(x_i),\mathbf{t}^*)\right)_{i,k=1}^N}{\det\left(x_i^{N-k}\right)_{i,k=1}^N}. \qquad (6.7.1)$$

Proof follows from the representation (6.5.1):

$$\tau_r(M,\mathbf{t}(\mathbf{x}^N),\mathbf{t}^*) = \Delta^{-1}\left(e^{\xi_{r'}(\mathbf{t}^*,x_1)}\cdots e^{\xi_{r'}(\mathbf{t}^*,x_N)}\cdot\Delta\right), \quad \Delta = \frac{\prod_{i<j}(x_i-x_j)}{(x_1\cdots x_N)^{N-1-M}}, \qquad (6.7.2)$$

from the explicit form of the Vandermond determinant: $\prod_{i<j}(x_i-x_j) = \det(x_i^{N-k})$ and from the representation (6.7.2) for the case when the number of variables x is equal to one as:

$$\tau_r(M-k+1,\mathbf{t}(x_i),\mathbf{t}^*) = e^{\xi_{r'}(\mathbf{t}^*,x_i)}\cdot x_i^{M-k+1} \qquad (6.7.3)$$

where $\mathbf{t}(x_i) = (x_i,\frac{x_i^2}{2},\frac{x_i^3}{3},\ldots)$.

In the case $\mathbf{t}=\mathbf{t}(\mathbf{x}^N), \mathbf{t}^*=\mathbf{t}^*(\mathbf{y}^N)$ we apply to (6.2.16) the Wick theorem to obtain a different Proposition

Proposition 14 *For $r \neq 0$ we take a tau function $\tau_r(M, \mathbf{t}(\mathbf{x}^N), \mathbf{t}^*(\mathbf{y}^N))$ and apply Wick's theorem. We get the determinant formula:*

$$\tau_r(M, \mathbf{t}(\mathbf{x}^N), \mathbf{t}^*(\mathbf{y}^N)) = \frac{c_M}{c_{M-N}} \frac{\det\left(\tau_r(M-N+1, x_i, y_j)\right)_{i,j=1}^N}{\Delta(\mathbf{x}^N)\Delta(\mathbf{y}^N)}, \tag{6.7.4}$$

where

$$c_n = \prod_{k=0}^{n-1} (r(k))^{k-n} \tag{6.7.5}$$

$$\tau_r(M-N+1, x_i, y_j) = 1 + r(M-N+1)x_i y_j + r(M-N+1)r(M-N+2)x_i^2 y_j^2 + \cdots \tag{6.7.6}$$

One can prove it also either from the representation (6.5.6) or from the representation (6.5.7).

6.8 Determinant formulae II (The case $\mathbf{r(0) = 0}$.)

There is also determinant formulae for tau functions in variables $\mathbf{t}, \mathbf{t}^*$. These one we get in case the function r has zeroes. If $r(0) = 0$ and $n > 0$ then

$$\tau_r(n, \mathbf{t}, \mathbf{t}^*) = c_n \det\left(\frac{\partial^{a+b}}{\partial t_1^a \partial t_1^{*b}} \tau_r(1, \mathbf{t}, \mathbf{t}^*)\right)_{a,b=0}^{n-1} \tag{6.8.1}$$

$$c_n = \prod_{k=0}^{n-1} (r(k))^{k-n} \tag{6.8.2}$$

6.9 Integral representations I (Hirota-Miwa variables) [12]

In the case $\mathbf{t} = \mathbf{t}(\mathbf{x}^N), \mathbf{t}^* = \mathbf{t}^*(\mathbf{y}^N)$ we get an integral representation formulae. For the fermions (6.2.5) we easily get the relations:

$$\int \psi(T, \alpha z) d\mu(\alpha) = \psi\left(T + T(\mu), z\right), \quad \int \psi^*\left(-T, \frac{1}{\alpha z}\right) d\tilde{\mu}(\alpha) = \psi^*\left(-T - T(\tilde{\mu}), \frac{1}{z}\right) \tag{6.9.1}$$

where $\mu, \tilde{\mu}$ are some integration measures, and shifts of times T_n are defined in terms of the moments:

$$\int \alpha^n d\mu(\alpha) = e^{-T_n(\mu)}, \quad \int \alpha^n d\tilde{\mu}(\alpha) = e^{-T_n(\tilde{\mu})}. \tag{6.9.2}$$

Therefore thanks to the bosonization formulae (6.2.16) we have the relations for the tau-function; below $\mathbf{t}^*$ is defined via Hirota-Miwa variables.

Proposition 15 *Integral representation formula holds*

$$\int \tilde{\Delta}_{\tilde{T}}(\tilde{\alpha}\mathbf{x}^N) \frac{\tau\left(M, \mathbf{t}(\tilde{\alpha}\mathbf{x}^N), T + \tilde{T}, \mathbf{t}(\alpha \mathbf{y}_{(N)})\right)}{\tau\left(M, \mathbf{0}, T + \tilde{T}, \mathbf{0}\right)} \Delta_T(\alpha \mathbf{y}_{(N)}) \prod_{i=1}^N d\tilde{\mu}(\tilde{\alpha}_i) \prod_{i=1}^N d\mu(\alpha_i)$$

$$= \tilde{\Delta}_{\tilde{T}+\tilde{T}(\tilde{\mu})}(\mathbf{x}^N) \frac{\tau\left(M, \mathbf{t}(\mathbf{x}^N), T + \tilde{T} + \tilde{T}(\tilde{\mu}) + T(\mu), \mathbf{t}(\mathbf{y}^N)\right)}{\tau\left(M, \mathbf{0}, T + \tilde{T} + \tilde{T}(\tilde{\mu}) + T(\mu), \mathbf{0}\right)} \Delta_{T+T(\mu)}(\mathbf{y}^N). \tag{6.9.3}$$

where $\Delta_T(\alpha\mathbf{y}_{(N)}) = \Delta^+(M,N,T,\alpha\mathbf{y}_{(N)})$*,* $\tilde{\Delta}_{\tilde{T}}(\tilde{\alpha}\mathbf{x}^N) = \tilde{\Delta}^+(M,N,\tilde{T},\tilde{\alpha}\mathbf{x}^N)$*,* $\alpha\mathbf{y}_{(N)} = (\alpha_1 y_1, \alpha_2 y_2, \dots, \alpha_N y_N)$ *and* $\tilde{\alpha}\mathbf{x}^N = (\tilde{\alpha}_1 x_1, \tilde{\alpha}_2 x_2, \dots, \tilde{\alpha}_N x_N)$*. In particular*

$$\int \frac{\tau\left(M,\mathbf{t},T,\mathbf{t}(\alpha\mathbf{y}_{(N)})\right)}{\tau\left(M,\mathbf{0},T,\mathbf{0}\right)}\Delta_T(\alpha\mathbf{y}_{(N)})\prod_{i=1}^N d\mu(\alpha_i) = \frac{\tau\left(M,\mathbf{t},T+T(\mu),\mathbf{t}(\mathbf{y}^N)\right)}{\tau\left(M,\mathbf{0},T+T(\mu),\mathbf{0}\right)}\Delta_{T+T(\mu)}(\mathbf{y}^N). \tag{6.9.4}$$

Remember that arbitrary linear combination of tau-functions is not a tau-function. Formulae (6.9.3) and also (6.9.4) give the integral representations for the tau-function (3.4.6). It may help to express a tau-function with the help of a more simple one.

We shall use the following integrals

$$\frac{1}{2\pi\sqrt{-1}}\int_C \psi^*(-T,\frac{\alpha}{x})e^{\alpha}\alpha^{-b}d\alpha = \psi^*(-T-T^b,\frac{1}{x}), \tag{6.9.5}$$

$$\int_0^\infty \psi^*(-T,\frac{1}{\alpha x})e^{-\alpha}\alpha^{a-1}d\alpha = \psi^*(-T-T^a,\frac{1}{x}), \tag{6.9.6}$$

$$\frac{1}{\Gamma(b-a)}\int_0^1 \psi^*(-T,\frac{1}{\alpha x})\alpha^{a-1}(1-\alpha)^{b-a-1}d\alpha = \psi^*(-T-T^c,\frac{1}{x}), \tag{6.9.7}$$

where C in (6.9.5) starts at $-\infty$ on the real axis, circles the origin in the counterclockwise direction and returns to the starting point, and

$$T_n^b = \ln\Gamma(b+n+1), \quad T_n^a = -\ln\Gamma(a+n+1), \quad T_n^c = \ln\frac{\Gamma(b+n+1)}{\Gamma(a+n+1)\Gamma(b-a)}. \tag{6.9.8}$$

Let us remind that q-versions of exponential function, of gamma function and of beta function are known as follows [25]

$$\exp_q(x) = \frac{1}{(x(1-q);q)_\infty}, \quad \Gamma_q(x) = (1-q)^{1-a}\frac{(q;q)_\infty}{(q^x;q)_\infty}, \quad B_q(x,y) = \frac{\Gamma_q(x)\Gamma_q(y)}{\Gamma_q(x+y)} \tag{6.9.9}$$

Gamma and beta functions have the q-integral representations

$$\Gamma_q(x) = -q^{-x}\int_0^\infty \frac{\alpha^{x-1}}{\exp_q(\alpha)}d_q\alpha, \quad B_q(x,y) = \int_0^1 \alpha^{x-1}\frac{(\alpha q;q)_\infty}{(\alpha q^y;q)_\infty}d_q\alpha \tag{6.9.10}$$

where q-integral are defined as

$$\int_0^\infty f(x)d_qx = (1-q)\sum_{j=-\infty}^{j=\infty} q^j f(q^j), \quad \int_0^c f(x)d_qx = c(1-q)\sum_{j=0}^{j=\infty} q^j f(cq^j) \tag{6.9.11}$$

The following q-integrals will be of use:

$$\frac{q^{-a}}{q-1}\int_0^\infty \psi^*(-T,\frac{q}{\alpha(1-q)x})\frac{\alpha^{a-1}}{\exp_q(\alpha)}d_q\alpha = \psi^*\left(-T-T(a,q),\frac{1}{x}\right), \tag{6.9.12}$$

$$\frac{1}{\Gamma_q(b-a)}\int_0^1 \psi^*(-T,\frac{1}{\alpha x})\alpha^{a-1}\frac{(\alpha q;q)_\infty}{(\alpha q^{b-a};q)_\infty}d_q\alpha = \psi^*(-T-T(a,b,q),\frac{1}{x}). \tag{6.9.13}$$

where, as one can check using (6.9.10),

$$T_n(a,q) = \ln \frac{1}{(1-q)^n \Gamma_q(a+n+1)}, \quad T_n(a,b,q) = \ln \frac{\Gamma_q(b+n+1)}{\Gamma_q(a+n+1)}. \tag{6.9.14}$$

In the same way one can consider Hirota-Miwa variables (1.0.8). In the *Examples* below we shall present hypergeometric functions listed in the *Subsections 1.2 and 1.3* as tau-functions of the type (6.2.16). Then we are able to write down integration formulae, namely (6.9.4), which express ${}_{p+1}\Phi_s$ and ${}_{p+1}\Phi_{s+1}$ in terms of ${}_p\Phi_s$ with the help of (6.9.12), (6.9.13) and (6.9.3). In [38] different integral representation formula was presented, which was based on the q-analog of Selberg's integral of Askey and Kadell. By taking the limit $q \to 1$ one can consider functions ${}_pF_s$. Using (6.9.8), one can express ${}_{p+1}\mathcal{F}_s$, ${}_{p+1}\mathcal{F}_{s+1}$ and ${}_p\mathcal{F}_{s+1}$ as integrals of ${}_p\mathcal{F}_s$ with the help of (6.9.6), (6.9.7) and (6.9.5) respectively.

6.10 Integral representations II

Now let us consider the case when τ_r depends on $T, \mathbf{t}^*$. Using (3.4.7) and (5.1.21) one gets the integral representation for τ_r in case the function r has zero. Let us again choose $r(0) = 0$. The have the representation described above

Proposition 16

$$\tau_r(M, \mathbf{t}, \mathbf{t}^*) = \int \cdots \int \Delta(z)\Delta(z^*) \prod_{k=1}^{M} e^{\sum_{n=1}^{\infty}(z_k^n t_n + z_k^{*n} t_n^*)} \mu_r(z_k {z^*}_k) dz_k dz_k^* \tag{6.10.1}$$

where

$$\Delta(z) = \prod_{i<j}^{M}(z_i - z_j), \quad \Delta(z^*) = \prod_{i<j}^{M}(z_i^* - z_j^*) \tag{6.10.2}$$

and μ_r is defined by (5.1.10),(5.1.2).

Now it is interesting to notice that if the series $\tau_{1/r}$ (which defines μ_r, see (5.1.10)) is a convergent series then the series τ_r may be divergent one. In this case one can consider the integral $I_r(M, \mathbf{t}, \mathbf{t}^*)$ is an analog of Borel sum for the divergent series τ_r, see [2].

6.11 Rational r and their q-deformations

Lemma 1 If

$$r(n) = \frac{\prod_{i=1}^{p}(n + a_i)}{\prod_{i=1}^{s}(n + b_i)} \tag{6.11.1}$$

then

$$r_\lambda(n) = (s_\lambda(\gamma(+\infty)))^{s-p} \frac{\prod_{k=1}^{p} s_\lambda(\gamma(a_k + n))}{\prod_{k=1}^{s} s_\lambda(\gamma(b_k + n))} \tag{6.11.2}$$

where

$$\gamma(\infty) = (1, 0, 0, \ldots), \quad \gamma(a) = \left(\frac{a}{1}, \frac{a}{2}, \frac{a}{3}, \ldots\right) \tag{6.11.3}$$

and $s_\lambda(\gamma)$ are Schur functions as functions of variables γ according to (2.1.9) and (2.1.12).□

Lemma 2 If

$$r(n) = \frac{\prod_{i=1}^{p}(1 + q^{n+a_i})}{\prod_{i=1}^{s}(1 + q^{n+b_i})} \tag{6.11.4}$$

then

$$r_\lambda(n) = (s_\lambda(\gamma(+\infty,q)))^{s-p}\,\frac{\prod_{k=1}^{p} s_\lambda(\gamma(a_k+n,q))}{\prod_{k=1}^{s} s_\lambda(\gamma(b_k+n,q))} \tag{6.11.5}$$

where

$$\gamma_m(a,q) = \frac{1-(q^a)^m}{m(1-q^m)}, \quad \gamma(+\infty,q) = \frac{1}{m(1-q^m)}, \quad m=1,2,\dots \tag{6.11.6}$$

□ For the proof we use the following relations (see [4]):

$$(q^a;q)_\lambda = \prod_{(i,j)\in\lambda}(1-q^{a+j-i}) = \frac{s_\lambda(\gamma(a,q))}{s_\lambda(\gamma(+\infty,q))}, \quad (a)_\lambda = \prod_{(i,j)\in\lambda}(a+j-i) = \frac{s_\lambda(\gamma(a))}{s_\lambda(\gamma(+\infty)}, \tag{6.11.7}$$

Now we see that for arbitrary complex b we have

Proposition 17

$$< e^{\gamma_1}, e^{\sum_{m=1}^\infty m\gamma_m t_m^*} >_{r,n} = < e^{(b+n)\sum_{m=1}^\infty m\gamma_m}, e^{\sum_{m=1}^\infty m\gamma_m t_m^*} >_{r_b,n}, \quad r_b(k) := \frac{r(k)}{b+k}. \tag{6.11.8}$$

Also we have

$$< e^{\sum_{m=1}^\infty \frac{\gamma_m}{1-q^m}}, e^{\sum_{m=1}^\infty m\gamma_m t_m^*} >_{r,n} = < e^{\sum_{m=1}^\infty \frac{\gamma_m(1-q^{b+n+m})}{1-q^m}}, e^{\sum_{m=1}^\infty m\gamma_m t_m^*} >_{r_b,n}, \quad r_b(k) := \frac{r(k)}{1-q^{b+k}}. \tag{6.11.9}$$

Let us note that parameters $\gamma_m(a,q)$ and $\gamma_m(a)$ are chosen via 'generalized Hirota-Miwa transforms' (6.11.6),(6.11.3) with 'multiplicity a'. Remember that $|q|<1$. Due to

$$s_\lambda(\gamma(+\infty,q)) = \lim_{a\to+\infty} s_\lambda(\gamma(a,q)) = \frac{q^{n(\lambda)}}{H_\lambda(q)}, \tag{6.11.10}$$

$$s_\lambda(\gamma(+\infty)) = \lim_{a\to+\infty} s_\lambda\left(\frac{\gamma_1(a)}{a}, \frac{\gamma_2(a)}{a^2}, \dots\right) = \lim_{a\to+\infty}\frac{1}{a^{|\lambda|}} s_\lambda(\gamma(a)) = \frac{1}{H_\lambda}. \tag{6.11.11}$$

This allows to rewrite the series (6.12.7) and (6.12.25) only in terms of Schur functions, see [12].

For functions ${}_pF_s$ of (6.12.7),(6.12.9) and ${}_p\Phi_s$ of (6.12.25),(6.12.26) we use the results of subsection 5.4 to derive the following representations

$${}_pF_s\left(\left.\begin{matrix} a_1+M,\dots,a_p+M \\ b_1+M,\dots,b_s+M\end{matrix}\right| \mathbf{x}^N\right) = \frac{1}{\Delta(x)} e^{x_1 r(D_1)+\cdots+x_N r(D_N)}\cdot\Delta(x) \tag{6.11.12}$$

$$= \frac{1}{\Delta(x)}\prod_{i=1}^N\left(1 - x_i\frac{r(D_i)}{b+D_i}\right)^{-b}\cdot\Delta(x) \tag{6.11.13}$$

$${}_p\Phi_s\left(\left.\begin{matrix} a_1+M,\dots,a_p+M \\ b_1+M,\dots,b_s+M\end{matrix}\right| q,\mathbf{x}^N\right) = \frac{1}{\Delta(x)}\prod_{i=1}^N \frac{1}{(x_i r(D_i);q)_\infty}\cdot\Delta(x) \tag{6.11.14}$$

$$= \frac{1}{\Delta(x)}\prod_{i=1}^N \frac{\left(x_i q^b r(D_i)(1-q^{b+D_i})^{-1};q\right)_\infty}{(x_i r(D_i);q)_\infty}\cdot\Delta(x) \tag{6.11.15}$$

where $D_i = x_i\frac{\partial}{\partial x_i}$, $\Delta(x)$ is defined in (6.5.8), the notation $(b;q)_\infty$ see in (8.0.14).

6.12 Examples of hypergeometric series [12]

The main point of this subsection is the observation that if $r(D)$ is a rational function of D then τ_r is a hypergeometric series. If $r(D)$ is a rational function of q^D we obtain q-deformed hypergeometric series. Concerning the hypergeometric functions see Appendix C. Now let us consider various $r(D)$.

Example 1 Let all parameters b_k be non integers.

$$ {}_p r_s(D) = \frac{(D+a_1)(D+a_2)\cdots(D+a_p)}{(D+b_1)(D+b_2)\cdots(D+b_s)}. \tag{6.12.1} $$

For the vacuum expectation value (3.4.8) we have:

$$ {}^p\tau_r^s(M,\mathbf{t},\mathbf{t}^*) = \sum_\lambda s_\lambda(\mathbf{t})s_\lambda(\mathbf{t}^*)\frac{(a_1+M)_\lambda\cdots(a_p+M)_\lambda}{(b_1+M)_\lambda\cdots(b_s+M)_\lambda}. \tag{6.12.2} $$

Let us notice that if we put

$$ t_1^* = 1, \qquad t_i^* = 0, \quad i > 1, \tag{6.12.3} $$

then $s_\lambda(\mathbf{t}^*) = H_\lambda^{-1}$ (see (6.11.11)), where H_λ is the following hook product:

$$ H_\lambda = \prod_{(i,j)\in\lambda} h_{ij}, \qquad h_{ij} = (\lambda_i + \lambda_j' - i - j + 1). \tag{6.12.4} $$

Thus for $\mathbf{t}^*$ as in (6.12.3) the tau function (6.12.2) yields the hypergeometric function related to Schur functions as follows:

$$ {}^p\tau_r^s(M,\mathbf{t},\mathbf{t}^*) = {}_pF_s\left(\begin{matrix} a_1+M,\ldots,a_p+M \\ b_1+M,\ldots,b_s+M\end{matrix}\middle| t_1,t_2,\ldots\right) = $$
$$ \sum_\lambda \frac{(a_1+M)_\lambda\cdots(a_p+M)_\lambda}{(b_1+M)_\lambda\cdots(b_s+M)_\lambda}\frac{s_\lambda(\mathbf{t})}{H_\lambda}. \tag{6.12.5} $$

As an example taking $a_1 = 0$, $\mathbf{t} = (t_1, 0, 0, \ldots)$, $\mathbf{t}^* = (t_1^*, 0, 0, \ldots)$ we obtain the well-known ordinary hypergeometric function of one variable:

$$ {}_{p-1}F_s(a_2\pm 1,\ldots,a_p\pm 1; b_1\pm 1,\ldots,b_s\pm 1; \pm t_1t_1^*) = \tau_r(\pm 1,\mathbf{t},\mathbf{t}^*), \quad r(n) = n\frac{\prod_{i=2}^p(a_i+n)}{\prod_{i=1}^s(b_i+n)} \tag{6.12.6} $$

Now we take $\mathbf{t} = \mathbf{t}(\mathbf{x}^N), \mathbf{t}^* = (1,0,0,\ldots)$ then the (6.12.5) turns out to be the known hypergeometric function of matrix argument $\mathbf{X}$ [39],[26]:

$$ {}_pF_s\left(\begin{matrix} a_1+M,\ldots,a_p+M \\ b_1+M,\ldots,b_s+M\end{matrix}\middle| \mathbf{X}\right) = {}^p\tau_r^s(M,\mathbf{t}(\mathbf{x}^N),\mathbf{t}^*) = $$
$$ \sum_{\substack{\lambda \\ l(\lambda)\le N}} \frac{(a_1+M)_\lambda\cdots(a_p+M)_\lambda}{(b_1+M)_\lambda\cdots(b_s+M)_\lambda}\frac{s_\lambda(\mathbf{x}^N)}{H_\lambda}. \tag{6.12.7} $$

This function depends only on the eigenvalues of the matrix $\mathbf{X}$ which are $\mathbf{x}^N = (x_1, \ldots, x_N)$. The hypergeometric function of matrix argument is also known as the hypergeometric function

(8.0.26) related to zonal polynomials for the symmetric space $GL(N,C)/U(N)$ [26],[4]. Here $x_i = z_i^{-1}, i = 1, ..., N$ are the eigenvalues of the matrix $\mathbf{X}$, and for zonal spherical polynomials there is the following matrix integral representation

$$Z_\lambda(\mathbf{X}) = Z_\lambda(\mathbf{I}_N) \int_{U(N,C)} \Delta^\lambda \left(U^* \mathbf{X} U\right) d_* U, \tag{6.12.8}$$

where $\Delta^\lambda\left(\mathbf{X}\right) = \Delta_1^{\lambda_1-\lambda_2} \Delta_2^{\lambda_2-\lambda_3} \cdots \Delta_N^{\lambda_N}$ and $\Delta_1, \ldots \Delta_N$ are main minors of the matrix $\mathbf{X}$, $d_* U$ is the invariant measure on $U(N,C)$, see [26]. In our case $Z_\lambda(\mathbf{X})$ coincides with the Schur function $s_\lambda(\mathbf{x}^N)$.

Due to (6.5.1) hypergeometric function (6.12.5) has the representation as follows

$${}_pF_s\left(\left.\begin{matrix} a_1+M, \ldots, a_p+M \\ b_1+M, \ldots, b_s+M \end{matrix}\right| \mathbf{x}^N\right) = \frac{1}{\Delta} \exp(x_1 r(D_1)) \cdots \exp(x_N r(D_N)) \cdot \Delta, \tag{6.12.9}$$

where $D_i = x_i \frac{\partial}{\partial x_i}$ and

$$\Delta = \frac{\prod_{i<j}(x_i - x_j)}{(x_1 \cdots x_N)^{N-M-1}} \tag{6.12.10}$$

(See also (6.11.13) for different representations).

Now let us take $\mathbf{t}^* = (t_1^*, 0, 0, \ldots)$. Then we get the same function (6.12.7), but with each x_i changed by $-t_1^* x_i$. Then one consider the equation (6.5.19) with $k = 1$. We put $D_i = x_i \partial_{x_i}$ and get

$$\left(\sum_{i=1}^N D_i - \frac{1}{\Delta}\left(\sum_{i=1}^N x_i r(D_i)\right) \Delta\right) {}_pF_s\left(\left.\begin{matrix} a_1+M, \ldots, a_p+M \\ b_1+M, \ldots, b_s+M \end{matrix}\right| \mathbf{x}^N\right) = 0 \tag{6.12.11}$$

where r is given by (6.12.1), and Δ see in (6.12.10).

Taking $N = 1, \mathbf{t}^* = (1, 0, 0, \ldots)$ we obtain the ordinary hypergeometric function of one variable $x = x_1$, which is (compare with (6.12.6)):

$${}_pF_s\left(a_1+M, \ldots, a_p+M; b_1+M, \ldots, b_s+M; x\right) = x^{-M} e^{x r(D)} \cdot x^M, \quad D = x\frac{d}{dx} \tag{6.12.12}$$

The ordinary hypergeometric series satisfies the known hypergeometric equation (we put $M = 0$) written in the following form:

$$\left(\partial_x - {}_pr_s(D)\right) {}_pF_s(a_1, \ldots, a_p; b_1, \ldots, b_s; x) = 0, \quad D := x\partial_x. \tag{6.12.13}$$

This relation helps us to understand a meaning of the function r.
It is known that the series (6.12.7) diverges if $p > s+1$ (until any of $a_i + M$ is non positive integer). In case $p = s+1$ it converges in certain domain in the vicinity of $\mathbf{x}^N = \mathbf{0}$. For $p < s+1$ the series (6.12.7) converges for all $\mathbf{x}^N$. These known facts (see [25]) can be also obtained with the help of the determinant representation (6.7.1) and properties of (8.0.10).

Now we have the following representations:

$${}_{p+1}F_s\left(\left.\begin{matrix} a, a_1, \ldots, a_p \\ b_1, \ldots, b_s \end{matrix}\right| x_1, \ldots, x_N\right) = \frac{1}{\prod_{i<j}^N (x_i - x_j)} \int_0^\infty \cdots \int_0^\infty \tag{6.12.14}$$

$$(\alpha_1 \cdots \alpha_N)^{a-N} e^{-\alpha_1 - \cdots - \alpha_N} {}_pF_s\left(\left.\begin{matrix} a_1, \ldots, a_p \\ b_1, \ldots, b_s \end{matrix}\right| \alpha_1 x_1, \ldots, \alpha_N x_N\right) \prod_{i<j}^N (\alpha_i x_i - \alpha_j x_j) d\alpha_1 \cdots d\alpha_N$$

Hankel type of representation

$$
{}_pF_{s+1}\left(\begin{matrix}a_1,\dots,a_p\\ b,b_1,\dots,b_s\end{matrix}\middle| x_1,\dots,x_N\right)=\frac{1}{\prod_{i<j}^N(x_i-x_j)}\frac{1}{(2\pi\sqrt{-1})^N}\int_C\cdots\int_C \tag{6.12.15}
$$

$$
(\alpha_1\cdots\alpha_N)^{1-N-b}e^{\alpha_1+\cdots+\alpha_N}{}_pF_s\left(\begin{matrix}a_1,\dots,a_p\\ b_1,\dots,b_s\end{matrix}\middle|\frac{x_1}{\alpha_1},\dots,\frac{x_N}{\alpha_N}\right)\prod_{i<j}^N(\alpha_i x_i-\alpha_j x_j)d\alpha_1\cdots d\alpha_N
$$

where C starts at $-\infty$ on the real axis, circles the origin in the counterclockwise direction and returns to the starting point.

Beta type representation:

$$
{}_{p+1}F_{s+1}\left(\begin{matrix}a,a_1,\dots,a_p\\ b,b_1,\dots,b_s\end{matrix}\middle| x_1,\dots,x_N\right)=\frac{1}{\prod_{i<j}^N(x_i-x_j)}\left(\frac{1}{\Gamma(b-a)}\right)^N\int_0^1\cdots\int_0^1 \tag{6.12.16}
$$

$$
{}_pF_s\left(\begin{matrix}a_1,\dots,a_p\\ b_1,\dots,b_s\end{matrix}\middle|\alpha_1x_1,\dots,\alpha_Nx_N\right)\prod_{i<j}^N(\alpha_ix_i-\alpha_jx_j)\prod_{i=1}^N\alpha_i^{a-N}(1-\alpha_i)^{b-a-1}d\alpha_i
$$

Let us notice that according to [14] the logarithmic derivative of the hypergeometric function (6.12.5) $u(x)=\frac{d}{dx}\log{}_2F_1\left(\begin{smallmatrix}p,q\\ n\end{smallmatrix}\middle| t_1=x,0,0,\dots\right)$ is a solution of the Painleve V equation.

Example 2. Let us consider N by N matrices $\mathbf{X},\mathbf{Y}$ with eigenvalues $\mathbf{x}^N=(x_1,\dots,x_N),\mathbf{y}^N=(y_1,\dots,y_N)$ respectively. In order to get a hypergeometric function of two matrix variables $\mathbf{X},\mathbf{Y}$, see (8.0.26) in the Appendix C, we put

$$
{}_pr_s(n)=\frac{\prod_{i=1}^p(a_i+n)}{\prod_{i=1}^s(b_i+n)}\frac{1}{N-M+n}, \tag{6.12.17}
$$

Notice that r now explicitly depends on N.

Taking into account the relations (6.11.7) and (6.11.11) we get

$$
(N)_\lambda=H_\lambda s_\lambda(\mathbf{t}(\mathbf{1}^N)),\quad \mathbf{1}^N=(1,1,\dots,1) \tag{6.12.18}
$$

and putting $\mathbf{t}=\mathbf{t}(\mathbf{x}^N)$ and $\mathbf{t}^*=\mathbf{t}^*(\mathbf{y}^N)$ we obtain the hypergeometric function of (8.0.26):

$$
\langle M|e^{H(\mathbf{t}(\mathbf{x}^N))}e^{-A(\mathbf{t}^*(\mathbf{y}^N))}|M\rangle={}_p\mathcal{F}_s\left(\begin{matrix}a_1+M,\dots,a_p+M\\ b_1+M,\dots,b_s+M\end{matrix}\middle|\mathbf{X},\mathbf{Y}\right)=
$$

$$
\sum_{\substack{\lambda\\ l(\lambda)\le N}}\frac{s_\lambda(\mathbf{x}^N)s_\lambda(\mathbf{y}^N)}{(N)_\lambda}\frac{(a_1+M)_\lambda\cdots(a_p+M)_\lambda}{(b_1+M)_\lambda\cdots(b_s+M)_\lambda}. \tag{6.12.19}
$$

Due to (6.5.8) hypergeometric function (6.12.19) has the representation as follows ($r={}_pr_s$):

$$
{}_p\mathcal{F}_s\left(\begin{matrix}a_1+M,\dots,a_p+M\\ b_1+M,\dots,b_s+M\end{matrix}\middle|\mathbf{x}^N,\mathbf{y}^N\right)=\frac{1}{\Delta}\prod_{i,j=1}^N(1-y_jx_ir(D_{x_i}))^{-1}\cdot\Delta \tag{6.12.20}
$$

Here and below the inverse operator $(1-y_jx_ir(D_{x_i}))^{-1}$ is a formal series $1+y_jx_ir(D_{x_i})+\cdots$.

For example when $N=1$ we get the ordinary hypergeometric function of one variable $xy=x_1y_1$':

$$
{}_pF_s\left(a_1+M,\dots,a_p+M;b_1+M,\dots,b_s+M;xy\right)=x^{-M}\left(1-xyr(D_x)\right)^{-1}\cdot x^M \tag{6.12.21}
$$

Looking at (6.12.9) one derives the following linear equation

$$\left(D_{y_j}-\sum_{i=1}^{N}\frac{1}{1-y_jx_ir(D_{x_i})}+N\right)\cdot\left(\Delta_p\mathcal{F}_s\left(\begin{matrix}a_1+M,\dots,a_p+M\\ b_1+M,\dots,b_s+M\end{matrix}\middle|\mathbf{x}^N,\mathbf{y}^N\right)\right)=0 \qquad (6.12.22)$$

Example 3 The q-generalization of the *Example 3* is Milne's hypergeometric function of the single set of variables. To get it we choose

$$ {}_pr_s^{(q)}(n)=\frac{\prod_{i=1}^{p}(1-q^{a_i+n})}{\prod_{i=1}^{s}(1-q^{b_i+n})}. \qquad (6.12.23)$$

We take $\mathbf{t}=\mathbf{t}(\mathbf{x}^N),\mathbf{t}^*=\mathbf{t}^*(\mathbf{y}^\infty)$. By putting

$$y_k=q^{k-1},\quad k=1,2,...,\quad t_m^*=\sum_{k=1}^{+\infty}\frac{y_k^m}{m}=\frac{1}{m(1-q^m)},\quad m=1,2,\dots \qquad (6.12.24)$$

we have $s_\lambda(\mathbf{t}^*)=q^{-n(\lambda)}H_\lambda(q)$, see (6.11.10). We get Milne's hypergeometric function (8.0.20):

$$\langle M|e^{H(\mathbf{t})}e^{-A(\mathbf{t}^*)}|M\rangle={}_p\Phi_s\left(\begin{matrix}a_1+M,\dots,a_p+M\\ b_1+M,\dots,b_s+M\end{matrix}\middle|q,\mathbf{x}^N\right)=$$

$$\sum_{\substack{\lambda\\ l(\lambda)\le N}}\frac{(q^{a_1+M};q)_\lambda\cdots(q^{a_p+M};q)_\lambda}{(q^{b_1+M};q)_\lambda\cdots(q^{b_s+M};q)_\lambda}\frac{q^{n(\lambda)}}{H_\lambda(q)}s_\lambda(\mathbf{x}^N). \qquad (6.12.25)$$

According to (6.5.8),(6.12.24) we have the following representation

$$ {}_p\Phi_s\left(\begin{matrix}a_1+M,\dots,a_p+M\\ b_1+M,\dots,b_s+M\end{matrix}\middle|q,\mathbf{x}^N\right)=\frac{1}{\Delta(x)}\prod_{i=1}^{N}\frac{1}{(x_ir(D_{x_i});q)_\infty}\cdot\Delta(x) \qquad (6.12.26)$$

$$=\frac{1}{\Delta}\exp_q(x_1r(D_1))\cdots\exp_q(x_Nr(D_N))\cdot\Delta$$

(the notations $(b;q)_\infty$, $\exp_q(\alpha)$ see in (8.0.14), (6.9.9), Δ is the same as in (6.12.10)). (See also (6.11.15) for different representation).

For this hypergeometric functions we have the linear equation (6.5.18) (where we choose r due to (6.12.23)), which is q-difference equation. For instance for the case $N=1,M=0$ we get (compare it with (6.12.13))

$$\left(\frac{1}{x}\left(1-q^D\right)-{}_pr_s^{(q)}(D)\right){}_p\Phi_s(a_1,\dots,a_p;b_1,\dots,b_s;q,x)=0,\quad D:=x\partial_x, \qquad (6.12.27)$$

where ${}_pr_s^{(q)}(D)$ is defined by (6.12.23). Milne-s function itself for $N=1$ is the ordinary basic hypergeometric function:

$$ {}_p\Phi_s\left(\begin{matrix}a_1+M,\dots,a_p+M\\ b_1+M,\dots,b_s+M\end{matrix}\middle|q,x\right)=\sum_{n=0}^{+\infty}\frac{(q^{a_1+M};q)_n\cdots(q^{a_p+M};q)_n}{(q^{b_1+M};q)_n\cdots(q^{b_s+M};q)_n}\frac{x^n}{(q;q)_n},\quad x=x_1(6.12.28)$$

Let us take $M=0$. We have the following integral representation

$$ {}_{p+1}\Phi_s\left(\begin{matrix}a,a_1,\dots,a_p\\ b_1,\dots,b_s\end{matrix}\middle|q,x_1,\dots,x_N\right)=\frac{(-1)^Nq^{\frac{N-N^2}{2}-Na}}{(1-q)^{\frac{3N-N^2}{2}}}\frac{1}{\prod_{i<j}^{N}(x_i-x_j)}\int_0^\infty\cdots\int_0^\infty \qquad (6.12.29)$$

$$\prod_{i<j}^{N}(\alpha_ix_i-\alpha_jx_j){}_p\Phi_s\left(\begin{matrix}a_1,\dots,a_p\\ b_1,\dots,b_s\end{matrix}\middle|q,\frac{\alpha_1(1-q)x_1}{q},\dots\frac{\alpha_N(1-q)x_N}{q}\right)\prod_{i=1}^{N}\frac{\alpha_i^{a-1}}{\exp_q(\alpha_i)}d_q\alpha_i$$

Also we have the Beta-type representation:

$$\begin{aligned}
{}_{p+1}\Phi_{s+1}\left(\begin{matrix}a,a_1,\dots,a_p\\ b,b_1,\dots,b_s\end{matrix}\,\middle|\, q,x_1,\dots,x_N\right) &= \left(\frac{1}{\Gamma_q(b-a)}\right)^N \frac{1}{\prod_{i<j}^N (x_i-x_j)}\int_0^1\cdots\int_0^1 \\
\prod_{i<j}^N(\alpha_i x_i-\alpha_j x_j)\,{}_p\Phi_s\left(\begin{matrix}a_1,\dots,a_p\\ b_1,\dots,b_s\end{matrix}\,\middle|\, q,\alpha_1x_1,\dots\alpha_N x_N\right) &\prod_{i=1}^N \alpha_i{}^{a-1}\frac{(\alpha_i q;q)_\infty}{(\alpha_i q^{b-a};q)_\infty}d_q\alpha_i
\end{aligned} \tag{6.12.30}$$

Example 4. To obtain Milne's hypergeometric function of two sets of variables $\mathbf{x}^N,\mathbf{y}^N$, we take $\mathbf{t}=\mathbf{t}(\mathbf{x}^N)$ and $\mathbf{t}^*=\mathbf{t}^*(\mathbf{y}^N)$. This choice restricts the sum over partitions λ with $l(\lambda)\le N$. We put

$${}_pr_s^{(q)}(n)=\frac{\prod_{i=1}^p(1-q^{a_i+n})}{\prod_{i=1}^s(1-q^{b_i+n})}\frac{1}{1-q^{N-M+n}}, \tag{6.12.31}$$

$$e^{-T_n}=\frac{1}{(1-q)^n\Gamma_q(n+N-M+1)}\frac{\prod_{i=1}^p(1-q)^n\Gamma_q(a_i+n+1)}{\prod_{i=1}^s(1-q)^n\Gamma_q(b_i+n+1)}, \tag{6.12.32}$$

$$\Gamma_q(a)=(1-q)^{1-a}\frac{(q;q)_\infty}{(q^a,q)_\infty},\qquad (q^a,q)_n=(1-q)^n\frac{\Gamma_q(a+n)}{\Gamma_q(a)}. \tag{6.12.33}$$

Here $\Gamma_q(a)$ is a q-deformed Gamma-function

$$\Gamma_q(a)=(1-q)^{1-a}\frac{(q;q)_\infty}{(q^a,q)_\infty},\qquad (q^a,q)_n=(1-q)^n\frac{\Gamma_q(a+n)}{\Gamma_q(a)}. \tag{6.12.34}$$

Using (6.11.7) and (6.11.10) (see III of [4]) we get

$$(q^{N-M};q)_\lambda=q^{-n(\lambda)}H_\lambda(q)s_\lambda(1,q,\dots,q^{N-1}) \tag{6.12.35}$$

(see Appendix C for the notations) and we obtain the Milne's formula (8.0.27)

$$\begin{aligned}
\tau_r(M,\mathbf{t}(\mathbf{x}^N),\mathbf{t}^*(\mathbf{y}^N)) &= {}_p\Phi_s\left(\begin{matrix}a_1+M,\dots,a_p+M\\ b_1+M,\dots,b_s+M\end{matrix}\,\middle|\, q,\mathbf{x}^N,\mathbf{y}^N\right)= \\
&\sum_{\substack{\lambda\\ l(\lambda)\le N}}\frac{q^{n(\lambda)}}{H_\lambda(q)}\frac{s_\lambda(\mathbf{x}^N)s_\lambda(\mathbf{y}^N)}{s_\lambda(1,q,\dots,q^{N-1})}\frac{(q^{a_1+M};q)_\lambda\cdots(q^{a_p+M};q)_\lambda}{(q^{b_1+M};q)_\lambda\cdots(q^{b_s+M};q)_\lambda}.
\end{aligned} \tag{6.12.36}$$

This is the KP tau-function (but not the TL one because (6.12.31) depends on TL variable M).

We have the same type of representation (6.12.20) and the same form of linear equation (6.12.22), however we shall use (6.12.31) for r in formulae (6.12.20),(6.12.22).

To receive the basic hypergeometric function of one set of variables we must put indeterminates $\mathbf{y}^N$ in (6.12.25) as $y_i=q^{i-1}, i=(1,\dots,N)$. Thus we have

$$\begin{aligned}
&{}_p\Phi_s\left(\begin{matrix}a_1+M,\dots,a_p+M\\ b_1+M,\dots,b_s+M\end{matrix}\,\middle|\, q,\mathbf{x}^N\right)= \\
&\sum_{\substack{\lambda\\ l(\lambda)\le N}}\frac{(q^{a_1+M};q)_\lambda\cdots(q^{a_p+M};q)_\lambda}{(q^{b_1+M};q)_\lambda\cdots(q^{b_s+M};q)_\lambda}\frac{q^{n(\lambda)}}{H_\lambda(q)}s_\lambda(\mathbf{x}^N)
\end{aligned} \tag{6.12.37}$$

And for $N=1$ we have the ordinary q-deformed hypergeometric function:

$${}_p\Phi_s\left(\begin{matrix}a_1+M,\dots,a_p+M\\ b_1+M,\dots,b_s+M\end{matrix}\,\middle|\, q,x,y\right)=\sum_{n=0}^{+\infty}\frac{(q^{a_1+M};q)_n\cdots(q^{a_p+M};q)_n}{(q^{b_1+M};q)_n\cdots(q^{b_s+M};q)_n}\frac{(xy)^n}{(q;q)_n},\ \ xy=x_1y_1 \tag{6.12.38}$$

6.13 Baker-Akhiezer functions and the elements of Sato Grassmannian related to the tau function of hypergeometric type

This subsection is based on [12]. Let us write down the expression for Baker-Akhiezer functions (6.1.5) in terms of Hirota-Miwa variables (1.0.8):

$$w_\infty(M, -\mathbf{t}(\mathbf{x}^N), \mathbf{t}^*, \frac{1}{z}) = \frac{\tau_r(M, -\mathbf{t}(\mathbf{x}^{N+1}), \mathbf{t}^*)}{\tau_r(M, -\mathbf{t}(\mathbf{x}^N), \mathbf{t}^*)} \prod_{i=1}^{N} (1 - \frac{x_i}{z}), \quad \mathbf{x}^{N+1} = (x_1, \ldots, x_N, z), \quad (6.13.1)$$

$$w_\infty^*(M, \mathbf{t}(\mathbf{x}^N), \mathbf{t}^*, \frac{1}{z}) = -\frac{\tau_r(M, \mathbf{t}(\mathbf{x^N}) + [\mathbf{z}], \mathbf{t}^*)}{\tau_r(M, \mathbf{x}^N, \mathbf{t}^*)} \prod_{i=1}^{N} \frac{1}{(1 - \frac{x_i}{z})} \frac{dz}{z}, \quad [\mathbf{z}] = (z, \frac{z^2}{2}, \ldots). \quad (6.13.2)$$

We see that the variables $x_k, k = 1, \ldots, N$ are zeroes of $w_\infty(z)$ and poles of $w_\infty^*(z)$.

Remark 1 *The associated linear problems for Baker-Akhiezer functions are read as*

$$(\partial_{t_1} - \partial_{t_1}\phi_n)\, w(n, \mathbf{t}, \mathbf{t}^*, z) = w(n+1, \mathbf{t}, \mathbf{t}^*, z), \quad (6.13.3)$$

$$\partial_{t_1^*} w(n, \mathbf{t}, \mathbf{t}^*, z) = r(n) e^{\phi_{n-1} - \phi_n} w(n-1, \mathbf{t}, \mathbf{t}^*, z). \quad (6.13.4)$$

where w is either w_∞ or w_0. The compatibility of these equations gives rise to the equation (6.3.11). Taking into account the second eq.(6.3.11), equations (6.13.3), (6.13.4) may be also viewed as the recurrent equations for the tau-functions which depend on different number of variables $\mathbf{x}^N$.

Let us write down a plane of Baker-Akhiezer functions (6.1.5), which characterizes elements of Sato Grassmannian related to the tau-function (6.3.1) $\tau_r(M, \mathbf{t}, \mathbf{t}^*)$. We take $x_k = 0, k = 1, \ldots, N$ in (6.13.1) and obtain:

$$w_\infty(n, \mathbf{0}, \mathbf{t}^*, z) = z^n (1 + \sum_{m=1}^{\infty} r(n) r(n-1) \cdots r(n-m+1) h_m(-\mathbf{t}^*) z^{-m}), \quad n = M, M+1, M+2, ... \quad (6.13.5)$$

The dual plane is

$$w_\infty^*(n, \mathbf{0}, \mathbf{t}^*, z) = z^{-n} (1 + \sum_{m=1}^{\infty} r(n) r(n+1) \cdots r(n+m-1) h_m(\mathbf{t}^*) z^{-m}) dz, \quad n = M, M+1, M+2, ... \quad (6.13.6)$$

About these formulae see also (6.14.22),(6.14.23).
We see that when r has zeroes, then the element of the Grassmannian defining the solution corresponds to the finite-dimensional Grassmannian.

If $r(0) = 0$ one can construct a plane

$$\{w_\infty^*(n, \mathbf{0}, \mathbf{t}^*, 1/z), n = 1, \ldots, M\} \quad (6.13.7)$$

which is a spanned by a finite number of vectors. The space spanned by these vectors and in addition by the vectors (6.13.5) gives the half infinite plane generated by $\{z^n, n \geq 0\}$.

6.14 Gauss factorization problem, additional symmetries, string equations and ΨDO on the circle [12]

Let us describe relevant string equations following Takasaki and Takebe [31],[33]. We shall also consider this topic in a more detailed paper.

Let us introduce infinite matrices to describe KP and TL flows and symmetries, see [27]. Zakharov-Shabat dressing matrices are K and $\bar{K}$. K is a lower triangular matrix with unit main diagonal: $(K)_{ii} = 1$. $\bar{K}$ is an upper triangular matrix. The matrices $K, \bar{K}$ depend on parameters $M, \mathbf{t}, \mathbf{T}, \mathbf{t}^*$. The matrices $(\Lambda)_{ik} = \delta_{i,k-1}$, $(\bar{\Lambda})_{ik} = \delta_{i,k+1}$. For each value of $\mathbf{t}, T, \mathbf{t}^*$ and $M \in Z$ they solve Gauss (Riemann-Hilbert) factorization problem for infinite matrices:

$$\bar{K} = KG(M, \mathbf{t}, T, \mathbf{t}^*), \quad G(M, \mathbf{t}, T, \mathbf{t}^*) = \exp\left(\xi(\mathbf{t}, \Lambda)\right) \Lambda^M G(\mathbf{0}, T, \mathbf{0}) \bar{\Lambda}^M \exp\left(\xi(\mathbf{t}^*, \bar{\Lambda})\right). \tag{6.14.1}$$

We put $\log(\bar{K}_{ii}) = \phi_{i+M}$, and a set of fields $\phi_i(\mathbf{t}, \mathbf{t}^*), (-\infty < i < +\infty)$ solves the hierarchy of higher two-dimensional TL equations.

Take $L = K\Lambda K^{-1}$, $\bar{L} = \bar{K}\bar{\Lambda}\bar{K}^{-1}$, and $(\Delta)_{ik} = i\delta_{i,k}$, $\widehat{M} = K\Delta K^{-1} + M + \sum n t_n L^n$, $\widehat{\bar{M}} = \bar{K}\Delta\bar{K}^{-1} + M + \sum n t_n^* \bar{L}^n$. Then the KP additional symmetries [31],[33],[34],[13], [35] and higher TL flows [27] are written as

$$\partial_{\mathbf{t}_n^*} K = -\left(\left(r(\widehat{M})L^{-1}\right)^n\right)_- K, \quad \partial_{\mathbf{t}_n^*} \bar{K} = \left(\left(r(\widehat{M})L^{-1}\right)^n\right)_+ \bar{K}, \tag{6.14.2}$$

$$\partial_{t_n^*} K = -\left(\bar{L}^n\right)_- K, \quad \partial_{t_n^*} \bar{K} = \left(\bar{L}^n\right)_+ \bar{K}. \tag{6.14.3}$$

Then the string equations are

$$\bar{L}L = r(\widehat{M}), \tag{6.14.4}$$

$$\widehat{\bar{M}} = \widehat{M}. \tag{6.14.5}$$

The first equation is a manifestation of the fact that the group time $\mathbf{t}_1^*$ of the additional symmetry of KP can be identified with the Toda lattice time t_1^*. In terms of tau-function we have the equation (6.6.7) in terms of vertex operator action [34],[13], or the equations (6.5.19) in case the tau-function is written in Hirota-Miwa variables.

The second string equation (6.14.5) is related to the symmetry of our tau-functions with respect to $\mathbf{t} \leftrightarrow \mathbf{t}^*$.

When

$$r(M) = M + a, \tag{6.14.6}$$

the equations (6.14.4),(6.14.5) describe $c = 1$ string , see [32],[33]. In this case we easily get the relation

$$[\bar{L}, L] = 1. \tag{6.14.7}$$

The string equations in the form of Takasaki allows us to notice the similarity to the different problem. The dispersionless limit of (6.14.7) (and also of (6.14.5), (6.14.4), where $r(M) = M^n$, and of (6.14.6)) will be written as

$$\bar{\lambda}\lambda = \mu^n, \quad n \in Z, \tag{6.14.8}$$

$$\bar{\mu} = \mu. \tag{6.14.9}$$

The case when $\bar{\lambda}$ and $\bar{\mu}$ are complex conjugate of λ and of μ respectively, is of interest. These string equations (mainly the case $n = 1$) were recently investigated to solve the so-called Laplacian growth problem, see [43]. We are grateful to A.Zabrodin for the discussion on this problem. For the dispersionless limit of the KP and TL hierarchies see [31].

In case the function $r(n)$ has zeroes (described by divisor $\mathbf{m} = (M_1, ...)$: $r(M_k) = 0$), one needs to produce the replacement:

$$\Lambda \to \Lambda(\mathbf{m}), \quad \bar{\Lambda} \to \bar{\Lambda}(\mathbf{m}), \tag{6.14.10}$$

where new matrices $\Lambda(\mathbf{m})$,$\bar{\Lambda}(\mathbf{m})$ are defined as

$$(\Lambda(\mathbf{m}))_{i,j} = \delta_{i,j-1}, j \neq M_k, \quad (\Lambda(\mathbf{m}))_{i,j} = 0, j = M_k, \tag{6.14.11}$$

$$(\bar{\Lambda}(\mathbf{m}))_{i,j} = \delta_{i,j+1}, i \neq M_k, \quad (\bar{\Lambda}(\mathbf{m}))_{i,j} = 0, i = M_k. \tag{6.14.12}$$

This modification describes the open TL equation (**??**):

$$\partial_{t_1}\partial_{\mathbf{t}_1^*}\varphi_n = \delta(n)e^{\varphi_{n-1}-\varphi_n} - \delta(n+1)e^{\varphi_n-\varphi_{n+1}}. \tag{6.14.13}$$

The set of fields $\phi_\infty, \dots, \phi_{M_1}, \phi_{M_1+1}\dots$ consists of the following parts due to the conditions

$$\sum_{n=-\infty}^{M_s} \phi_n = 0, \qquad \sum_{n=M_k+1}^{M_{k+1}} \phi_n = 0, \qquad \sum_{n=\infty}^{M_1+1} \phi_n = 0, \tag{6.14.14}$$

which result from $\tau(M_k, \mathbf{t}, \mathbf{t}^*) = 1$.

Remark 2 *The matrix* $r(\widehat{M})$ *contains* $(\widehat{M} - b_i)$ *in the denominator. The matrix* $(\widehat{M} - b_i)^{-1}$ *is* $K(\Delta - b_i)^{-1}K^{-1}(1 + O(\mathbf{t}))$ *(compare the consideration of the inverse operators with [35]).*

The KP tau-function (3.4.8) can be obtained as follows.

$$G(M, \mathbf{t}, T, \mathbf{t}^*) = G(\mathbf{0}, T, \mathbf{0})U(M, \mathbf{t}, \mathbf{t}^*), \quad U(M, \mathbf{t}, \mathbf{t}^*) = U^+(\mathbf{t})U^-(M, \mathbf{t}^*). \tag{6.14.15}$$

$$U^+(\mathbf{t}) = \exp\left(\xi(\mathbf{t}, \Lambda)\right), \quad U^-(M, \mathbf{t}^*) = \exp\left(\xi(\mathbf{t}^*, \bar{\Lambda}r\left(\Delta + M\right))\right), \tag{6.14.16}$$

The matrix $G(\mathbf{0}, T, \mathbf{0})$ is related to the transformation of the eq.(6.3.14) to the eq.(6.3.11). By taking the projection [27] $U \mapsto U_{--}$ for non positive values of matrix indices we obtain a determinant representation of the tau-function (3.4.8):

$$\tau_r(M, \mathbf{t}, \mathbf{t}^*) = \frac{\det U_{--}(M, \mathbf{t}, \mathbf{t}^*)}{\det\left(U^+_{--}(\mathbf{t})\right)\det\left(U^-_{--}(M, \mathbf{t}^*)\right)} = \det U_{--}(M, \mathbf{t}, \mathbf{t}^*), \tag{6.14.17}$$

since both determinants in the denominator are equal to one. Formula (6.14.17) is also a Segal-Wilson formula for $GL(\infty)$ 2-cocycle [46] $C_M\left(U^+(-\mathbf{t}), U^-(-\mathbf{t}^*)\right)$. Choosing the function r as in *Section 3.2* we obtain hypergeometric functions listed in the *Introduction.*

Remark 3 *Therefore the hypergeometric functions which were considered above have the meaning of* $GL(\infty)$ *two-cocycle on the two multi-parametric group elements* $U^+(\mathbf{t})$ *and* $U^-(M, \mathbf{t}^*)$. *Both elements* $U^+(\mathbf{t})$ *and* $U^-(M, \mathbf{t}^*)$ *can be considered as elements of group of pseudo-differential operators on the circle. The corresponding Lie algebras consist of the multiplication operators* $\{z^n; n \in N_0\}$ *and of the pseudo-differential operators* $\{\left(\frac{1}{z}r(z\frac{d}{dz} + M)\right)^n; n \in N_0\}$. *Two sets of group times* $\mathbf{t}$ *and* $\mathbf{t}^*$ *play the role of indeterminates of the hypergeometric functions (6.12.2). Formulas (3.4.8) and (6.3.8) mean the expansion of* $GL(\infty)$ *group 2-cocycle in terms of corresponding Lie algebra 2-cocycle*

$$c_M(z, \frac{1}{z}r(D)) = r(M), \quad c_M(\tilde{r}(D)z, \frac{1}{z}r(D + M)) = \tilde{r}(M)r(M). \tag{6.14.18}$$

Japanese cocycle is cohomological to Khesin-Kravchenko cocycle [51] for the ΨDO *on the circle:*

$$c_M \sim c_0 \sim \omega_M, \tag{6.14.19}$$

which is

$$\omega_M(A,B) = \frac{1}{2\pi\sqrt{-1}} \oint \boldsymbol{res}_\partial A[\log(D+M), B]dz, \quad A, B \in \Psi DO. \tag{6.14.20}$$

For the group cocycle we have

$$C_M\left(e^{-\sum z^n t_n}, e^{-\sum (z^{-1}r(D))^n \mathbf{t}^*_n}\right) = \tau_r(M, \mathbf{t}, \mathbf{t}^*), \tag{6.14.21}$$

where we imply that the order of ΨDO $r(D)$ *is 1 or less.*

Remark 4 *It is interesting to note that in case of hypergeometric functions* ${}_pF_s$ *(8.0.17) the order of* r *is* $p-s$ *(see* Example 3*), and the condition* $p-s \leq 1$ *is the condition of the convergence of this hypergeometric series, see [25]. Namely the radius of convergence is finite in case* $p-s=1$*, it is infinite when* $p-s<1$ *and it is zero for* $p-s>1$ *(this is true for the case when no one of* a_k *in (8.0.17) is nonnegative integer).*

Remark 5 *The set of functions* $\{w(n,z), n = M, M+1, M+2, \ldots\}$*, where*

$$w(n,z) = \exp\left(-\sum_{m=0}^{\infty} t^*_m \left(\frac{1}{z}r(D)\right)^m\right) \cdot z^n, \tag{6.14.22}$$

may be identified with Sato Grassmannian (6.13.5) related to the cocycle (6.14.21). The dual Grassmannian (6.13.6) is the set of one forms $\{w^*(n,z), n = M, M+1, M+2, \ldots\}$*,*

$$w^*(n,z) = \exp\left(\sum_{m=0}^{\infty} t^*_m \left(\frac{1}{z}r(-D)\right)^m\right) \cdot z^{-n}dz. \tag{6.14.23}$$

7 Appendices B

7.1 A scalar product in the theory of symmetric functions [4]

We consider a ring of symmetric functions of variables $x_i, i = 1, 2, \ldots, N$, where N may be infinity. Let us remember the notion of scalar product. Different choices of scalar product give rise to different systems of bi-orthogonal symmetric polynomial functions (like Schur functions, Macdonald functions and so on).

Actually we consider two sets of variables, the first one is $\mathbf{x} = (x_1, x_2, \ldots, x_N)$, the second is $\mathbf{p} = (p_1, p_2, \ldots)$. These two sets are related via

$$p_m = \sum_{i=1}^{N} x_i^m, \quad m = 1, 2, \ldots \tag{7.1.1}$$

If one takes

$$e^{\sum_{m=1}^\infty \frac{1}{m} v_m^{-1} p_m p^*_m} = \sum_\lambda P_\lambda(\mathbf{p}) Q_\lambda(\mathbf{p}^*) = \sum_\lambda \frac{p_\lambda p^*_\lambda}{z^{(v)}_\lambda}, \tag{7.1.2}$$

then for any partitions μ, λ

$$\langle P_\mu, Q_\lambda \rangle_v = \delta_{\mu,\lambda} \tag{7.1.3}$$

with respect to the scalar product

$$\langle p_\mu, p_\lambda \rangle_v = z_\lambda^{(v)} \delta_{\mu,\lambda} \tag{7.1.4}$$

When $v = 1$ then

$$z_\lambda^{(v)} = z_\lambda, \quad z_\lambda = \prod_i i^{m_i} m_i! \tag{7.1.5}$$

where $m_i = m_i(\lambda)$ is the number of parts of λ equal to i

$$p_\mu = \prod_i p_{\mu_i}, \quad p_{\mu_i} = \sum_k x_k^{\mu_i}, \quad m t_m = p_m \tag{7.1.6}$$

If one takes

$$v_m = \frac{1 - t^m}{1 - q^m} \tag{7.1.7}$$

he gets Macdonald's polynomials $Q_\lambda(\mathbf{p}) = Q_\lambda(\mathbf{p}; q, t), P_\lambda(\mathbf{p}) = P_\lambda(\mathbf{p}; q, t)$. In this case

$$z_\lambda^{(v)} = z_\lambda(q,t) = z_\lambda \prod_{i=1}^{l(\lambda)} \frac{1 - q^{\lambda_i}}{1 - t^{\lambda_i}}, \quad z_\lambda = \prod_i i^{m_i} m_i! \tag{7.1.8}$$

where $l(\lambda)$ is the length of λ and $m_i = m_i(\lambda)$ is the number of parts of λ equal to i.

7.2 A deformation of the scalar product and series of hypergeometric type

For different choice of function v we choose numbers $a^{(v)}, b^{(v)}$ and a function $r^{(v)}$ which is a function of one variable M, say. We define

$$r_\lambda^{(v)}(M) = \prod_{i,j \in \lambda} r^{(v)}(M + a^{(v)} i + b^{(v)} j) \tag{7.2.1}$$

as a product over all boxes of the Young diagram. In what follows we shall omit the superscript (v).

Now let us deform the scalar product (7.1.3) in a following way

$$\langle P_\mu, Q_\lambda \rangle_{v,r,M} = r_\lambda(M) \delta_{\mu,\lambda} \tag{7.2.2}$$

To get series of hypergeometric type we consider

$$\langle e^{\sum_{m=1}^\infty \sum_{k=1}^N v_m^{-1} x_k^m \frac{p_m}{m}}, e^{\sum_{m=1}^\infty \sum_{k=1}^N v_m^{-1} y_k^m \frac{p_m}{m}} \rangle_{v,r,M} \tag{7.2.3}$$

Using (7.1.2)

$$\langle e^{\sum_{m=1}^\infty \sum_{k=1}^N v_m^{-1} x_k^m \frac{p_m}{m}}, e^{\sum_{m=1}^\infty \sum_{k=1}^N v_m^{-1} y_k^m \frac{p_m}{m}} \rangle_{v,r,M} = \sum_\lambda r_\lambda(M) P_\lambda(\mathbf{x}^N) Q_\lambda(\mathbf{y}^N) \tag{7.2.4}$$

In this paper we shall consider two cases of (7.1.7). First one is

$$v_n = 1 \tag{7.2.5}$$

for all n. The second one is

$$v_{2n} = 0, \quad v_{2n+1} = 2 \tag{7.2.6}$$

The first case corresponds to Schur function series. The second one corresponds to projective Schur function series. These are the cases when the scalar product (7.2.2) can be presented as a vacuum expectation of certain fermionic fields, see *Section 4* and [52].

7.3 Scalar product of tau functions. A conjecture.

Before we considered tau functions of hypergeometric type (3.4.8). In this subsection we shall consider more general TL tau functions. Let $\tau_1(\mathbf{t},\mathbf{t}^*),\tau_2(\mathbf{t},\mathbf{t}^*)$ be TL tau functions of the form

$$\tau_1(\mathbf{t},\mathbf{t}^*)=e^{X(\mathbf{t})}\cdot e^{\sum_{m=1}^\infty mt_mt_m^*},\quad \tau_2(\mathbf{t},\mathbf{t}^*)=e^{Y(\mathbf{t}^*)}\cdot e^{\sum_{m=1}^\infty mt_mt_m^*} \tag{7.3.1}$$

where $e^X=1+X+\cdots,e^Y=1+Y+\cdots$, and where the operators $X(\mathbf{t})$ and $Y(\mathbf{t}^*)$ are linear combinations of vertex operators (6.1.1) and (6.1.2) respectively:

$$X_i(\mathbf{t})=\int f(z,z')V_\infty(\mathbf{t},z)V_\infty^*(\mathbf{t},z')dzdz' \tag{7.3.2}$$

$$Y_i(\mathbf{t}^*)=\int g(z,z')V_0(\mathbf{t}^*,z)V_\infty^*(\mathbf{t}^*,z')dzdz' \tag{7.3.3}$$

If the TL tau functions (7.3.1) have the typical forms (1.0.9) (see [28],[29],[30])

$$\tau_1(\mathbf{t},\mathbf{t}^*)=\sum_{\lambda,\mu}K_{\lambda\mu}s_\lambda(\mathbf{t})s_\mu(\mathbf{t}^*),\quad \tau_2(\mathbf{t},\mathbf{t}^*)=\sum_{\lambda,\mu}M_{\lambda\mu}s_\lambda(\mathbf{t})s_\mu(\mathbf{t}^*) \tag{7.3.4}$$

then their (standard) scalar product (2.1.15) is obviously equal to

$$<\tau_1(\mathbf{t},\gamma),\tau_2(\gamma,\mathbf{t}^*)>=\sum_{\lambda,\mu}N_{\lambda\mu}s_\lambda(\mathbf{t})s_\mu(\mathbf{t}^*),\quad N_{\lambda\mu}=\sum_\nu K_{\lambda\nu}M_{\nu\mu} \tag{7.3.5}$$

It is not immediately clear that what we got is a tau function again (without solving Hirota equations). However on the other hand we have

$$<\tau_1(\mathbf{t},\gamma),\tau_2(\gamma,\mathbf{t}^*)>=e^{X(\mathbf{t})}\cdot e^{Y(\mathbf{t}^*)}\cdot<e^{\sum_{m=1}^\infty mt_m\gamma_m},e^{\sum_{m=1}^\infty m\gamma_mt_m^*}>=e^{X(\mathbf{t})}\cdot e^{Y(\mathbf{t}^*)}\cdot e^{\sum_{m=1}^\infty mt_mt_m^*} \tag{7.3.6}$$

The last relation shows that we get tau function again according to general results of [7]. (It was shown there that the action of vertex operators e^X to KP tau function gives rise to a new KP tau function. TL tau function is the tau function of the pair of KP hierarchies related to the sets of times $\mathbf{t}$ and $\mathbf{t}^*$, see [27]).

Due to (5.7.1) it means that if the following integral over complex planes $\gamma_m, m=1,2,\dots$

$$\int\tau_1(\mathbf{t},\gamma)e^{-\sum_{m=1}^\infty m|\gamma_m|^2}\tau_2(-\bar\gamma,\mathbf{t}^*)\prod_{m=1}^\infty\frac{md^2\gamma_m}{\pi} \tag{7.3.7}$$

exists, then it is a new TL tau function. Here the variables $\bar\gamma=(\bar\gamma_1,\bar\gamma_1,\dots)$ are complex conjugated to $\gamma=(\gamma_1,\gamma_2,\dots)$.

The proof can be done also directly as follows

$$\int\tau_1(\mathbf{t},\gamma)e^{-\sum_{m=1}^\infty m|\gamma_m|^2}\tau_2(-\bar\gamma,\mathbf{t}^*)\prod_{m=1}^\infty\frac{md^2\gamma_m}{\pi}= \tag{7.3.8}$$

$$e^{X(\mathbf{t})}\cdot e^{Y(\mathbf{t}^*)}\cdot\int e^{\sum_{m=1}^\infty mt_m\gamma_m-m|\gamma_m|^2-m\bar\gamma_mt_m^*}\prod_{m=1}^\infty\frac{md^2\gamma_m}{\pi}= \tag{7.3.9}$$

$$e^{X(\mathbf{t})}\cdot e^{Y(\mathbf{t}^*)}\cdot e^{\sum_{m=1}^\infty mt_mt_m^*} \tag{7.3.10}$$

which is tau function.

Let us try to consider more general situation. Let $\tau_0(\mathbf{t},\mathbf{t}^*)$ be a TL tau function with the following properties. First, $\tau_0(\mathbf{t},\mathbf{t}^*)=\tau_0(\mathbf{t}^*,\mathbf{t})$. Second, each integral over complex plane $\gamma_m, m=1,2,\dots$

$$\int \tau_0(\gamma,-\bar{\gamma})d^2\gamma_m = a_m \tag{7.3.11}$$

where $\bar{\gamma}_m$ is complex conjugated of γ_m, is convergent and non-vanishing. Then

$$\int \tau_0(\gamma,-\bar{\gamma})\prod_{m=1}^{\infty}\frac{d^2\gamma_m}{a_m} = 1 \tag{7.3.12}$$

Given τ_0, we introduce the following scalar product

$$< f,g >_0 = \int f(\gamma)\tau_0(\gamma,-\bar{\gamma})g(-\bar{\gamma})\prod_{m=1}^{\infty}\frac{d^2\gamma_m}{a_m} \tag{7.3.13}$$

Let us present a conjecture that scalar product of TL tau functions is TL tau function again.

Conjecture. Let $\tau_1(\mathbf{t},\mathbf{t}^*),\tau_2(\mathbf{t},\mathbf{t}^*)$ be TL tau functions. If the integral over complex planes $\gamma_m, m=1,2,\dots$

$$\int \tau_1(\mathbf{t},\gamma)\tau_0(\gamma,-\bar{\gamma})\tau_2(-\bar{\gamma},\mathbf{t}^*)\prod_{m=1}^{\infty}\frac{d^2\gamma_m}{a_m}, \tag{7.3.14}$$

exists, then it is a new TL tau function. □

7.4 Integral representation of scalar products

For the scalar product (7.2.2) one can present the following integral realization.

In case $r=1$

$$\langle f,g\rangle = \int\cdots\int f(\mathbf{t})g(\bar{\mathbf{t}})e^{-\sum_{m=1}^{\infty}m|t_m|^2}\prod_{m=1}^{\infty}\frac{d^2t_m}{a_m} \tag{7.4.1}$$

where mt_m are power sum functions, $m\bar{t}_m$ are complex conjugated variables, and integration is going over complex planes $(t_m,\bar{t}_m)$. Normalization factors a_m depend on m:

$$a_m = \frac{\pi}{m} \tag{7.4.2}$$

In case $r(n)=n$ there is a representation

$$\langle f,g\rangle_{r,n} = \int\cdots\int f(\mathbf{z})g(\bar{\mathbf{z}})e^{-\sum_{k=1}^{N}|z_k|^2}|\Delta(z)|^2\prod_{k=1}^{N}\frac{d^2z_k}{b_k} \tag{7.4.3}$$

where $\bar{\mathbf{z}}=(\bar{z}_1,\dots,\bar{z}_N)$ are complex conjugated to $\mathbf{z}=(z_1,\dots,z_N)$, normalizing constants are

$$b_k = \pi \tag{7.4.4}$$

Δ is Vandermond determinant

$$\Delta(z) = \prod_{i<k}^{N}(z_k-z_i) \tag{7.4.5}$$

and one puts $\Delta(z)=1$ for $N=1$. The integration is going over complex planes $(z_k,\bar{z}_k)$.

For the scalar product (7.1.4) we present the representation

$$\langle f,g\rangle_v = \int\cdots\int f(\mathbf{t})g(\bar{\mathbf{t}})e^{-\sum_{m=1}^{\infty}\frac{m}{v_m}t_m\bar{t}_m}\prod_{m=1}^{\infty}\frac{dt_m d\bar{t}_m}{a_m} \tag{7.4.6}$$

where mt_m are power sum functions, $m\bar{t}_m$ are complex conjugated variables, and integration is going over complex planes $(t_m, \bar{t}_m)$. Normalization factors a_m depend on m:

$$a_m = \frac{2\pi v_m}{m}\sqrt{-1} \tag{7.4.7}$$

In particular for Macdonald's polynomials $Q_\lambda(\mathbf{p};q,t), P_\lambda(\mathbf{p};q,t)$, see (7.1.7), we get

$$\int\cdots\int Q_\lambda(\mathbf{p};q,t)P_\lambda(\bar{\mathbf{p}};q,t)e^{-\sum_{m=1}^\infty \mathbf{p}_m\bar{\mathbf{p}}_m\frac{1-q^m}{m(1-t^m)}}\prod_{m=1}^\infty \frac{dp_m d\bar{p}_m}{a_m} = \delta_{\mu,\lambda} \tag{7.4.8}$$

$$a_m = 2\pi\frac{1-q^m}{m(1-t^m)}\sqrt{-1} \tag{7.4.9}$$

8 Appendix C. Hypergeometric functions

Ordinary hypergeometric functions First let us remember that generalized hypergeometric function of one variable x is defined as

$${}_pF_s\left(a_1,\ldots,a_p;b_1,\ldots,b_s;x\right) = \sum_{n=0}^\infty \frac{(a_1)_n\cdots(a_p)_n}{(b_1)_n\cdots(b_s)_n}\frac{x^n}{n!}. \tag{8.0.10}$$

Here $(a)_n$ is Pochhammer's symbol:

$$(a)_n = \frac{\Gamma(a+n)}{\Gamma(a)} = a(a+1)\cdots(a+n-1). \tag{8.0.11}$$

This hypergeometric function solves the equation

$$\left(\prod_{k=0}^s\left(x\frac{d}{dx}+b_k-1\right) - x\prod_{j=1}^p\left(x\frac{d}{dx}+a_j\right)\right){}_pF_s\left(a_1,\ldots,a_p;b_1,\ldots,b_s;x\right) = 0, \quad b_0 = 1 \tag{8.0.12}$$

Given number q, $|q|<1$, the so-called basic hypergeometric series of one variable is defined as

$${}_p\Phi_s\left(a_1,\ldots,a_p;b_1,\ldots,b_s;q,x\right) = \sum_{n=0}^\infty \frac{(q^{a_1};q)_n\cdots(q^{a_p};q)_n}{(q^{b_1};q)_n\cdots(q^{b_s};q)_n}\frac{x^n}{(q;q)_n}. \tag{8.0.13}$$

Here $(q^a,q)_n$ is q-deformed Pochhammer's symbol:

$$(b;q)_0 = 1, \quad (b;q)_n = (1-b)(1-bq^1)\cdots(1-bq^{n-1}). \tag{8.0.14}$$

Both series converge for all x in case $p < s+1$. In case $p = s+1$ they converge for $|x| < 1$. We refer these well-known hypergeometric functions as ordinary hypergeometric functions.

The multiple hypergeometric series related to Schur polynomials [26],[25],[38]. There are several well-known different multi-variable generalizations of hypergeometric series of one variable [26, 25]. If one replaces the sum over n to a sum over partitions $\lambda = (n_1, n_2, \ldots)$, and replaces the single variable x to a Hermitian matrix X, he get one of generalizations of hypergeometric series which is called *hypergeometric function of matrix argument* $\mathbf{X}$ *with indices* $\mathbf{a}$ *and* $\mathbf{b}$ [39],[26]:

$${}_pF_s\left(\begin{matrix}a_1,\ldots,a_p\\ b_1,\ldots,b_s\end{matrix}\middle|\,\mathbf{X}\right) = \sum_{\substack{\lambda\\ l(\lambda)\le N}} \frac{(a_1)_\lambda\cdots(a_p)_\lambda}{(b_1)_\lambda\cdots(b_s)_\lambda}\frac{Z_\lambda(\mathbf{X})}{|\lambda|!}. \tag{8.0.15}$$

Here the sum is over all different partitions $\lambda = (n_1, n_2, \ldots, n_k)$, where $n_1 \geq n_2 \geq \cdots \geq n_k$, $k \leq |\lambda|$, $|\lambda| = n_1 + \cdots + n_r$ and whose length $l(\lambda) = k \leq N$ ($n_k \neq 0$). $\mathbf{X}$ is a Hermitian $N \times N$ matrix, and $Z_\lambda(\mathbf{X})$ are zonal spherical polynomial for the symmetric spaces of the following types: $GL(N,R)/SO(N)$, $GL(N,C)/U(N)$, or $GL(N,H)/Sp(N)$ see [25]. The definition of symbol $(a)_\lambda$ depends on the choice of the symmetric space:

$$(a)_\lambda = (a)_{n_1}(a - \frac{1}{\alpha})_{n_2} \cdots (a - \frac{k-1)}{\alpha})_{n_k}, \quad (a)_{\mathbf{0}} = 1, \tag{8.0.16}$$

where $\alpha = 2$, $\alpha = 1$ and $\alpha = \frac{1}{2}$ for the symmetric spaces $GL(N,R)/SO(N)$, $GL(N,C)/U(N)$, and $GL(N,H)/Sp(N)$ respectively. The function (8.0.15) actually depends on the eigenvalues of matrix X which are $\mathbf{x}^N = (x_1, \ldots, x_N)$. In what follows we consider only the case of $GL(N,C)/U(N)$ symmetric space. Then zonal spherical polynomial $Z_\lambda(\mathbf{X})$ is proportional to the Schur function $s_\lambda(x_1, x_2, ..., x_N)$ corresponding to a partition λ [4], $s_\lambda(x_1, x_2, ..., x_N)$ is a symmetric function of variables x_k.

For this choice of the symmetric space the hypergeometric function can be rewritten as follows

$${}_pF_s\left(\begin{matrix} a_1, \ldots, a_p \\ b_1, \ldots, b_s \end{matrix} \middle| \mathbf{X}\right) = \sum_{\substack{\lambda \\ l(\lambda) \leq N}} \frac{(a_1)_\lambda \cdots (a_p)_\lambda}{(b_1)_\lambda \cdots (b_s)_\lambda} \frac{s_\lambda(\mathbf{x}^N)}{H_\lambda}. \tag{8.0.17}$$

where H_λ is 'hook product':

$$H_\lambda = \prod_{(i,j)\in\lambda} h_{ij}, \quad h_{ij} = (n_i + n'_j - i - j + 1) \tag{8.0.18}$$

and

$$(a)_\lambda = (a)_{n_1}(a-1)_{n_2} \cdots (a-k+1)_{n_k}, \quad (a)_{\mathbf{0}} = 1 \tag{8.0.19}$$

Taking $N = 1$ we get (8.0.10).

For this hypergeometric functions we suggested different representations like (6.11.12),(6.11.13) or like (6.12.14),(6.12.15),(6.12.16).

Let $|q| < 1$. The multiple *basic* hypergeometric series related to Schur polynomials were suggested by L.G.Macdonald and studied by S.Milne [38]. These series are as follows

$${}_p\Phi_s\left(a_1, \ldots, a_p; b_1, \ldots, b_s; q, \mathbf{x}^N\right) = \sum_{\substack{\lambda \\ l(\lambda) \leq N}} \frac{(q^{a_1}; q)_\lambda \cdots (q^{a_p}; q)_\lambda}{(q^{b_1}; q)_\lambda \cdots (q^{b_s}; q)_\lambda} \frac{q^{n(\lambda)}}{H_\lambda(q)} s_\lambda\left(\mathbf{x}^N\right), \tag{8.0.20}$$

where the sum is over all different partitions $\lambda = (n_1, n_2, \ldots, n_k)$, where $n_1 \geq n_2 \geq \cdots \geq n_k \geq 0$, $k \leq |\lambda|$, $|\lambda| = n_1 + \cdots + n_k$ and whose length $l(\lambda) = k \leq N$. Schur polynomial $s_\lambda\left(\mathbf{x}^N\right)$, with $N \geq l(\lambda)$, is a symmetric function of variables $\mathbf{x}^N$ and defined as follows [4]:

$$s_\lambda(\mathbf{x}^N) = \frac{a_{\lambda+\delta}}{a_\delta}, \quad a_\lambda = \det(x_i^{n_j})_{1\leq i,j\leq N}, \quad \delta = (N-1, N-2, \ldots, 1, 0). \tag{8.0.21}$$

Coefficient $(q^c; q)_\lambda$ associated with partition λ is expressed in terms of the q-deformed Pochhammer's Symbols $(q^c; q)_n$ (8.0.14):

$$(q^c; q)_\lambda = (q^c; q)_{n_1}(q^{c-1}; q)_{n_2} \cdots (q^{c-k+1}; q)_{n_k}. \tag{8.0.22}$$

The multiple $q^{n(\lambda)}$ defined on the partition λ:

$$q^{n(\lambda)} = q^{\sum_{i=1}^k (i-1)n_i}, \tag{8.0.23}$$

and q-deformed 'hook polynomial' $H_\lambda(q)$ is

$$H_\lambda(q) = \prod_{(i,j)\in\lambda} \left(1 - q^{h_{ij}}\right), \quad h_{ij} = (n_i + n'_j - i - j + 1), \tag{8.0.24}$$

where λ' is the conjugated partition (for the definition see [4]).

For this hypergeometric functions we suggested different representations like (6.11.14),(6.11.15) or like (6.12.29),(6.12.30).

For $N = 1$ we get (8.0.13).

Let us note that in the limit $q \to 1$ series (8.0.20) reduces to (8.0.17), see [25].

Hypergeometric series of double set of arguments [26],[25],[38]

Another generalization of hypergeometric series is so-called hypergeometric function of two matrix arguments $\mathbf{X}, \mathbf{Y}$ with indices $\mathbf{a}$ and $\mathbf{b}$:

$${}_p\mathcal{F}_s\left(a_1, \ldots, a_p; b_1, \ldots, b_s; \mathbf{X}, \mathbf{Y}\right) = \sum_\lambda \frac{(a_1)_\lambda \cdots (a_p)_\lambda}{(b_1)_\lambda \cdots (b_s)_\lambda} \frac{Z_\lambda(\mathbf{X}) Z_\lambda(\mathbf{Y})}{|\lambda|! Z_\lambda(\mathbf{I}_N)}. \tag{8.0.25}$$

Here $\mathbf{X}, \mathbf{Y}$ are Hermitian $N \times N$ matrices and $Z_\lambda(\mathbf{X}), Z_\lambda(\mathbf{Y})$ are zonal spherical polynomials for the symmetric spaces $GL(N,C)/U(N)$, $GL(N,R)/SO(N)$ and $GL(N,H)/Sp(N)$ see [25]. The notations are the same as in previous subsection. Again we shall consider only the case of $GL(N,C)/U(N)$ symmetric spaces. In this case (8.0.25) it may be written as

$${}_p\mathcal{F}_s\left(a_1, \ldots, a_p; b_1, \ldots, b_s; \mathbf{x}^{(N)}, \mathbf{y}^{(N)}\right) = \sum_\lambda \frac{(a_1)_\lambda \cdots (a_p)_\lambda}{(b_1)_\lambda \cdots (b_s)_\lambda} \frac{s_\lambda(\mathbf{x}^N) s_\lambda(\mathbf{y}^N)}{H_\lambda s_\lambda(\mathbf{1}^N)}, \tag{8.0.26}$$

where $\mathbf{1}^N = (1, \ldots, 1)$.

We suggest a different representations like (6.12.20) or different integral representations.

The q-deformation of the hypergeometric function (8.0.26) is as follows

$${}_p\Phi_s\left(\begin{matrix} a_1, \ldots, a_p \\ b_1, \ldots, b_s \end{matrix}\middle| q, \mathbf{x}^N, \mathbf{y}^N\right) =$$

$$\sum_{\substack{\lambda \\ l(\lambda)\le N}} \frac{(q^{a_1}; q)_\lambda \cdots (q^{a_p}; q)_\lambda}{(q^{b_1}; q)_\lambda \cdots (q^{b_s}; q)_\lambda} \frac{q^{n(\lambda)}}{H_\lambda(q)} \frac{s_\lambda\left(\mathbf{x}^N\right) s_\lambda\left(\mathbf{y}^N\right)}{s_\lambda\left(1, q, q^2, ..., q^{N-1}\right)} \tag{8.0.27}$$

This is the multi-variable basic hypergeometric function of two sets of variables which was also studied by S.Milne, see [38], [25].

There are also hypergeometric functions related to Jack polynomials $C_\lambda^{(d)}$ [25]:

$${}_p\mathcal{F}_s^{(d)}\left(a_1, \ldots, a_p; b_1, \ldots b_s; \mathbf{x}^N, \mathbf{y}^N\right) =$$

$$\sum_\lambda \frac{(a_1)_\lambda^{(d)} \cdots (a_p)_\lambda^{(d)}}{(b_1)_\lambda^{(d)} \cdots (b_s)_\lambda^{(d)}} \frac{C_\lambda^{(d)}(\mathbf{x}^N) C_\lambda^{(d)}(\mathbf{y}^N)}{|\lambda|! C_\lambda^{(d)}(1^n)}, \tag{8.0.28}$$

where

$$(a)_\lambda^{(d)} = \prod_{i=1}^{l(\lambda)} \left(a - \frac{d}{2}(i-1)\right)_{n_i}. \tag{8.0.29}$$

Here $(c)_k = c(c+1)\cdots(c+k-1)$. For the special value $d = 2$ the last expression (8.0.28) coincides with (8.0.20), and reduces to (8.0.26) as $|q| \to 1$. The open problem is to get a fermionic representation of (8.0.28) for arbitrary value of the parameter d.

9 Further generalization. Examples of Gelfand-Graev hypergeometric functions

9.1 Scalar product and series in plane partitions

Let us consider the scalar product (3.3.9)

$$\langle s_\mu, s_\lambda\rangle_{r,n} = r_\lambda(n)\delta_{\mu,\lambda} \tag{9.1.1}$$

For the simplicity of notations sometimes we shall write $\langle , \rangle_r$ instead of $\langle , \rangle_{r,n}$.

We introduce notations

$$(\mathbf{t},\mathbf{p}) = \sum_{m=1}^{\infty} p_m t_m, \quad (\mathbf{p},\beta_i) = \sum_{m=1}^{\infty} p_m \beta_{im}, \quad (\mathbf{p},\beta_i^*) = \sum_{m=1}^{\infty} p_m \beta_{im}^* \tag{9.1.2}$$

Lemma

$$\langle s_\lambda, e^{(\mathbf{p},\beta)} s_\mu\rangle_{r,n} = \langle s_\mu e^{(\mathbf{p},\beta)}, s_\lambda\rangle = r_\lambda(n) s_{\lambda/\mu}(\beta) \tag{9.1.3}$$

One sees that for $\beta = 0$ and $r = 1$ the right hand side is equal to $\delta_{\mu,\lambda}$. □

Proof One develops $e^{(p,\beta_i)}$ using

$$e^{(p,\beta_i)} = \sum_\nu s_\nu(\gamma) s_\nu(\beta_i) \tag{9.1.4}$$

and taking into account the orthogonality of Schur functions (9.1.1) and the formulas (see section 5 of chapter I of [4])

$$s_\nu s_\mu = \sum_\lambda c_{\mu\nu}^\lambda s_\lambda, \quad s_{\lambda/\mu} = \sum_\nu c_{\mu\nu}^\lambda s_\nu \tag{9.1.5}$$

□

Let us consider a set of scalar products (9.1.1) which differ by the choice of r. The scalar product of functions of p_i will be labelled by script i: we shall denote this scalar product as $\langle , \rangle_{r^{(i)}}$.

Lemma

$$e^{(\mathbf{p}_i,\beta_i+\gamma_{i-1})} = \sum_{\nu_1,\nu_2} s_{\nu_1}(\gamma_i) s_{\nu_1/\nu_2}(\beta_i) s_{\nu_2}(\gamma_{i-1}) \tag{9.1.6}$$

and for given partition ν_{i-1} we have

$$\langle e^{(\mathbf{p}_i,\beta_i+\gamma_{i-1})}, s_{\nu_{i-1}}(\gamma_{i-1})\rangle_{r^{(i-1)},n_{i-1}} = \sum_{\nu_i} s_{\nu_i}(\gamma_i) s_{\nu_i/\nu_{i-1}}(\beta_i) r_{\nu_{i-1}}^{(i-1)}(n_{i-1}) \tag{9.1.7}$$

where

$$\gamma_i = (\frac{p_{i1}}{1}, \frac{p_{i2}}{2}, \frac{p_{i3}}{3}, \dots) \tag{9.1.8}$$

□ **Proof** follows from the relations (see [4])

$$e^{\sum_{n=1}^{\infty}(p_{in}\beta_{in}+\frac{1}{n}p_{in}p_{i-1,n})} = \sum_{\nu_1} s_{\nu_1}(\beta_i+\gamma_{i-1}) s_{\nu_1}(\gamma_i), \quad s_{\nu_2}(\beta_i+\gamma_{i-1}) = \sum_{\nu_2} s_{\nu_1/\nu_2}(\beta_i) s_{\nu_2}(\gamma_{i-1}) \tag{9.1.9}$$

and definition (3.3.9). □

Let us consider a set of $\{\mathbf{p}_i, i = 0, 1, 2, \dots, k+1\}$ and a set of $\{\beta_i, i = 1, 2, \dots, k+1\}$ with conditions

$$\mathbf{p}_0 = 0, \quad \mathbf{p}_{k+1} = \mathbf{p} \tag{9.1.10}$$

With the help of Lemma we consider scalar product of scalar products (compare with (3.4.7)):

$$\langle e^{(\mathbf{p}_k,\beta_k+\gamma_{k-1})}\cdots,\langle e^{(\mathbf{p}_3,\beta_3+\gamma_2)},\langle e^{(\mathbf{p}_2,\beta_2+\gamma_1)},e^{(\mathbf{p}_1,\beta_1+\gamma_0)}\rangle_{r^{(1)},n_1}\rangle_{r^{(2)},n_2}\cdots\rangle_{r^{(k-1)},n_k} \tag{9.1.11}$$

$$= \sum_{\nu_k\supseteq\cdots\supseteq\nu_0} r^{(k-1)}_{\nu_{k-1}}(n_{k-1})r^{(k-2)}_{\nu_{k-2}}(n_{k-2})\cdots r^{(2)}_{\nu_2}(n_2)r^{(1)}_{\nu_1}(n_1)$$

$$s_{\nu_k}(\gamma_k)s_{\nu_k/\nu_{k-1}}(\beta_k)s_{\nu_{k-1}/\nu_{k-2}}(\beta_{k-1})\cdots s_{\nu_2/\nu_1}(\beta_2)s_{\nu_1/\nu_0}(\beta_1)s_{\nu_0}(\gamma_0) \tag{9.1.12}$$

$$= I_k(\mathbf{p},\mathbf{r}^{k-1},\lambda^{k-1},\beta,\mathbf{p}_0) \tag{9.1.13}$$

where $\mathbf{r}^{k-1}=(r^{(1)},r^{(2)},\ldots,r^{(k-1)})$, $\lambda_{k-1}=(n_1,\ldots,n_{k-1})$, $s_{\nu_0}(\gamma_0=0)=\delta_{0,\nu_0}$.
This is the sum over all *plane partitions* [4] with the largest part k: $\pi^{(k)}=(\nu_1\subseteq\cdots\subseteq\nu_k)$.

Now let us consider scalar product of different I and with respect to standard scalar product (9.1.1) . Putting $\mathbf{p}_0=0$ we get a function of $\beta=\{\beta_{in},i=1,\ldots,k,\ n=1,2,\ldots\}$, $\beta^*=\{\beta^*_{in},i=1,\ldots,l,n=1,2,\ldots\}$, which will appear to be a tau function

$$\tau_{\tilde{\mathbf{r}},\mathbf{r}}(\lambda_{l-1},\lambda^*_k,\beta,\beta^*)=\langle I_k(\mathbf{p},\tilde{\mathbf{r}}_{k-1},\beta,0),I_l(\mathbf{p},\mathbf{r}_{l-1},\beta^*,0)\rangle_{r^{(k)},n_k} \tag{9.1.14}$$

where $\tilde{\mathbf{r}}=(\tilde{r}^{(1)},\tilde{r}^{(2)},\ldots,\tilde{r}^{(k-1)}),\mathbf{r}=(r^{(1)},r^{(2)},\ldots,r^{(k-1)},r^{(k)})$ and $\lambda_{l-1}=(n_1,\ldots,n_{l-1}),\lambda^*_k=(n^*_1,\ldots,n^*_k)$. We see that (9.1.14) is a sum over a pair of plane partitions.

9.2 Integral representation for $\tau_{\tilde{\mathbf{r}},\mathbf{r}}$

In case the function r has zero the scalar product (9.1.1) is degenerate one. For simplicity let us take $r(0)=0,r(k)\neq 0,k\leq n$. The scalar product (9.1.1) is non generate on the subspace of symmetric functions spanned by Schur functions $\{s_\lambda,l(\lambda)\leq n\}$, $l(\lambda)$ is the length of partition λ.

$$\langle f,g\rangle_{r,n}=\int\cdots\int\prod_{i=1}^n d\bar{z}_i dz_i\mu_r(\bar{z}_iz_i)\Delta_n(\bar{z})\Delta_n(z)f(\bar{z}^n)g(z^n) \tag{9.2.1}$$

where $\bar{z}^n=(\bar{z}_1,\ldots,\bar{z}_n),z^n=(z_1,\ldots,z_n)$ and

$$\Delta_n(\bar{z})=\prod_{i<j}^n(\bar{z}_i-\bar{z}_j),\quad \Delta_n(z)=\prod_{i<j}^n(z_i-z_j) \tag{9.2.2}$$

Thus get the following integral representation for (9.1.14)

$$\tau_{\tilde{\mathbf{r}},\mathbf{r}}(\lambda_{l-1},\lambda^*_k,\beta,\beta^*)= \tag{9.2.3}$$

$$\int\cdots\int\prod_{j=1}^{l-1}\prod_{i=1}^{n_j}d\bar{z}_{ji}dz_{ji}\mu_{\tilde{r}^{(j)}}(\bar{z}_{ji}z_{ji})D_j\delta(z_{ki}-z^*_{l-1,i})\delta(\bar{z}_{ki}-\bar{z}^*_{l-1,i})\prod_{j=1}^{k}\prod_{i=1}^{n^*_j}d\bar{z}^*_{ji}dz^*_{ji}\mu_{r^{(j)}}(\bar{z}^*_{ji}z^*_{ji})D^*_j \tag{9.2.4}$$

$$D_j=\Delta_{n_j}(\bar{z}^*_j)\Delta_{n_j}(z^*_j)e^{\xi_j},\quad D^*_j=\Delta_{n^*_j}(\bar{z}^*_j)\Delta_{n_j}(z^*_j)e^{\xi^*_j} \tag{9.2.5}$$

$$\xi_j=\sum_{m=1}^\infty\sum_{k=1}^{n_j}(z_{jk})^m\left(\beta_{jm}+\frac{1}{m}\sum_{i=1}^{n_{j-1}}(\bar{z}_{j-1,i})^m\right),j>1,\quad \xi_1=\sum_{m=1}^\infty\sum_{k=1}^{n_1}(z_{1k})^m\beta_{1m} \tag{9.2.6}$$

$$\xi^*_j=\sum_{m=1}^\infty\sum_{k=1}^{n^*_j}(\bar{z}^*_{1k})^m\left(\beta^*_{jm}+\frac{1}{m}\sum_{i=1}^{n^*_{j-1}}(z^*_{j-1,i})^m\right),j>1,\quad \xi^*_1=\sum_{m=1}^\infty\sum_{k=1}^{n_1}(\bar{z}^*_{1k})^m\beta^*_{1m} \tag{9.2.7}$$

9.3 Multi-matrix integrals

In case $\tilde{r}^{(j)}(n) = n + a_j, r^{(j)}(n) = n + a_j^*$ we get that the tau function (9.2.3) is equal to the following multi matrix integral

$$\int \cdots \int \prod_{j=1}^{l-1} dM_j d\bar{M}_j e^{TrM_j\bar{M}_j} e^{V_j} \delta(M_k - M_{l-1}^*)\delta(\bar{M}_{ki} - \bar{M}_{l-1,i}^*) \prod_{j=1}^{k} dM_j^* d\bar{M}_j^* e^{TrM_j^*\bar{M}_j^*} e^{V_j^*} \tag{9.3.1}$$

$$V_j = \sum_{m=1}^{\infty} Tr(M_j)^m \left(\beta_{jm} + \frac{1}{m} Tr(\bar{M}_{j-1})^m\right), j > 1, \quad V_1 = \sum_{m=1}^{\infty} Tr(M_1)^m \beta_{1m} \tag{9.3.2}$$

$$V_j^* = \sum_{m=1}^{\infty} Tr(\bar{M}_j^*)^m \left(\beta_{jm}^* + \frac{1}{m} Tr(M_{j-1}^*)^m\right), j > 1, \quad V_1^* = \sum_{m=1}^{\infty} Tr(\bar{M}_1^*)^m \beta_{1m}^* \tag{9.3.3}$$

where $M_j, \bar{M}_j$ has a size $n_j = M + a_j$ and $M_j^*, \bar{M}_j^*$ has a size $n_j^* = M + a_j^*$.

9.4 Fermionic representation [12]

Due to (3.3.10) scalar product (9.1.14) is equal to a vacuum expectation value. Let us write down it explicitly using [12].

Formula (6.3.1) is related to 'Gauss decomposition' of operators inside vacuums $\langle M| \ldots |M\rangle$ into diagonal operator $e^{H_0(T)}$ and upper triangular operator $e^{H(\mathbf{t})}$ and lower triangular operator $e^{-H^*(\mathbf{t}^*)}$ the last two have the Toeplitz form. Now let us consider more general two-dimensional Toda chain tau function

$$\tau = \langle M| e^{H(\mathbf{t})} g e^{-A(\mathbf{t}^*)} |M\rangle, \tag{9.4.1}$$

where we decompose g in the following way:

$$g(\beta, \beta^*) = e^{\tilde{A}_1(\beta_1)} \cdots e^{\tilde{A}_k(\beta_k)} e^{-A_l(\beta_l^*)} \cdots e^{-A_1(\beta_1^*)}, \tag{9.4.2}$$

where

$$\tilde{\beta}_i^* = (\beta_{i1}, \beta_{i2}, \ldots), \quad \beta_i^* = (\beta_{i1}^*, \beta_{i2}^*, \ldots) \tag{9.4.3}$$

$$\tilde{A}_k(\beta_k) = \sum_{m=1}^{\infty} \tilde{A}_{km}\beta_{km}, \quad A_k({\beta^*}_k) = \sum_{m=1}^{\infty} A_{km}{\beta^*}_{km} \tag{9.4.4}$$

Here each of $A_k(\beta_i^*)$ has a form as in (3.5.3) and corresponds to operator $r^{(k)}(D)$, while each of $\tilde{A}_k(\tilde{\beta^*}_k)$ has a form of (3.5.4) and corresponds to operator $\tilde{r}^{(k)}(D)$:

$$\tilde{A}_{km} = -\frac{1}{2\pi\sqrt{-1}} \oint \psi^*(z) \left(\tilde{r}^{(k)}(D) z\right)^m \psi(z), \quad m = 1, 2, \ldots \tag{9.4.5}$$

$$A_{km} = \frac{1}{2\pi\sqrt{-1}} \oint \psi^*(z) \left(\frac{1}{z} r^{(k)}(D)\right)^m \psi(z), \quad m = 1, 2, \ldots, \tag{9.4.6}$$

Collections of variables $\beta = \{\beta_{in}\}, \beta^* = \{\beta_{in}^*\}$ play the role of coordinates for some wide enough class of Clifford group elements g. This tau function is related to rather involved generalization of the hypergeometric functions we considered above. Tau function (9.4.1),(9.4.2) may be considered as the result of applying of the additional symmetries to the vacuum tau function, which is 1, see *Appendix "The vertex operator action"*.

Let us calculate this tau function. First of all we introduce a set consisting of $m + 1$ partitions:

$$(\lambda^1, \ldots, \lambda^m, \lambda^{m+1} = \lambda), \quad 0 = \lambda^0 \subseteq \lambda^1 \subseteq \lambda^2 \subseteq \cdots \subseteq \lambda^m \subseteq \lambda^{m+1} = \lambda, \tag{9.4.7}$$

see [4] for the notation $\subseteq$ for the partitions. The corresponding set

$$\Theta^m_\lambda = (\lambda^1, \theta^1, \ldots, \theta^m), \quad \theta^i = \lambda^{i+1} - \lambda^i, \quad i = 1, \ldots, m \tag{9.4.8}$$

depends on the partition λ and the number $m+1$ of the partitions. We take as $s_\Theta(\mathbf{t}^*, \beta^*)$ the product which is relevant to the set Θ^m_λ and depending on the set of variables $\mu_i = \{\mu_{ij}\}$ $(i = (1, \ldots, m+1),\ j = (1, 2 \ldots))$

$$s_{\Theta^m_\lambda}(\mu) = s_{\lambda^1}(\mu_1) s_{\theta^1}(\mu_2) \cdots s_{\theta^m}(\mu_{m+1}). \tag{9.4.9}$$

Here s_{θ^i} is a skew Schur function (see [4]). Further we define function $r_{\Theta^m_\lambda}(M)$:

$$r_{\Theta^m_\lambda}(M) = r_{\lambda^1}(M) r^1_{\theta^1}(M) \cdots r^m_{\theta^m}(M), \tag{9.4.10}$$

where the function $r^i_{\theta^i}(M)$, a skew analogy of $r_\lambda(M)$ of (2.2.1), is

$$r_{\theta^i}(M) = \prod_{j=1}^{s} r(n^{(i)}_j - j + 1 + M) \cdots r(n^{(i+1)}_j - j + M), \tag{9.4.11}$$

where $\lambda^{i+1} = (n^{(i+1)}_1, \ldots, n^{(i+1)}_s)$. If the function $r^i(m)$ has no poles and zeroes at integer points then the relation

$$r^i_{\theta^i}(M) = \frac{r^i_{\lambda^{i+1}}(M)}{r^i_{\lambda^i}(M)}, \quad i = 1, \ldots, m \tag{9.4.12}$$

is correct. To calculate the tau function we need the *Lemma*
Lemma 3 Let partitions $\lambda = (i_1, \ldots, i_s | j_1 - 1, \ldots, j_s - 1)$ and $\tilde\lambda = (\tilde i_1, \ldots, \tilde i_r | \tilde j_1 - 1, \ldots, \tilde j_r - 1)$ satisfy the relation $\lambda \supseteq \tilde\lambda$. The following is valid:

$$\begin{aligned} \langle 0 | \psi^*_{\tilde i_1} \cdots \psi^*_{\tilde i_r} \psi_{-\tilde j_r} \cdots \psi_{-\tilde j_1} e^{A^i(\beta_i)} \psi^*_{-j_1} \cdots \psi^*_{-j_s} \psi_{i_s} \cdots \psi_{i_1} | 0 \rangle = \\ = (-1)^{\tilde j_1 + \cdots + \tilde j_r + j_1 + \cdots + j_s} s_\theta(\beta_i) r_\theta(0), \qquad \theta = \lambda - \tilde\lambda. \end{aligned} \tag{9.4.13}$$

$$\begin{aligned} \langle 0 | \psi^*_{i_1} \cdots \psi^*_{i_s} \psi_{-j_s} \cdots \psi_{-j_1} e^{\tilde A^i(\beta^*_i)} \psi^*_{-\tilde j_1} \cdots \psi^*_{-\tilde j_r} \psi_{\tilde i_r} \cdots \psi_{\tilde i_1} | 0 \rangle = \\ = (-1)^{\tilde j_1 + \cdots + \tilde j_r + j_1 + \cdots + j_s} s_\theta(\beta_i) r_\theta(0), \qquad \theta = \lambda - \tilde\lambda. \end{aligned} \tag{9.4.14}$$

□ **Proof:** One proof is achieved by a development of , $e^A = 1 + A + \cdots$, $e^{\tilde A} = 1 + \tilde A + \cdots$ and a direct evaluation of vacuum expectations (9.4.13),(9.4.14). (see Example 22 in Sec 5 of [4] for help).
The second proof of (9.4.13),(9.4.14) is achieved using the developments (3.5.17), (3.5.18) respectively.□

Then we obtain the generalization of *Proposition 1*:

Proposition 18

$$\tau_M(\mathbf{t}, \mathbf{t}^*; \beta, \beta^*) = \sum_\lambda \sum_{\Theta^k_\lambda} \sum_{\Theta^l_\lambda} \tilde r_{\Theta^k_\lambda}(M) r_{\Theta^l_\lambda}(M) s_{\Theta^k_\lambda}(\mathbf{t}, \beta) s_{\Theta^l_\lambda}(\mathbf{t}^*, \beta^*), \tag{9.4.15}$$

where $\tilde r_{\Theta^k_\lambda}(M)$ and $r_{\Theta^l_\lambda}(M)$ are given by (9.4.11).

With the help of this series one can obtain different hypergeometric functions.

In the end of the subsection we put $\beta = 0$. For the case $\mathbf{t} = \mathbf{t}(\mathbf{x}^N)$, all $\beta = 0$ one can obtain the analog of (3.4.2)

$$e^{A(\mathbf{t}^*)}g^{-1}(\beta = 0, \beta^*)\psi(z)g(\beta = 0, \beta^*)e^{-A(\mathbf{t}^*)} = e^{-\xi_{r_l}(\beta^*_l, z^{-1})} \cdots e^{-\xi_{r_1}(\beta^*_1, z^{-1})}e^{-\xi_r(\mathbf{t}^*, z^{-1})}\psi(z) \tag{9.4.16}$$

$$e^{A(\mathbf{t}^*)}g^{-1}(\beta = 0, \beta^*)\psi^*(z)g(\beta = 0, \beta^*)e^{-A(\mathbf{t}^*)} = e^{\xi_{r'_l}(\beta^*_l, z^{-1})} \cdots e^{\xi_{r'_1}(\beta^*_1, z^{-1})}e^{\xi_{r'}(\mathbf{t}^*, z^{-1})}\psi^*(z) \tag{9.4.17}$$

where

$$\xi_{r_k}(\mathbf{t}^*, z^{-1}) = \sum_{m=1}^{+\infty} t_m \left(\frac{1}{z}r_k(D)\right)^m, \quad D = z\frac{d}{dz}, \quad {r'}_k(D) = r_k(-D) \tag{9.4.18}$$

In (9.4.16),(9.4.17) ξ_r are operators which act on z variable of ψ, ψ^*.

In cases $\mathbf{t} = \mathbf{t}(\mathbf{x}^N)$ (using (6.2.9) and (3.4.3)) and $\mathbf{t} = -\mathbf{t}(\mathbf{x}^N)$ (using (6.2.10) and (3.4.2)) one gets the following representations

Proposition 19

$$\tau(M, \mathbf{t}(\mathbf{x}^N), \mathbf{t}^*; \beta = 0, \beta^*) = \Delta^{-1}e^{{\eta_l}'} \cdots e^{{\eta_1}'}e^{\eta'}\Delta, \quad \Delta = \frac{\prod_{i<j}(x_i - x_j)}{(x_1 \cdots x_N)^{N-1-M}} \tag{9.4.19}$$

$$\tau(M, -\mathbf{t}(\mathbf{x}^N), \mathbf{t}^*; \beta = 0, \beta^*) = \Delta^{-1}e^{-\eta_l} \cdots e^{-\eta_1}e^{-\eta}\Delta, \quad \Delta = \frac{\prod_{i<j}(x_i - x_j)}{(x_1 \cdots x_N)^{N-1+M}} \tag{9.4.20}$$

where

$$\eta = \sum_{i=1}^{N} \xi_r(\mathbf{t}^*, x_i), \quad \eta_k = \sum_{i=1}^{N} \xi_{r_k}(\mathbf{t}^*, x_i) \tag{9.4.21}$$

$$\eta' = \sum_{i=1}^{N} \xi_{r'}(\mathbf{t}^*, x_i), \quad {\eta_k}' = \sum_{i=1}^{N} \xi_{{r'}_k}(\mathbf{t}^*, x_i) \tag{9.4.22}$$

and (cf (3.4.4))

$$\xi_{{r'}_k}(\mathbf{t}^*, x_i) = \sum_{m=1}^{+\infty} t_m \left(x_i r_k(D_i)\right)^m, \quad \xi_{r_k}(\mathbf{t}^*, x_i) = \sum_{m=1}^{+\infty} t_m \left(x_i r_k(-D_i)\right)^m, \quad D_i = x_i\frac{\partial}{\partial x_i} \tag{9.4.23}$$

Let us note that $[\xi_{{r'}_k}(\mathbf{t}^*, x_i), \xi_{{r'}_k}(\mathbf{t}^*, x_j)] = 0$ and $[\xi_{r_k}(\mathbf{t}^*, x_i), \xi_{r_k}(\mathbf{t}^*, x_j)] = 0$ for all k, i, j, while in general $[\eta_k, \eta_n] \neq 0$.

(9.4.20) may be also obtained from (9.4.19) with the help of (3.4.10).

Looking at (9.4.19) one easily derives a set of linear equations, which are differential equations with respect to variables β^* and $\mathbf{t}^*$:

$$\left(\frac{\partial}{\partial \beta^*_{km}} - e^{{\eta_l}'} \cdots e^{{\eta_{k-1}}'} \cdot \left(\sum_{i=1}^{N} x_i r_k(D_i)\right)^m \cdot e^{-{\eta_{k-1}}'} \cdots e^{-{\eta_l}'}\right)\left(\Delta\tau(M, \mathbf{t}(\mathbf{x}^N), \mathbf{t}^*; \beta = 0, \beta^*)\right) = 0 \tag{9.4.24}$$

$$\left(\frac{\partial}{\partial t^*_m} - e^{{\eta_l}'} \cdots e^{{\eta_1}'} \cdot \left(\sum_{i=1}^{N} x_i r_k(D_i)\right)^m \cdot e^{-{\eta_1}'} \cdots e^{-{\eta_l}'}\right)\left(\Delta\tau(M, \mathbf{t}(\mathbf{x}^N), \mathbf{t}^*; \beta = 0, \beta^*)\right) = 0 \tag{9.4.25}$$

where Δ is the same as in (9.4.19).

Proposition 20 *For the tau function $\tau_r(M, \mathbf{t}, \mathbf{t}^*)$ of (3.4.6) and the vertex operators $\Omega^{(\infty)}_{\tilde{r}1}, \Omega^{(0)}_{rl}$ defined by (6.6.1) we have*

$$e^{-\Omega^{(\infty)}_{\tilde{r}1}(\beta_1)} \cdots e^{-\Omega^{(\infty)}_{\tilde{r}k}(\beta_k)} \cdots e^{\Omega^{(\infty)}_{rl}(\beta^*_l)} \cdots e^{\Omega^{(\infty)}_{r1}(\beta^*_1)} \cdot \tau_r(M, \mathbf{t}, \mathbf{t}^*) = \tau(M, \mathbf{t}, \mathbf{t}^*; \beta, \beta^*), \tag{9.4.26}$$

$$e^{\Omega^{(0)}_{r1}(\beta^*_1)} \cdots e^{\Omega^{(0)}_{rl}(\beta^*_l)} \cdots e^{-\Omega^{(0)}_{\tilde{r}k}(\beta_k)} \cdots e^{-\Omega^{(0)}_{\tilde{r}1}(\beta_1)} \cdot \tau_r(M, \mathbf{t}, \mathbf{t}^*) = \tau(M, \mathbf{t}, \mathbf{t}^*; \beta, \beta^*). \tag{9.4.27}$$

9.5 Loop matrix integrals

Let us consider three matrix integral used in [61]

$$I = \int e^{\sum_{m=1}^{\infty}(\beta_m \mathrm{Tr} X_-^m + \beta_m^* \mathrm{Tr} X_+^m)} e^{\sqrt{-1}\mathrm{Tr} X_- X_+} e^{-q\sqrt{-1}\mathrm{Tr} X_- \Omega X_+ \Omega^{-1}} dX_- dX_+ d\Omega \tag{9.5.1}$$

Here $X_\pm$ are Hermitian n by n matrices, and $\Omega \in U(n)$. (In (9.5.1) we use β_m (and β_m^*) instead of t_m (and of t_{-m}) in [61]).

Using (2.2.11),(5.3.8) and (9.1.5) we get

$$I = \sum_{\lambda,\nu,\mu} s_\lambda(\beta) \frac{< s_\lambda(X_-) s_\nu(X_-), s_\nu(X_+) s_\mu(X_+) >_{r,n}}{(-q)^{-|\nu|}(n)_\nu} s_\mu(\beta^*) = \sum_{\lambda \geq \nu} \frac{(n)_\lambda s_{\lambda/\nu}(\beta) s_{\lambda/\nu}(\beta^*)}{(n)_\nu (-q)^{-|\nu|}} \tag{9.5.2}$$

As we see $I =<< \tau >>$ (see (3.6.8)), where

$$\tau(n, \mathbf{t}, \mathbf{t}^*; \beta, \beta^*) = \langle n | e^{H(\mathbf{t})} g e^{H^*(\mathbf{t}^*)} | n \rangle = \sum_{\lambda \geq \nu,\mu} \frac{(n)_\lambda s_{\lambda/\nu}(\beta) s_{\lambda/\mu}(\beta^*)}{(n)_\mu} s_\nu(\mathbf{t}) s_\mu(\mathbf{t}^*) (-q)^{|\mu|} \tag{9.5.3}$$

where

$$g = (-q)^n e^{H(\beta)} e^{A(\beta^*)} e^{-\log(-q)\sum_{m=-\infty}^{\infty} m :\psi_m \psi_m:} \tag{9.5.4}$$

where A is the same as in the tau function for the two-matrix model ($r(m) = m$). We have $I = \mathrm{Tr}_{F_n} g = \sum_\lambda g_{\lambda\lambda}$.

9.6 The example of Gelfand,Graev and Retakh hypergeometric series [12]

Below we also put $\beta = 0$. Let us consider the tau function:

$$\tau(M, \tilde{\beta}, \beta; 0, \beta^*) = \langle M | e^{\tilde{A}(\tilde{\beta})} e^{-A_l(\beta_l^*)} \cdots e^{-A_1(\beta_1^*)} e^{-A(\beta)} | M \rangle. \tag{9.6.1}$$

We put

$$\tilde{\beta} = (x, \frac{x^2}{2}, \frac{x^3}{3}, \ldots), \quad \beta = (y_1, 0, 0, \ldots), \quad \beta_i^* = (y_{i+1}, 0, 0, \ldots) \quad i = (1, \ldots, l). \tag{9.6.2}$$

We obtain the series

$$\tau(M, x, y_1, \ldots, y_{l+1}) = \sum_{n_1,\ldots,n_{l+1}=0}^{+\infty} \tilde{r}_{(n_1+\cdots+n_{l+1})}(M) r_{\Theta_\lambda^l}(M) \frac{(xy_1)^{n_1} \cdots (xy_{l+1})^{n_{l+1}}}{n_1! \cdots n_{l+1}!} = \tag{9.6.3}$$

$$\sum_{n_1,\ldots,n_{l+1} \in Z} c(n_1, \ldots, n_{l+1}) (xy_1)^{n_1} \cdots (xy_{l+1})^{n_{l+1}}, c(n_1, \ldots, n_{l+1}) = \frac{\tilde{r}_{(n_1+\cdots+n_{l+1})}(M) r_{\Theta_\lambda^l}(M)}{\Gamma(n_1+1) \cdots \Gamma(n_{l+1}+1)}, \tag{9.6.4}$$

where Θ_λ^l corresponds to the set of simple partitions-rows

$$\lambda^1 = (n_1), \lambda^2 = (n_1 + n_2), \ldots, \lambda_{l+1} = (n_1 + \cdots + n_{l+1}) \tag{9.6.5}$$

When functions $b_i(n_1, \ldots, n_{l+1})$ defined as

$$b_i(n_1, \ldots, n_{l+1}) = \frac{c(n_1, \ldots, n_i + 1, \ldots, n_{l+1})}{c(n_1, \ldots, n_{l+1})}, \quad i = 1, \ldots, l+1 \tag{9.6.6}$$

are rational functions of $(n_1, \ldots, n_{l+1})$, then tau function (9.6.3) is a Horn hypergeometric series [25].

Above series for the special choice of functions $r^i(D)$ can be deduced from the Gelfand, Graev and Retakh series [54] defined on the special lattice and corresponding to the special set of parameters. Let us take the rational functions $r^i(D)$:

$$r^i(D) = \frac{\prod_{j=1}^{p^{(i)}}(D + a_j^{(i)})}{\prod_{m=1}^{s^{(i)}}(D + b_m^{(i)})}, \quad (i = 0, \ldots, l), \qquad r^0(D) = r(D) \tag{9.6.7}$$

$$\tilde{r}(D) = \frac{\prod_{j=1}^{p^{(l+1)}}(D + a_j^{(l+1)})}{\prod_{m=1}^{s^{(l+1)}}(D + b_m^{(l+1)})} \tag{9.6.8}$$

Let define $N = p^{(0)} + s^{(0)} + 2\sum_{j=1}^{l}(p^{(j)} + 2s^{(j)}) + p^{(l+1)} + s^{(l+1)} + l + 1$ and consider complex space C^N. In this space we consider the $l+1$-dimensional basis B and the vector υ consisting of parameters.

$$\begin{gathered} p_0 = s_0 = 0, \quad p_i = p^{(i-1)} + s^{(i)}, \quad s_i = s^{(i-1)} + p^{(i)}, \quad i = (1, \ldots, l) \\ p_{l+1} = p^{(l)} + p^{(l+1)}, \quad s_{l+1} = s^{(l)} + s^{(l+1)}, \quad N = \sum_{j=1}^{l+1}(p_j + s_j) + l + 1 \end{gathered} \tag{9.6.9}$$

$$\begin{gathered} \mathbf{f}^i = -(\mathbf{e}_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+1} + \cdots + \mathbf{e}_{p_1+s_1+\cdots+p_{i-1}+s_{i-1}+p_i}) + \\ +(\mathbf{e}_{p_1+s_1+\cdots+p_{i-1}+s_{i-1}+p_i+1} + \cdots + \mathbf{e}_{p_1+s_1+\cdots+p_{i-1}+s_{i-1}+p_i+s_i}), \quad i = 1, \ldots, l+1 \end{gathered} \tag{9.6.10}$$

where $\mathbf{e}_i = \underbrace{(0, \ldots, 0, \overset{i}{\hat{1}}, 0, \ldots)}_{\text{N}}$. The lattice $B \in C^N$ is generated by the vector basis of dimension $l+1$:

$$\mathbf{b}^i = \mathbf{f}^i + \cdots + \mathbf{f}^{l+1} + \mathbf{e}_{N-l-1+i}, \qquad i = 1, \ldots, l+1 \tag{9.6.11}$$

Vector $\upsilon \in C^N$ is defined as follows (compare with (9.6.10)):

$$\begin{aligned} \upsilon^i = -(&a_1^{(i-1)}\mathbf{e}_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+1} + \cdots + a_{p^{(i-1)}}^{(i-1)}\mathbf{e}_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p^{(i-1)}} + \\ &+b_1^{(i)}\mathbf{e}_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p^{(i-1)}+1} + \cdots + b_{s^{(i)}}^{(i)}\mathbf{e}_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p_i}) + \\ +((&b_1^{(i-1)} - 1)\mathbf{e}_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p_i+1} + \cdots + (b_{s^{(i-1)}}^{(i-1)} - 1)\mathbf{e}_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p_i+s^{(i-1)}} + \\ &+(a_1^{(i)} - 1)\mathbf{e}_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p_i+s^{(i-1)}+1} + \cdots + (a_{s^{(i)}}^{(i)} - 1)\mathbf{e}_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p_i+s_i}) \end{aligned} \tag{9.6.12}$$

for $i = (1, \ldots l)$, and

$$\begin{aligned} \upsilon^{l+1} = -(&a_1^{(l)}\mathbf{e}_{p_0+s_0+\cdots+p_l+s_l+1} + \cdots + a_{p^{(l)}}^{(l)}\mathbf{e}_{p_0+s_0+\cdots+p_l+s_l+p^{(l)}} + \\ &+a_1^{(l+1)}\mathbf{e}_{p_0+s_0+\cdots+p_l+s_l+p^{(l)}+1} + \cdots + a_{s^{(l+1)}}^{(l+1)}\mathbf{e}_{p_0+s_0+\cdots+p_l+s_l+p_{l+1}}) + \\ +((&b_1^{(l)} - 1)\mathbf{e}_{p_0+s_0+\cdots+p_l+s_l+p_{l+1}+1} + \cdots + (b_{s^{(l)}}^{(l)} - 1)\mathbf{e}_{p_0+s_0+\cdots+p_l+s_l+p_{l+1}+s^{(l)}} + \\ &+(b_1^{(l+1)} - 1)\mathbf{e}_{p_0+s_0+\cdots+p_l+s_l+p_{l+1}+s^{(l)}+1} + \cdots + (b_{s^{(l+1)}}^{(l+1)} - 1)\mathbf{e}_{p_0+s_0+\cdots+p_l+s_l+p_{l+1}+s_{l+1}}) \end{aligned} \tag{9.6.13}$$

Vector υ is:

$$\upsilon = \upsilon^1 + \cdots + \upsilon^{l+1} \tag{9.6.14}$$

Now we can write down Gelfand, Graev and Retakh hypergeometric series corresponding to the lattice B and vector υ:

$$F_B(\upsilon;z) = \sum_{\mathbf{b}\in B}\prod_{j=1}^{N}\frac{z_j^{\upsilon_j+b_j}}{\Gamma(\upsilon_j+b_j+1)}+$$

$$\sum_{n_1,\dots,n_{l+1}\in Z}\prod_{j=1}^{N}\frac{z_j^{\upsilon_j+n_1b_j^1+\cdots+n_{l+1}b_j^{l+1}}}{\Gamma(\upsilon_j+n_1b_j^1+\cdots+n_{l+1}b_j^{l+1}+1)} \tag{9.6.15}$$

Let us compare this series with tau function (9.6.3):

$$F_B(\upsilon;\mathbf{z}) = c_1(a,b)g_1(\mathbf{z})\cdots c_{l+1}(a,b)g_{l+1}(\mathbf{z})\tau(M,x,y_1,\dots,y_{l+1}) \tag{9.6.16}$$

where

$$c_i^{-1}(a,b) = \Gamma(1-a_1^{(i-1)})\cdots\Gamma(1-a_{p^{(i-1)}}^{(i-1)})\Gamma(1-b_1^{(i)})\cdots\Gamma(1-b_{s^{(i)}}^{(i)})\times$$
$$\times\Gamma(b_1^{(i-1)})\cdots\Gamma(b_{s^{(i-1)}}^{(i-1)})\Gamma(a_1^{(i)})\cdots\Gamma(a_{s^{(i)}}^{(i)}),\quad i=1,\dots,l \tag{9.6.17}$$

$$c_{l+1}^{-1}(a,b) = \Gamma(1-a_1^{(l)})\cdots\Gamma(1-a_{p^{(l)}}^{(l)})\Gamma(1-a_1^{(l+1)})\cdots\Gamma(1-a_{s^{(l+1)}}^{(l+1)})\times$$
$$\times\Gamma(b_1^{(l)})\cdots\Gamma(b_{s^{(l)}}^{(l)})\Gamma(b_1^{(l+1)})\cdots\Gamma(b_{s^{(l+1)}}^{(l+1)}) \tag{9.6.18}$$

$$\frac{z_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p_i+1}\cdots z_{p_1+s_1+\cdots+p_{i-1}+s_{i-1}+p_i+s_i}}{(-z_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+1})\cdots(-z_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p_i})}=1,\quad i=2,\dots,l \tag{9.6.19}$$

$$\frac{z_{p_1+1}\cdots z_{p_1+s_1}z_{N-l}}{(-z_1)\cdots(-z_{p_1})}=y_1 \tag{9.6.20}$$

$$y_i = z_{N-l-1+i},\quad i=2,\dots,l+1 \tag{9.6.21}$$

$$\frac{z_{p_1+s_1+\cdots+p_l+s_l+p_{l+1}+1}\cdots z_{p_1+s_1+\cdots+p_l+s_l+p_{l+1}+s_{l+1}}}{(-z_{p_1+s_1+\cdots+p_l+s_l+1})\cdots(-z_{p_1+s_1+\cdots+p_l+s_l+p_{l+1}})}=x \tag{9.6.22}$$

$$g_i(\mathbf{z}) = z_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+1}^{\left(-a_1^{(i-1)}\right)}\cdots z_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p^{(i-1)}}^{\left(-a_{p^{(i-1)}}^{(i-1)}\right)}\times$$
$$\times z_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p^{(i-1)}+1}^{\left(-b_1^{(i)}\right)}\cdots z_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p_i}^{\left(-b_{s^{(i)}}^{(i)}\right)}\times$$
$$\times z_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p_i+1}^{\left(b_1^{(i-1)}-1\right)}\cdots z_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p_i+s^{(i-1)}}^{\left(b_{s^{(i-1)}}^{(i-1)}-1\right)}\times$$
$$\times z_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p_i+s^{(i-1)}+1}^{\left(a_1^{(i)}-1\right)}\cdots z_{p_0+s_0+\cdots+p_{i-1}+s_{i-1}+p_i+s_i-1}^{\left(a_{s^{(i)}}^{(i)}-1\right)} \tag{9.6.23}$$

for $i=(1,\dots l)$, and

$$g_{l+1}(\mathbf{z}) = z_{p_1+s_1+\cdots+p_l+s_l+1}^{\left(--a_1^{(l)}\right)}\cdots z_{p_1+s_1+\cdots+p_l+s_l+p^{(l)}}^{\left(-a_{p^{(l)}}^{(l)}\right)}\times$$
$$\times z_{p_1+s_1+\cdots+p_l+s_l+p^{(l)}+1}^{\left(-a_1^{(l+1)}\right)}\cdots z_{p_1+s_1+\cdots+p_l+s_l+p_{l+1}}^{\left(-a_{s^{(l+1)}}^{(l+1)}\right)}\times$$
$$\times z_{p_1+s_1+\cdots+p_l+s_l+p_{l+1}+1}^{\left(b_1^{(l)}-1\right)}\cdots z_{p_1+s_1+\cdots+p_l+s_l+p_{l+1}+s^{(l)}}^{\left(b_{s^{(l)}}^{(l)}-1\right)}\times$$
$$\times z_{p_1+s_1+\cdots+p_l+s_l+p_{l+1}+s^{(l)}+1}^{\left(b_1^{(l+1)}-1\right)}\cdots z_{p_1+s_1+\cdots+p_l+s_l+p_{l+1}+s_{l+1}-1}^{\left(b_{s^{(l+1)}}^{(l+1)}-1\right)} \tag{9.6.24}$$

10 Acknowledgements

It is my pleasure to thank John Harnad for helpful numerous discussions and Takahiro Shiota for helpful initial discussions. I thank S.Milne for sending [38], L.A.Dickey the discussion and for sending [53]. I thank M.Bertola, B.Eynard and J.Harnad for the discussion of the preprint [49]. I thank also Anton Zabrodin and Mark Mineev for the numerous discussions of the paper [43] and Aleksej Morozov for giving [69]. I thank grant RFBR 02-02-17382a. I thank the hospitality of the University of Montreal (CRM, October-November 2001) and of the Los Alamos National Laboratory (February 2002), I thank J.Harnad (CRM) and M.Mineev (Los Alamos) and M.Chertkov (Los Alamos) for the organizations of my visits there. I thank Yan Fyodorov for giving the reference [66].

References

[1] A. Yu. Orlov, "Soliton theory, Symmetric Functions and Matrix Integrals" SI/0207030 v1

[2] J. Harnad and A. Yu. Orlov "Matrix Integrals as Borel sums of Schur Function Expansions", nlin.SI/0209035; J. Harnad and A. Yu. Orlov "Schur Function Expansions of Matrix Integrals", preprint CRM December 2001

[3] A. Yu. Orlov, "New Solvable Matrix Integrals", nlin. SI/0209063

[4] I. G. Macdonald, Symmetric Functions and Hall Polynomials, Clarendon Press, Oxford, 1995

[5] M. L. Mehta, *Random Matrices* Academic Press, Inc, 1991

[6] V. E. Zakharov and A. B. Shabat, J. Funct. Anal. Appl. **8**, 226 (1974), **13**, 166 (1979)

[7] E. Date, M. Jimbo, M. Kashiwara and T. Miwa, "Transformation groups for soliton equations", in: Nonlinear integrable systems – classical theory and quantum theory eds. M. Jimbo, and T. Miwa, World Scientific, 39–120, 1983

[8] V. Kazakov, Phys Lett **150B**, 282 (1985)

[9] A. Yu. Orlov and M. D. Scherbin "Milne's hypergeometric functions in terms of free fermions" J.Phys A:Math.Gen.**34** (2001) 2295-2310

[10] A. Yu. Orlov and D. M. Scherbin "Multivariate hypergeometric functions as tau functions of Toda lattice and Kadomtsev-Petviashvili equation", Physics **D**, March 2001

[11] A. Yu. Orlov and D. M. Scherbin "Hypergeometric solutions of soliton equations" Teor Mat Phys **128** No 1 pp 84-108 (2001)

[12] A. Yu. Orlov and D. M. Scherbin " Fermionic representation for basic hypergeometric functions related to Schur polynomials" nlin.SI/0001001

[13] M. Adler, T. Shiota and P. van Moerbeke, Phys.Lett. A 194 (1994), no. 1-2, 33-34

[14] M. Adler and P. van Moerbeke, "Integrals over Grassmannians and Random permutations" math.CO/0110281

[15] P. van Moerbeke, "Integrable Lattices: Random Matrices and Random Purmutations" CO/0010135

[16] Harish Chandra, A.J.Math **80** 241 (1958); C.Itzykson and J.B.Zuber, J.Math.Phys. **21** 411 (1980)

[17] V. V. Kadomtsev and V. I. Petviashvili, Soviet Phys. Dokl. **15**, 539 (1970)

[18] V. E. Zakharov, S. V. Manakov, S. P. Novikov and L. P. Pitaevsky *The Theory of Solitons. The Inverse Scattering Method*

[19] A. Yu. Morozov, "Integrability and Matrix Models" Uspehi Fizicheskih Nauk, vol 164 (1994) pp.3-62

[20] L. A. Dickey, *Soliton Equations and Hamiltonian System*, World Scientific, Singapore, 1991

[21] V. Dryuma, JETF Lett.,19, 219 (1974)

[22] A. V. Mikhailov, "On the Integrability of two-dimensional Generalization of the Toda Lattice", Letters in Journal of Experimental and Theoretical Physics, v.30, p. 443-448, 1979

[23] M. Jimbo and T. Miwa, Publ. RIMS Kyoto Univ. **19**, 943 (1983)

[24] T. Miwa, "On Hirota's difference equations", Proc. Japan Acad. **58** Ser.A (1982) 9-12

[25] N. Ya. Vilenkin and A. U. Klimyk, Representation of Lie groups and Special Functions, Recent Advances, Kluwer Academic Publishers, 1995

[26] N. Ya. Vilenkin and A. U. Klimyk *Representation of Lie Groups and Special Functions. Volume 3: Classical and Quantum Groups and Special Functions*, Kluwer Academic Publishers, 1992

[27] K. Ueno and K. Takasaki, Adv. Stud. Pure Math. **4**, 1-95 (1984)

[28] K. Takasaki, "Initial value problem for the Toda lattice hierarchy", Adv. Stud. Pure Math. **4**, pp. 139-163 (1984)

[29] T. Takebe, "Representation Theoretical Meaning of Initial Value Problem for the Toda Lattice Hierarchy I", LMP **21**: 77-84, 1991

[30] T. Takebe, "Representation Theoretical Meaning of Initial Value Problem for the Toda Lattice Hierarchy II", Publ. RIMS, Kyoto Univ. **27** (1991), 491-503

[31] T. Takebe and K. Takasaki, "Integrable Hierarchies and Dispersionless Limit", Rev. Math. Phys. 7 (1995) 743-808; hep-th/94050096

[32] T. Nakatsu, K. Takasaki and S. Tsujimaru, "Quantum and Classical Aspects of Deformed $c = 1$ Strings", Nucl. Phys. B443 (1995) 155; hep-th/9501038, January 1995

[33] K. Takasaki, "The Toda Lattice Hierarchy and Generalized String Equation", Comm.Math.Phys. 181 (1996) 131; hep-th/9506089, June 1995

[34] L. A. Dickey, Modern Physics Letters A, Vol. 8, No 14, 1357-1377 (1993)

[35] A. Yu. Orlov and P. Winternitz, Theoretical and Mathematical Physics, Vol 113, No2, pp.1393-1417 (1997)

[36] A. Yu. Orlov, "Vertex Operators, ∂ bar Problem, Symmetries, Hamiltonian and Lagrangian Formalism of (2+1) Dimensional Integrable Systems", in *Plasma Theory and Nonlinear and Turbulent Processes in Physics, Proc. III Kiev. Intern. Workshop (1987)*, vol. I, pp. 116–134, World Scientific Pub, Singapore, 1988, (editors: V. G. Bar'yakhtar,...,V. E. Zakharov)

[37] P. G. Grinevich and A. Yu. Orlov, "Virasoro Action on Riemann Surfaces, Grassmannians, $det\bar{\partial}_j$ and Segal-Wilson tau function", in *Problems of Modern Quantum Field Theory* pp. 86-106, Springer, 1989 (editors: A. A. Belavin, A. U. Klimyk, A.B.Zamolodchikov)

[38] S. C. Milne, "Summation theorems for basic hypergeometric series of Schur function argument", in *Progress in Approximation Theory*, Eds. A. A. Gonchar and E. B. Saff, pp. 51-77, Springer-Verlag, New-York, 1992

[39] K. I. Gross and D. S. Richards "Special functions of matrix arguments.I: Algebraic induction, zonal polynomials, and hypergeometric functions", Transactions Amer Math Soc, **301**, No 2 (1987) 781-811

[40] A. P. Prudnikov, Yu. A. Brychkov and O. I. Marichev "Integrals and series. Additional chapters", Moscow, Nauka 1986 (in Russian)

[41] V. Kac and A. Radul, "Representation theory of the vertex algebra $W_{1+\infty}$" hep-th/9512150

[42] B. Konopelchenko and L. M. Alonso, "The KP Hierarchy in Miwa coordinates", solv-int/9905005

[43] M. Mineev-Weinstein, P. Wiegmann and A. Zabrodin, "Integrable structure of interface dynamics", LAUR-99-0703

[44] A.Gerasimov, A. Marshakov, A. Mironov, A. Morozov and A. Orlov, Nuclear Physics **B357** (1991) 565-618

[45] A. Zabrodin, S. Kharchev, A. Mironov, A. Marshakov and A. Orlov "Matrix Models among Integrable Theories: Forced Hierarchies and Operator Formalizm" Nuclear Physics B366 (1991) 569-601

[46] G. Segal and G. Wilson, Publ. Math. IHES **61**, 5 (1985)

[47] P. Zinn-Justion "HCIZ integral and 2D Toda lattice hierarchy" math-ph/0202045

[48] P. Zinn-Justion and J.-B.Zuber, "On some integrals over the U(N) unitary group and their large limit", math-ph/0209019

[49] M. Bertola, B. Eynard and J. Harnad, "Partition Functions for Matrix Models ans Izomonodromic Tau Functions" preprint SI/0204054

[50] B. Eynard "Random Matrices"Cours de Physique Theorique de Sacley" Sacley-T01/014 CRM-2708

[51] O. Kravchenko and B. Khesin, Funct. Anal. Appl., **25**, 83 (1991)

[52] A. Yu. Orlov "BKP hierarchy and hypergeometric tau functions"

[53] L. A. Dickey, “Joint solution to the string equation and equations of the n-reduced KP hierarchy (Kontsevich integral)”, Preprint, 1992; L. A. Dickey, ”Chains of KP, semi-infinite 1-Toda lattice hierarchy and Kontsevich integral”, Journal of Applied Mathematics 1:4 (2001) 175-193

[54] I. M. Gelfand, M. I. Graev and V. S. Retakh, *General hypergeometrical systems of equations and series of hypergeometrical kind*, UMN, vol.47, 4, 1992

[55] V. Kazakov, ”Solvable matrix models” hep-th/003064

[56] V. Kazakov, M. Staudacher and T.Wynter ”Character Expension Method for Matrix Models of Dually Weighted Graphs” hep-th/9502132 v 2 (1995)

[57] Ivan K. Kostov, M. Staudacher and T. Wynter, ”Complex Matrix Models and Statistics of Branched Covering of 2D Surfaces” heph-th/9703189 March 1997

[58] Ivan K. Kostov, ”Exact Solution of the Six-Vertex Model on a Random Lattice” hep/9911023

[59] Ivan K. Kostov, ”$U(N)$ Gauge Theory and Lattice Strings” hep-th/9306110

[60] Ivan K. Kostov ”Conformal Field Theory Techniques in Random Matrix models” hep-th/9907060

[61] Ivan K. Kostov, ”Integrable Flows in $c = 1$ String Theory” hep-th/0208034

[62] R. Dijgraph and C. Vafa, ”On Geometry and Matrix Models” hep-th/0207106

[63] E. Brezin and S. Hikami, ”Characteristic polynomials of random matrices” math-ph/9910005

[64] E.Brezin, S.Hikami, ”An extension of the Harish-Chandra-Itzykson-Zubet integral” math-ph/0208002

[65] Y. V. Fyodorov and E. Strahov, ”Characteristic polynomials of random Hermitian matrices and Duistermaat-Heckman localization on noncompact Kaehler manifolds”, math-ph/0201045

[66] Y. V. Fyodorov and E. Strahov, ”An exact formula for general spectral correlation function of random Hermitian matrices” math-ph/0204051

[67] M. Feigin and A. P. Veselov, ”Quasiinvariants of Coxeter groups and m-harmonic polynomials” math-ph/0105014

[68] D. J. Gross and E. Witten, ”Possible Third Order Phase Tranzition in the Large N Lattice Gauge Theory” Phys Rev D **21**, 446 (1980)

[69] A. Mironov, A. Morozov and G. Semenoff ”Unitary Matrix Integrals in the Framework of the Generalized Kontsevich Model” Intern J Mod Phys A, Vol 11 No 28 (1996) 5031-5080

[70] L-L. Chau and O. Zaboronsky ”On the Structure of Correlation Functions in the Normal Matrix Models” hep-th/9711091 (1997)

[71] I. M. Gessel, ”Symmetric Functions and P-recursiveness” J. Comb. Theory Ser A **53** (1990) 257-285